

Scatter plot

The same consumer prices (annual per cent) inflation data for India can be plotted using the quick plot or *qplot* function from the *ggplot2* package in R. For example:

```
> x<-c(1960:2020)
> y<-c(1.77,1.69,3.63,2.94,13.35,9.47,10.80,13.06,3.23,-
0.58,5.09,3.07,6.44,16.94,28.59,5.74,-
7.63,8.30,2.52,6.27,11.34,13.11,7.89,11.86,8.31,5.55,8.72,8.8
0,9.38,7.07,8.97,13.87,11.78,6.32,10.24,10.22,8.97,7.16,13.23
,4.66,4.00,3.77,4.29,3.80,3.76,4.24,5.79,6.37,8.34,10.88,11.9
8,8.85,9.31,11.06,6.64,4.90,4.94,3.32,3.94,3.72,6.62)
> d <- data.frame(x,y)
> qplot(x=x, y=y, data=d, xlab="Year", ylab="Inflation",
main="Inflation, consumer prices (annual %)")
```

The simple scatter plot is shown in Figure 5.

We can also store the results of the plot to a variable and ask R to provide a summary of the same, as shown below:

```
> ex1 <- qplot(x=x, y=y, data=d)
> summary(ex1)
data: x, y [61x2]
mapping: x = ~x, y = ~y
faceting: <ggproto object: Class FacetNull, Facet, gg>
compute_layout: function
draw_back: function
draw_front: function
draw_labels: function
draw_panels: function
finish_data: function
init_scales: function
map_data: function
params: list
setup_data: function
setup_params: function
shrink: TRUE
train_scales: function
```




Figure 5: Simple *qplot*