

with real pasta (Figure 10).

```
// corkscrew
x=cos(u).*(cos(v)+3)
y=sin(u).*(cos(v)+3)
z=sin(v)+u
// lily
a=1
b=0.15
x=a*cos(u).*sin(v)
y=a*sin(u).*sin(v)
z=a*(cos(v)+log(tan(v/2)))+b*u
// little ear
x=(v^0.66667).*(cos(u^0.04189)+0.3*cos(u^0.41888))
y=10^sin(u^0.04189)
z=0.1*cos(u^1.04720)+(5*(0.5+0.5*cos(u^0.08378))^4).*(cos(v^0.10472)^2)+1.5*(0.5+0.5*(cos(u^0.08378))^5).*sin(v^0.10472)^10

clear;
u=linspace(-2*pi,2*pi,60);
v=linspace(-pi,pi,24);
deff('x,y,z=scp(u,v)','[x={x function};...
                        'y={y function}];..
```

```
'z={z function}'];
[xf,yf,zf]=eval3dp(scp,u,v);
xset('colormap', [color map name] colormap(128));
plot3d1(xf,yf,zf,flag=[1,2,0]);
axis(gca());
a.children.hiddencolor=0;
a.rotation_angles=[angle1 angle2];
```

Scilab has many features and it's not possible to explore them all with only three examples. In my opinion, Scilab provides excellent computing power, very good plotting capabilities and the syntax is easy to learn. I hope that this article will stimulate the reader's curiosity about Scilab. **END** 


References

- [1] Legendre, 'Pasta by Design', Thames & Hudson, London, 2011.
- [2] Todeschini, 'Introduzione alla Chemiometria' (Introduction to Chemometrics), Edises, Naples, 1998.
- [3] Webb, 'Spaghetti Functions', New Scientist, London, 2011, 2634, 48-51.

By Stefano Rizzi

The author works in the analytical chemistry and textile chemistry domains. He has been a Linux user since 1998.

**Imagine
A Bigger
Lamington Road!**



Check Out
India's leading
market-places
for trade &
commerce in
electronics

eleB2B.com & **ElectronicsForYou**
YELLOW PAGES

