


THEROPODS (THERRO-pods)


All meat-eating dinosaurs were saurischians, and they formed a group called the theropods. This is the group from which birds are descended. They ranged in size from the chicken-sized *Compsognathus* to monsters such as the mighty *Spinosaurus*.



◀ SPINOSAURUS
had a massive skin
"sail" that ran the
length of its back.

SAUROPODOMORPHS (SORE-oh-POD-oh-morfs)

This group contained the heaviest and longest animals ever to walk on Earth. They were herbivores (plant-eaters) and would have had to graze constantly to obtain the energy they needed.



▲ BRACHIOSAURUS
Sauropods had tiny heads
compared to their bodies.

ORNITHOPODS (OR-nith-oh-pods)


These plant-eaters all roamed on two legs, so were able to use their forelimbs to grasp their food. They were immensely successful and very common. Fossils of these types of dinosaur have been found all over the world.



▲ IGUANODON, an
ornithomimid, was the second
dinosaur to be named.

MARGINOCEPHALIANS (MAR-jee-no-sa-FAY-lee-ans)

This group of plant-eaters had heads that sported bony frills. (Marginocephalian means "fringed head.") Some walked on two legs, some on four. They were common dinosaurs in the Cretaceous and included the well-known *Triceratops*.



▲ EINIOSAURUS
had a forward-
curving horn.

