

```

10. variance = sum(squared_deviations) / len(returns)
11. volatility = variance ** 0.5
12.
13. # Calculate maximum drawdown
14. cumulative_returns = [1]
15. for r in returns:
16.     cumulative_returns.append(cumulative_returns[-1] * (1 + r))
17.
18. peak = cumulative_returns[0]
19. max_drawdown = 0
20.
21. for value in cumulative_returns:
22.     if value > peak:
23.         peak = value
24.     drawdown = (peak - value) / peak
25.     if drawdown > max_drawdown:
26.         max_drawdown = drawdown
27.
28. return total_return, average_return, volatility, max_drawdown
29.
30. When calling this function, you can capture all returned values:
31. # Strategy returns (as decimal percentages)
32. strategy_returns = [0.01, -0.005, 0.02, 0.015, -0.01]
33.
34. # Get all metrics at once
35. total, average, vol, drawdown = calculate_strategy_metrics(strategy_returns)
36.
37. print(f"Total Return: {total:.2%}")
38. print(f"Average Return: {average:.2%}")
39. print(f"Volatility: {vol:.2%}")
40. print(f"Maximum Drawdown: {drawdown:.2%}")
41.

```

This ability to return multiple related values makes functions particularly powerful for comprehensive financial analysis.