

NEREUS: An AI-Driven Water Quality Intelligence System

A Comprehensive Analysis Using Satellite-Derived Indices, ML
Regression, and Generative AI

Authors: Hemang Seth & Satyam Devangan
IIIT Bangalore

Date: December 5, 2025

Contents

| | |
|---|-----------|
| Abstract | 2 |
| 1 Introduction | 3 |
| 2 Dataset Description | 4 |
| 2.1 Sample Data Table | 4 |
| 3 Correlation Analysis | 5 |
| 3.1 Correlation Heatmap | 5 |
| 4 Temporal Trend Analysis | 6 |
| 4.1 Chla Trend Over Time (Figure 2) | 6 |
| 4.2 Turbidity Trend Over Time (Figure 3) | 7 |
| 4.3 Shrinkage (NDWI) Trend Over Time (Figure 4) | 8 |
| 5 Distribution Analysis | 9 |
| 5.1 Chla Distribution by Year (Figure 5) | 9 |
| 5.2 Turbidity Distribution by Year (Figure 6) | 10 |
| 5.3 Shrinkage Distribution by Year (Figure 7) | 11 |
| 6 Machine Learning Model Evaluation | 12 |
| 6.1 Model Accuracy Comparison (Figure 11) | 12 |
| 7 Discussion | 13 |
| 8 Conclusion | 14 |

Abstract

This report presents **NEREUS**, an integrated AI-remote sensing water quality analytics system. Using Chla, Turbidity (NTU), and NDWI-derived Shrinkage metrics collected longitudinally from 2020–2024, the system performs ML modeling, trend decomposition, and predictive environmental assessment. A generative AI engine (Gemini 2.5) interprets dataset patterns and provides advanced ecological insights. This document outlines the data pipeline, methods, plots (10 total), and detailed analysis.

Chapter 1

Introduction

Monitoring inland water systems requires quantifying optical, biological, and hydrological properties using satellite indices. This project integrates:

- **Chla** — Chlorophyll-a and algal bloom proxy
- **Turbidity (NTU)** — Suspended matter concentration
- **NDWI / Shrinkage** — Water body area fluctuation

In combination with advanced ML (Random Forest, Gradient Boosting, etc.), the system delivers predictive analytics up to 90–93% accuracy.

Chapter 2

Dataset Description

The dataset contains daily observations from 2020–2024. Additional temporal features (Year, Month, DayOfYear) were engineered.

2.1 Sample Data Table

| Date | Chla | Turbidity (NTU) | Shrinkage (%) |
|------------|------|-----------------|---------------|
| 2020-08-01 | 4.20 | 9.0 | 1.83 |
| 2020-08-02 | 4.52 | 9.1 | 1.90 |
| 2020-08-03 | 4.55 | 10.0 | 2.10 |

Table 2.1: Sample rows from the dataset.

Chapter 3

Correlation Analysis

3.1 Correlation Heatmap




Figure 3.1: Correlation Matrix Across Chla, Turbidity, Shrinkage, and Temporal Features.

Analysis: Chla shows moderate positive correlation with Year (0.53), suggesting long-term gradual increase in chlorophyll content. Turbidity and shrinkage show weak pairwise correlations, indicating independent seasonal drivers.

Chapter 4

Temporal Trend Analysis

4.1 Chla Trend Over Time (Figure 2)




Figure 4.1: Chla Trend (2020–2024).

Analysis: Chla values show gradual upward drift, indicating progressive chlorophyll intensification and potential eutrophication risk.

4.2 Turbidity Trend Over Time (Figure 3)




Figure 4.2: Turbidity Trend (2020–2024).

Analysis: Turbidity variability is wider, suggesting rainfall, sediment transport, and wind-driven resuspension events.

4.3 Shrinkage (NDWI) Trend Over Time (Figure 4)




Figure 4.3: NDWI-Based Shrinkage Trend (2020–2024).

Analysis: Seasonal shrinkage patterns align with monsoon/non-monsoon transitions.

Chapter 5

Distribution Analysis

5.1 Chla Distribution by Year (Figure 5)




Figure 5.1: Chla Distribution (2020–2024).

Analysis: Distributions gradually shift upward with narrower variance in later years.

5.2 Turbidity Distribution by Year (Figure 6)




Figure 5.2: Turbidity Distribution (2020–2024).

5.3 Shrinkage Distribution by Year (Figure 7)




Figure 5.3: Shrinkage Distribution (2020–2024).

Analysis: Violin shapes reveal seasonal clustering and water-level seasonality.

Chapter 6

Machine Learning Model Evaluation

6.1 Model Accuracy Comparison (Figure 11)




Figure 6.1: Comparison of ML model accuracies: Random Forest, Gradient Boosting, AdaBoost, Linear Regression, SVR, KNN.

Analysis: Random Forest achieves the highest accuracy (92–93%), making it the best-performing model for Chla prediction.

Chapter 7

Discussion

The integrated system reveals:

- Long-term upward drift in Chla → rising chlorophyll concentrations.
- Moderate turbidity variability → seasonal hydrodynamics.
- NDWI shrinkage rhythm → monsoon recovery followed by dry-season decline.
- ML models accurately predict Chla and turbidity.

Chapter 8

Conclusion

NEREUS successfully integrates satellite-derived indices, machine learning, and generative AI to provide a full water-quality intelligence framework. With 10 analytical figures, predictive modeling, and AI explanations, it offers a robust approach to freshwater monitoring.