

Q1.

a) $x_0 = \int_0^1 t^0 (t+7)^{-1} dt$
 $= \int_0^1 (t+7)^{-1} dt$

Let $u = t+7$

$du = dt$

$$x_0 = \int \frac{1}{u} du = [\ln|t+7|]_0^1$$

$$= \ln 8 - \ln 7 = \ln \frac{8}{7}$$

for $n=1$, $x_1 = \int_0^1 t^1 (t+7)^{-1} dt$

$$= \int_0^1 \frac{t}{t+7} dt = \int_0^1 \left(1 - \frac{7}{t+7}\right) dt = 1 - 7x_0$$

$$x_n = \frac{1}{n} - 7x_{n-1} = \int_0^1 t^n (t+7)^{-1} dt$$

$$\frac{1}{n} = \int_0^1 \frac{t^n}{t+7} dt + 7 \int_0^1 \frac{t^{n-1}}{t+7} dt$$

$$RHS = \int_0^1 \frac{t^n + 7t^{n-1}}{t+7} dt$$

$$= \int_0^1 \frac{t^{n-1}(t+7)}{t+7} dt = \int_0^1 t^{n-1} dt = \left[\frac{t^n}{n} \right]_0^1 = \frac{1}{n}$$

$\therefore LHS$

\therefore proved

Q1 b)

```
In [1]: 1 import numpy as np  
2 import scipy.integrate  
3 import matplotlib.pyplot as plt
```

```
In [2]: 1 def f(x):  
2     return pow(x,18) / (x + 7)  
3 a = 0; b = 1
```

```
In [3]: 1 x18, err = scipy.integrate.quad(f, a, b)  
2 print("The approximation of x18 is ", x18, ", the error is ", e)
```

The approximation of x18 is 0.006620563824579688 , the error is 7.350302394048756e-17

```
In [4]: 1 def fb(x,n):  
2     return (1/n - x) / 7  
3 def ff(x,n):  
4     return 1/n - 7*x
```

```
In [5]: 1 xb = np.zeros(19)  
2 xb[18] = x18  
3 for i in range(len(xb)-2,-1,-1):  
4     xb[i] = fb(xb[i+1],i+1)  
5 print(f"xn by backward recurrence formulas:\n{xb}")
```

xn by backward recurrence formulas:
[0.13353139 0.06528025 0.04303824 0.03206566 0.02554036 0.02121749
 0.01814422 0.01584763 0.01406657 0.01264509 0.01148435 0.01051866
 0.0097027 0.00900417 0.00839942 0.00787076 0.00740469 0.00699071
 0.00662056]

```
In [6]: 1 xf = np.zeros(19)  
2 xf[0] = np.log(8/7)  
3 for i in range(1,len(xf)):  
4     xf[i] = ff(xf[i-1],i)  
5 print(f"xn by forward recurrence formulas:\n{xf}")
```

xn by forward recurrence formulas:
[0.13353139 0.06528025 0.04303824 0.03206566 0.02554036 0.02
 121749
 0.01814422 0.01584763 0.01406657 0.0126451 0.01148433 0.01
 051875
 0.00970209 0.00900842 0.00836967 0.008079 0.00594698 0.01
 719466
 -0.06480705]


The result given by backwards recurrence formulas is much more accurate than that of forward recurrence formulas.

```
In [7]: 1 float_info = np.finfo(float)
2 x0err = xf[0] - xb[0]
3 x18err = xf[18]-xb[18]
4 print(f" the difference between x0 times 7^18 {x0err*pow(7,18)})
```

the difference between x0 times 7^18 -0.09039511350064361
 the difference between x18 -0.0714276134172489

```
In [8]: 1 error = abs(xf-xb)
2 print(f"absolute value of errors are \n{error}")
3 n = np.linspace(0,18,19)
4 plt. figure(figsize= (13,8))
# plt.axes(yscale = "log")
5 plt.scatter(n,np.log(error),label='log(error)')
6 plt.title('Errors against number of iterations')
7 plt.xlabel('n')
8 plt.ylabel('log(error)')
9 plt.xlim(0,18)
10 plt.ylim(-40,0)
11 plt.legend(loc='best')
12 plt.grid()
13 plt.show()
```

absolute value of errors are
 [5.55111512e-17 3.05311332e-16 2.15105711e-15 1.50435220e-14
 1.05315062e-13 7.37212374e-13 5.16047621e-12 3.61233231e-11
 2.52863263e-10 1.77004285e-09 1.23902999e-08 8.67320995e-08
 6.07124696e-07 4.24987288e-06 2.97491101e-05 2.08243771e-04
 1.45770640e-03 1.02039448e-02 7.14276134e-02]



```
In [9]: 1 float_info = np.finfo(float)
2 x0err = xf[0] - xb[0]
3 x18err = xf[18]-xb[18]
4 print(f" the difference between x0 times 7^18 {x0err*pow(7,18)}")
```

```
the difference between x0 times 7^18 -0.09039511350064361
the difference between x18 -0.0714276134172489
```

The errors clearly forms a line in log scale, this means the error is multiplied by a certain number when we using the recurrence formulae to find the result. We can clearly find that when applying forwards and backwards recurrence formulas, together with the value of x_n , the error is multiplied by 7 and $1/7$ in each iteration respectively. Therefore after a number of iterations, the error by forwards recurrence formula is multiplied by 7^{18} which is obviously a large number, while the error in backwards recurrence formula is multiplied by $(1/7)^{18}$, which makes the starting error extremely small. The above results proves the error is roughly multiplied by 7^{18} after 18 iterations.

Q2.

$$\text{a) } X^{-1} = \begin{bmatrix} I - A^{-1}B \\ 0 & I \end{bmatrix} \begin{bmatrix} A^{-1} & 0 \\ 0 & (D - CA^{-1}B)^{-1} \end{bmatrix} \begin{bmatrix} I & 0 \\ -CA^{-1} & I \end{bmatrix}$$
$$= \begin{bmatrix} A^{-1} - A^{-1}B(D - CA^{-1}B)^{-1} \\ 0 & (D - CA^{-1}B)^{-1} \end{bmatrix} \cdot \begin{bmatrix} I & 0 \\ -CA^{-1} & I \end{bmatrix}$$

$$= \begin{bmatrix} A^{-1} + A^{-1}B(D - CA^{-1}B)^{-1} \cdot CA^{-1} & -A^{-1}B(D - CA^{-1}B)^{-1} \\ -(D - CA^{-1}B)^{-1} \cdot CA^{-1} & (D - CA^{-1}B)^{-1} \end{bmatrix}$$

$$\text{Let } (D - CA^{-1}B) = P, (D - CA^{-1}B)^{-1} = P^{-1}$$

$$\therefore X \cdot X^{-1} = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \cdot \begin{bmatrix} A^{-1} + A^{-1}BP^{-1}CA^{-1} & -A^{-1}BP^{-1} \\ -P^{-1} \cdot CA^{-1} & P^{-1} \end{bmatrix}$$
$$= \begin{bmatrix} I + BP^{-1}CA^{-1} - BP^{-1}CA^{-1} & -BP^{-1} + BP^{-1} \\ CA^{-1} + CA^{-1}BP^{-1}CA^{-1} - DP^{-1}CA^{-1} & -CA^{-1}BP^{-1} + DP^{-1} \end{bmatrix}$$
$$= \begin{bmatrix} I & 0 \\ CA^{-1} + (CA^{-1}B - D)P^{-1}CA^{-1} & (D - CA^{-1}B)P^{-1} \end{bmatrix}$$
$$= \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix}$$

Since we only need to compute the inverse of 2 smaller square matrices instead of a bigger one, the calculation is much easier.

Q2.

b) $A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, C = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$

$$D = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

$\therefore A^{-1} = I^{-1} = I, A^{-1}B = B, CA^{-1} = C$

$$\begin{aligned} P &= D - CA^{-1}B = D - CB = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

$$P^{-1} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

$$X^{-1} = \begin{bmatrix} I & -B \\ 0 & I \end{bmatrix} \begin{bmatrix} I & 0 \\ 0 & P^{-1} \end{bmatrix} \cdot \begin{bmatrix} I & 0 \\ -C & I \end{bmatrix}$$

$$= \begin{bmatrix} I & -BP^{-1} \\ 0 & P^{-1} \end{bmatrix} \cdot \begin{bmatrix} I & 0 \\ -C & I \end{bmatrix} = \begin{bmatrix} I + BP^{-1}C & -BP^{-1} \\ -P^{-1}C & P^{-1} \end{bmatrix}$$

$$BP^{-1}C = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$$

$$BP^{-1} = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix}, P^{-1}C = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\therefore X^{-1} = \begin{bmatrix} 1 & 0 & 0 & -1 \\ 1 & 1 & -1 & 1 \\ -1 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Q3.

a) The condition number of a matrix A

is given by $k(A) = \|A\| \|A^{-1}\|$

$$\|A\|_1 = \|A\|_\infty = \max(|\alpha|+1, 2) \quad A^{-1} = \frac{1}{\alpha-1} \begin{bmatrix} 1 & -1 \\ -1 & \alpha \end{bmatrix}$$

$$\therefore \|A^{-1}\|_1 = \|A^{-1}\|_\infty = \frac{\max(|\alpha|+1, 2)}{|\alpha|-1}$$

$$\therefore k(A) = \frac{(\max(|\alpha|+1, 2))^2}{|\alpha|-1}, \text{ for } k(A) \text{ to be large,}$$

$\alpha-1$ should be close to 0

but $\alpha-1$ comes from A^{-1} ,

which means when A is closer to singular,

$k(A)$ is more likely to be bigger

and $k(A)$ is infinity when A is singular

where $\alpha = 1$

Q3.

b) $\|A\| = \max \left\{ \frac{\|Ax\|}{\|x\|}, x \in \mathbb{R}^n, x \neq 0 \right\}$, when $x=0$
 $0 > 0$ exists

Since $\|Ax\| \geq \theta \|x\| \forall x \text{ such } \theta > 0$

$$\forall x \neq 0, \|A\| = \frac{\|Ax\|}{\|x\|} \geq \theta$$

a singular matrix A can turn $Ax=y$ to zero but
can't bring x back by $A^{-1}y$ because A^{-1} don't exist.

\therefore if A is singular $\min \frac{\|Ax\|}{\|x\|} = 0$ instead of
some $\theta > 0$,

\therefore A is nonsingular

$$\therefore \min \frac{\|Ax\|}{\|x\|} = \min \frac{\|y\|}{\|A^{-1}y\|} = \frac{1}{\max \frac{\|A^{-1}y\|}{\|y\|}}$$

since $\|Ax\| \leq \|A\| \|x\|$,

similarly $\|A^{-1}y\| \leq \|A^{-1}\| \|y\|$

$$\therefore \max \frac{\|A^{-1}y\|}{\|y\|} = \|A^{-1}\|$$

$$\therefore \min \frac{\|Ax\|}{\|x\|} = \frac{1}{\|A^{-1}\|}, \text{ since } \frac{\|Ax\|}{\|x\|} \geq \theta$$

$$\therefore \frac{1}{\|A^{-1}\|} \geq \theta, \|A^{-1}\| \leq \theta^{-1}$$

Q4.

Book1 — Saved to my Mac

Home Insert Draw Page Layout Formulas Data Review View Add-ins Tell me

Cut Copy Format Calibri (Body) 12 A Wrap Text General Conditional Format as Table Normal Bad Good Neutral Calculation Check Cell Auto-sum Fill Sort & Filter Insert Delete Format Clear Analyse Data Sensitivity

G1 fx 0

G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W	X	Y	Z	AA	AB	AC	AD	AE	AF	AG				
1	b																				Ly = b	Ux = y								
2	0.100	-1.732	1	0	0	0	0	0	0	0		1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	y	xhat	x	xhat-x						
3	0.100	1	-1.732	1	0	0	0	0	0	0		-0.5774	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.1	-0.3890475	0.15766089	-0.546708382					
4	0.100	0	1	-1.732	1	0	0	0	0	0		0.000	-0.8661	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.15774	-0.5738302	0.37306867	-0.946898918				
5	0.100	0	0	1	-1.732	1	0	0	0	0		0.000	0.000	-1.155	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.23661861	-0.5050339	0.58849404	-1.093527978				
6	0.100	0	0	0	1	-1.732	1	0	0	0		0.000	0.000	0.000	-1.733	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.3732945	-0.2006903	0.74620301	-0.946893281			
7	0.100	0	0	0	0	1	-1.732	1	0	0		0.000	0.000	0.000	0.000	1640	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.74691937	0.25745607	0.80392957	-0.546473497		
8	0.100	0	0	0	0	1	-1.732	1	0	0		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-1224.8478	0.74676234	0.74620301	0.000559336					
9	0.100	0	0	0	0	0	1	-1.732	1	0		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.6462997	0.58924551	0.58849404	0.000751472					
10	0.100	0	0	0	0	0	0	1	-1.732	1		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.2733027	0.37368424	0.37306867	0.000615571					
11	0.100	0	0	0	0	0	0	0	1	-1.732		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.1367348	0.15792888	0.15766089	0.000267987					
12																														
13																														
14																														
15																														
16																														
17																														
18																														
19																														
20																														
21																														
22																														
23																														
24																														
25																														
26																														
27																														
28																														
29																														
30																														
31																														
32																														
33																														
34																														
35																														
36																														
37																														
38																														
39																														
40																														
41																														
42																														
43																														
44																														
45																														
46																														
47																														
48																														
49																														
50																														
51																														
52																														

Q4.

$$\left[\begin{array}{ccccccccc} -1.732 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -1.732 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1.732 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1.732 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1.732 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1.732 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -1.732 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -1.732 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -1.732 \end{array} \right] \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \\ x_7 \\ x_8 \\ x_9 \end{pmatrix} = \begin{pmatrix} 0.100 \\ 0.100 \\ 0.100 \\ 0.100 \\ 0.100 \\ 0.100 \\ 0.100 \\ 0.100 \\ 0.100 \end{pmatrix}$$

a) the green colored matrices

b) $\|\hat{x} - x\|_2 \approx 1.894$

c) $\|\hat{x} - x\|_2$ of LUP decomposition

is 0.0001481, much smaller than LU factorisation