



Long-Term Changes of Optical Properties of Mineral Dust And Its Mixtures Derived From Raman Polarization Water Vapor Lidar in Central Europe

D.M. Szczepanik¹, W. Kumala¹, C.F. Olusegun ¹, E. Tetoni², V. Amiridis³, D. Nicolae⁴, D. Althausen⁵, I.S. Stachlewska¹

¹Faculty of Physics, University of Warsaw (UW), Poland

²Deutsches Zentrum für Luft- und Raumfahrt (DLR), Germany

³National Observatory of Athens (NOA), Greece

⁴National Institute for Research and Development of Optoelectronics Bucharest INOE 2000, Romania

⁵Leibniz Institute for Tropospheric Research (TROPOS), Germany

03. Atmospheric aerosol and clouds properties

28 June 2022 (Tuesday), 12:00 UTC

Poster P21

→ *The Earth's climate and its changes are one of the hot topics in nowadays science.*

How and how fast is the climate changing?

Do the climate changes affect the aerosol loading within the atmosphere?

Are there any changes in the aerosol-climate feedback observed?

→ *Desert dust – one of the major natural aerosols – was investigated over Warsaw (Central Europe).*

→ *Unique database of lidar-derived optical properties of mineral dust and its mixtures was derived and analyzed.*



Figure 1: *Polly^{XT} lidar in the RS-Lab station in Warsaw, Poland.*
(photo: I. Stachlewska)

Methodology

- Polly^{XT} lidar at the RS-Lab Warsaw, Poland (52.21 °N, 20.98 °E, 96 m a.s.l.) conducts 24h/7 automatic measurements at 1064, 532, and 355 nm (elastic), 532 and 355nm (elastic cross), 607 and 387 nm (Raman N₂) and 407 nm (Raman H₂O, Engelman et al. 2016).
- Data between July 2013 and December 2020 were examined for long-range transport of mineral dust (high depolarization in the free troposphere and raised air temperatures).
- Mineral dust inflows identification based on models: backward trajectory model - HYSPLIT, mineral dust prediction model - BSC-DREAM8b, and aerosol prediction model - NAAPS.
- Data analyzed using the classical Raman approach, which enables obtaining the full sets of optical properties ($3\beta+2\alpha+2\delta$).

(more details in Szczepanik et al. 2021)

- Our study revealed **20 independent episodes** of mineral dust in free troposphere over Warsaw.
- For the identified cases, we obtained **120 sets of profiles** containing at least particle backscattering and depolarization.
- Three groups were classified: **pure mineral dust (MD)**, **mineral dust with sulfates (MD+S)**, and **mineral dust with biomass burning aerosols (MD+BB)**.

In Warsaw, the δ values for clean mineral dust are lower than for the dust observed close to the source (Mamouri & Ansmann, 2014).

The mixed in additional aerosol affects the optical properties of mineral dust:

- **sulfate's presence** slightly lowers values of all properties;
- **Biomass burning** admixture increases α values significantly, while δ values are the lowest;
- The amount of the additive may also play a role here, but it is hard to estimate it.

***Table 1.** Average values of optical properties and layer height for: pure desert dust (MD) and its mixtures with sulfates (MD+S) and with aerosol originating from biomass combustion (MD+BB) observed over Warsaw in 2013-2020.*

Parameter	MD	MD+S	MD+BB
β 355 nm [$\text{Mm}^{-1}\text{sr}^{-1}$]	1.60 ± 0.76	1.32 ± 0.92	3.16 ± 0.85
β 532 nm [$\text{Mm}^{-1}\text{sr}^{-1}$]	1.22 ± 0.50	0.93 ± 0.53	1.80 ± 0.80
β 1064 nm [$\text{Mm}^{-1}\text{sr}^{-1}$]	0.82 ± 0.49	0.45 ± 0.31	1.08 ± 0.57
α 355 nm [Mm^{-1}]	69 ± 29	78 ± 41	248 ± 73
α 532 nm [Mm^{-1}]	48 ± 16	49 ± 28	173 ± 85
δ 355 nm [%]	10.53 ± 3.86	6.77 ± 2.99	2.23 ± 0.83
δ 532 nm [%]	15.46 ± 5.49	10.79 ± 3.22	4.49 ± 2.16

Frequency of dust occurrences and its length analysis

- ↓ The number of dust inflows over Warsaw is increasing from 1-2 episodes per year before 2017 to even 6 cases in 2019;
- ↓ The length of the observed dust inflow is getting longer starting from 2016, reaching up to 36 h of mineral dust observations per event.




Figure 2: Histogram showing the number of inflows and their average duration for the identified episodes of the desert dust inflow over Warsaw.

Tracking the stage of inflow evolution




Figure 3: Values of δ and LR for 355 nm (left) and 532 nm (right) depending on dust inflow evolution (number of points is 83 for 355nm; 120 for 532nm).

- ↑ The highest δ values are during the full evolution of the dust inflow, while the lowest is for the fadeaway phase.
- ↑ Some outlying points stand for the cases of observation of a mixture of dust with another aerosol.

Acknowledgements

- We thank **Volker Freudenthaler** at the MIM-LMU for evaluating the QA tests and **Lucia Mona** at the IMMA-CNR for running the QC procedures.
- We acknowledge the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT transport and dispersion model and/or READY website (<https://www.ready.noaa.gov>), the NAAPS model output obtained via NRL/ Monterey Aerosol Page (<https://www.nrlmry.navy.mil>), and the BSC-DREAM8b Barcelona Supercomputing Center which were used via webpage <https://ess.bsc.es/bsc-dust-daily-forecast>.
- The work was supported by European Commission grants: ACTRIS (no. 262254), ACTRIS-2 (no. 654109), and ACTRIS PPP (no. 739530).
- The research was partly funded by the NCN project PRELUDIUM-19 no: 2020/37/N/ST10/02682 entitled: *Study of differences in optical and microphysical properties of local agricultural dust and long range transported desert dust*.

Literature

- Engelman, R., Kanitz, T., Baars, H., Heese, B., Althausen, D., Skupin, A., Wandinger, U., Komppula, M., Stachlewska, I. S., Amiridis, V., Marinou, E., Mattis, I., Linné, H., and Ansmann, A.: *The automated multiwavelength Raman polarization and water-vapor lidar PollyXT: The next generation.*, *Atmos. Meas. Tech.*, 9, 1767-1784, doi:10.5194/amt-9-1767-2016, 2016.
- Mamouri, R. E., and Ansmann, A.: *Fine and coarse dust separation with polarization lidar*, *Atmos. Meas. Tech.*, 7, 3717-3735, <https://doi.org/10.5194/amt-7-3717-2014>, 2014.
- Szczepanik, D. M., Stachlewska, I. S., Tetoni, E., and Althausen, D.: *Properties of Saharan Dust versus Local Urban Dust – a case study*. *Earth and Space Science*, 10.1029/2021EA001816, 2021.