



Cloud Base Height Comparison of Co-Located Micro-pulse LiDAR and Lufft Ceilometer

Victoria Pinnegar¹, Paul Christiaans¹, Joe Clarke¹, Alexander Haefele², E. J. Welton³, Robert Sica¹

¹ Department of Physics and Astronomy, University of Western Ontario, London, ON, Canada

² Federal Office of Meteorology and Climatology MeteoSwiss, Payerne, Switzerland

³ Goddard Space Flight Center, NASA, Greenbelt, MD 20771, USA

03- Atmospheric Aerosol and Cloud Properties

27th of June, 14:15

Monday_ 03_P12



MPLNET Home



E-PROFILE Home



MPLCAN Home

- MPLCAN: Micro-Pulse Lidar CANada Network
- 5 Network sites at London ON, Sherbrooke QU, Halifax NS, Eureka NU and Toronto ON (PI Debra Wunch)
- Objectives of MPLCAN included developing cross country smoke tracking, interaction and improvements in Aerosol characterization and Air Quality
- Micro-Pulse Lidar and Lufft Ceilometer operate together at an MPLCAN site partnered with MPLNET and EUMETNET's E-PROFILE
- London site has been operating since December 2020

Motivations

1. To accurately compare the data products of MPLNET and E-PROFILE
2. To produce a Cloud Base product that can be applied to both instruments and thus across the network
3. To minimize the difference in height detection between instruments


CBH correlation between Co-Located MPL and CHM15k



Lidar Optical System	Micro-Pulse Lidar	CHM15k
Type	Polarization	Elastic Backscatter
Wavelength	532 nm	1064 nm
Energy per Pulse	3 – 4 μJ	8 μJ
Range Resolution	5/15/30/75 m	5 m
Power Aperture Product	4.398E-05 W m ²	1.909E-05 W m ²
Mirror Diameter	80 mm	90 mm
Range	100m - 30 km	5m – 15 km
Cloud Detection	around 15 km	around 12 km



- Simultaneous operation since December 2020
- CHM 15k connected to E-PROFILE
- miniMPL data hosted on MPLNET
- Using the Cloud Base Height (CBH) Product produced by MPLNET and the Lufft (CBH), the linear regression between the two instruments has a high comparison.
- Higher comparison in low aerosol winter



Case 1: Lufft CHM15k Cloud Base Altitude vs MPLNET V3 Cloud Base Altitude. Equinox (Spring, Autumn), Summer and Winter 2021-2022

- Utilizes profiles in five-minute intervals
- If $\frac{dR(z)}{dz} > 0$, there is a minimum 60 m increasing gradient, and the value of $\beta > a_{thres}$, z value is marked as Cloud Base.
- If profile is clear, and previous profile has a cloud, profile is rechecked with $\beta > 0.5 a_{thres}$
- Based on MPLNET V2 [1]

$$R(z) = \frac{\beta(z)T^2(z)}{\beta_{mol}(z)T_{mol}^2(z)} \quad (1)$$

$$a_{thres} = C\bar{\beta} \quad (2)$$

$\beta(z)$ = Normalized Backscatter

$T^2(z)$ = Two – way Atmospheric Transmission

$\bar{\beta}$ = Spacial Average of Normalized Backscatter

a_{thres} = Threshold

$R(z)$ = attenuated backscatter ratio

C = empirical value






Figure: Example cloud located between 2 and 3 km with marked cloud base observation.





Case 2: Gradient Based Detection
(applied on MPL data) vs
MPLNET Cloud Base Height.
Equinox (Spring, Autumn),
Summer and Winter 2021-2022



Case 3: Gradient Based Detection
(applied on MPL data) vs Lufft
Cloud Base Height. Equinox
(Spring, Autumn), Summer and
Winter 2021-2022

Summary

- Initial network comparison is overall good
- Algorithm has comparable statistical values, and can be applied to instruments in both networks
- Open source and will be available on GitHub



Acknowledgments:

Thank you to NASA MPLNET for the completion of the Calibration, Université de Sherbrooke for the use of their facilities, Western University, Dalhousie University, University of Toronto, MeteoSwiss and L'Université du Québec à Montréal for their participation in the project

