



Laboratory Evaluation of the (355, 532) Depolarization Ratio of Pure Pollen

Danaël CHOLLETON, Patrick RAIROUX, Alain MIFFRE

Institute of Light and Matter (iLM), ATMOS Research group, Lyon University, France

03. Atmospheric aerosol and clouds properties

Tuesday June 28th, 12:00 UTC

P24

■ Health and economical impact


- Allergic reactions (asthma, rhinitis)
- 1/3 of the population in France (ANSES, 2021)
- Health cost of pollen : 151 billion €/year in Europe (Lake et al. 2017)

■ Climatic impact







- Regional warming through scattering and absorption (Spänkuch et al., 2002)
- Cloud and ice condensation nuclei (Pope et al., 2010; Wozniak et al. 2018)

■ Evolution in future years

- Higher pollen concentrations
- Longer pollen season
- Larger geographical locations



- Pure pollen samples



(a) To (d) *Ambrosia*, *Betula*,
Fraxinus, *Pinus*,
SEM Images, Univ Lyon 1
(e) And (f) : *Cupressus*, *Picea*
Paldat.org


- State of the art on pollen lidar remote sensing

- Depolarization lidar profiles of pollen mixtures :
Sassen 2008, Noh et al. 2013, Sicard et al. 2016,
Shang et al., 2022.
- First evaluation of the pollen PDR by Cao et al.
2010 at close but not exact backscattering angle.
- No accurate light-scattering numerical simulation
for such large and complex-shaped particles.

→ **Laboratory evaluation of pure pollen Lidar PDR
at 180.0° lidar backscattering angle.**

- Unique laboratory polarimeter at 180.0 lidar backscattering angle (Miffre et al. 2016, 2019)


➡ **See A. Miffre's talk tomorrow at 15:45 !**



$$PDR_{\lambda} = \frac{1 - F_{22}^{\lambda}/F_{11}^{\lambda}}{1 + F_{22}^{\lambda}/F_{11}^{\lambda}}$$


- **Time-resolved:** ns-pulsed laser source (as in lidar applications)
- **High sensitivity:** polarization cross-talks are fully negligible (as low as 10^{-7}).
- **Multi-spectral:** 355 and 532 nm lidar wavelengths.
- **Accurate (1%) spectroscopic evaluation of the PDR of pure pollen.**

- Polarimeter validated on spherical particles $\rightarrow PDR_{\lambda} = 0$ as expected by Lorenz-Mie theory
- Evaluation of the PDR by modulating the backscattered intensity with a retarder plate (angle ψ) :



$$PDR_{355} = 5.2 \pm 0.5 \%$$


$$PDR_{532} = 32.7 \pm 0.3 \%$$



Pure pollen taxon	PDR_{355} [%]	PDR_{532} [%]
Ragweed	5.2 ± 0.5	32.7 ± 0.3
Birch	3.8 ± 0.7	31.1 ± 0.6
Ash tree	5.0 ± 0.6	43.8 ± 0.2
Pine	22.2 ± 0.7	41.3 ± 0.5
Spruce	25.0 ± 1.0	37.2 ± 0.8
Cypress	17.4 ± 0.5	39.8 ± 0.3

Precision on lidar PDR of pure pollen : helps interpreting lidar observations of pollen mixtures (Sassen 2008, Noh et al. 2013, Sicard et al. 2016), and discussing the validity of inversion methodologies (Shang et al., 2022).

Pure Pollen Particle Depolarization Ratio [%]



To be submitted to Remote Sensing
Special Issue 2022 ELC Conference