

FIGURES

| | | |
|------|--|----|
| 2.1 | Various ways of interpreting the example ambiguous transformation: (a) change of coordinates; (b) transformation of plane onto itself; and (c) transformation from one plane to another. | 11 |
| 2.2 | The solutions of the linear equation $3x_1 + 2x_2 = 6$. | 26 |
| 2.3 | Three possible two-equation linear system solutions. | 27 |
| 2.4 | Schematic diagram of a function. | 46 |
| 2.5 | Composition of two functions. | 47 |
| 2.6 | One-to-one, onto, and isomorphic maps. | 48 |
| 2.7 | One-to-one and onto functions. | 48 |
| 2.8 | An invertible mapping. | 49 |
| 2.9 | Least squares example. | 57 |
| 3.1 | Vectors as directed line segments. | 64 |
| 3.2 | Two vectors. | 65 |
| 3.3 | Two vectors drawn head-to-tail. | 65 |
| 3.4 | Vector addition. | 66 |
| 3.5 | Vector addition chain. | 66 |
| 3.6 | Vector subtraction. | 67 |
| 3.7 | Vector multiplication. | 67 |
| 3.8 | Vector negation. | 68 |
| 3.9 | Commutativity of vector addition. | 68 |
| 3.10 | Associativity of vector addition. | 68 |
| 3.11 | Distributivity of addition over multiplication. | 69 |
| 3.12 | Distributivity of multiplication over addition. | 69 |
| 3.13 | The span of two vectors in 3D space is a plane. | 70 |
| 3.14 | A vector as the linear combination of basis vectors. | 72 |
| 3.15 | Angle between vectors. | 74 |
| 3.16 | The right-hand rule for orientation. | 75 |
| 3.17 | A vector as the linear combination of two different sets of basis vectors. | 76 |
| 3.18 | The sine function. | 77 |

| | | |
|------|---|-----|
| 3.19 | Linear transformation “scale by two.” | 78 |
| 3.20 | Nonuniform scale linear transformation. | 79 |
| 3.21 | Rotation transformation. | 79 |
| 3.22 | Shear transformation. | 79 |
| 3.23 | Definition of point subtraction. | 81 |
| 3.24 | The Head-to-Tail axiom. | 81 |
| 3.25 | Affine combination of two points. | 83 |
| 3.26 | Affine combination of several points. | 85 |
| 3.27 | Angle between vectors and vector length. | 86 |
| 3.28 | Parallelogram rule for vector addition. | 87 |
| 3.29 | Vector projection. | 87 |
| 3.30 | $\cos \theta$ negative. | 89 |
| 3.31 | The vector product. | 92 |
| 3.32 | The right-hand rule. | 93 |
| 3.33 | Parallelepiped defined by three vectors. | 94 |
| 3.34 | The scalar triple product. | 95 |
| 3.35 | Coordinates of an affine point, relative to an arbitrary frame. | 98 |
| 3.36 | Affine maps preserve relative ratios. | 99 |
| 3.37 | Vector sum. | 101 |
| 3.38 | Vector scale. | 102 |
| 3.39 | Sum of point and vector. | 102 |
| 3.40 | Composition of affine maps (rotation). | 104 |
| 3.41 | Affine (a) and barycentric (b) coordinates. | 105 |
| 3.42 | The first three simplexes: a line (a), a triangle (b), and a tetrahedron (c). | 107 |
| 4.1 | $P = p_1\vec{i} + p_2\vec{j} + p_3\vec{k} + \mathcal{O} = [p_1 \ p_2 \ p_3 \ 1]$. | 114 |
| 4.2 | $\vec{v} = v_1\vec{i} + v_2\vec{j} + v_3\vec{k} = [v_1 \ v_2 \ v_3 \ 0]$. | 114 |
| 4.3 | The “perp” operator. | 122 |
| 4.4 | The perp dot product reflects the signed angle between vectors. | 124 |
| 4.5 | The perp dot product is related to the signed area of the triangle formed by two vectors. | 125 |
| 4.6 | Representing P in \mathcal{A} and \mathcal{B} . | 129 |
| 4.7 | Representing \mathcal{O} in \mathcal{G} . | 130 |
| 4.8 | Representing \vec{v}_i in \mathcal{B} . | 131 |
| 4.9 | Translation. | 134 |
| 4.10 | Translation of a frame. | 136 |
| 4.11 | Simple rotation of a frame. | 137 |
| 4.12 | General rotation. | 140 |

| | | |
|------|---|-----|
| 4.13 | General rotation shown in the plane \mathcal{A} perpendicular to \hat{u} and containing P . | 141 |
| 4.14 | Scaling a frame. | 143 |
| 4.15 | Uniform scale. | 145 |
| 4.16 | Nonuniform scale. | 146 |
| 4.17 | Mirror image. | 148 |
| 4.18 | Simple reflection in 2D. | 150 |
| 4.19 | Simple reflection in 3D. | 151 |
| 4.20 | General reflection in 3D. | 152 |
| 4.21 | Mirror image in 2D. | 153 |
| 4.22 | Mirror image in 3D. | 154 |
| 4.23 | Shearing in 2D. | 155 |
| 4.24 | $T_{xy,\theta}$. | 155 |
| 4.25 | General shear specification. | 157 |
| 4.26 | Orthographic (orthogonal) projection. | 160 |
| 4.27 | Oblique projection. | 161 |
| 4.28 | Edge-on view of oblique projection. | 162 |
| 4.29 | Perspective projection. | 163 |
| 4.30 | Cross-ratio. | 164 |
| 4.31 | Perspective map for vectors. | 165 |
| 4.32 | The plane $x + y = k$. | 166 |
| 4.33 | Incorrectly transformed normal. | 167 |
| 4.34 | Normal as cross product of surface tangents. | 167 |
| 5.1 | Examples of (a) a line, (b) a ray, and (c) a segment. | 172 |
| 5.2 | Implicit definition of a line. | 173 |
| 5.3 | The two possible orderings for a triangle. | 175 |
| 5.4 | The domain and range of the parametric form of a triangle. | 176 |
| 5.5 | The domain and range of the barycentric form of a triangle. | 176 |
| 5.6 | The domain and range for the parametric form of a rectangle. | 177 |
| 5.7 | The symmetric form of a rectangle. | 178 |
| 5.8 | A typical polyline. | 178 |
| 5.9 | Examples of (a) a simple concave polygon and (b) a simple convex polygon. | 179 |
| 5.10 | Examples of nonsimple polygons. (a) The intersection is not a vertex. (b) The intersection is a vertex. The polygon is a polysolid. | 180 |
| 5.11 | Examples of polygonal chains: (a) strictly monotonic; (b) monotonic, but not strict. | 180 |

| | | |
|------|---|-----|
| 5.12 | A monotone polygon. The squares are the vertices of one chain. The triangles are the vertices of the other chain. The circles are those vertices on both chains. | 181 |
| 5.13 | Solutions to the quadratic equation depending on the values for $d_0 \neq 0$, $d_1 \neq 0$, and r . | 183 |
| 5.14 | Solutions to the quadratic equation depending on the values for $d_0 \neq 0$, $d_1 = 0$, e_1 , and r . | 184 |
| 5.15 | Circles defined in distance (implicit) and parametric forms. | 184 |
| 5.16 | Definition of an ellipse. | 185 |
| 5.17 | A cubic Bézier curve. | 186 |
| 5.18 | A cubic B-spline curve. | 187 |
| 6.1 | Closest point $X(\bar{t})$ on a line to a specified point Y . | 190 |
| 6.2 | Closest point on a ray to a given point: (a) $X(\bar{t})$ closest to Y ; (b) P closest to Y . | 192 |
| 6.3 | Closest point on a segment to a given point: (a) $X(\bar{t})$ closest to Y ; (b) P_0 closest to Y ; (c) P_1 closest to Y . | 192 |
| 6.4 | The segment S_0 generated the current minimum distance μ between the polyline and Y . S_1 and S_2 cannot cause μ to be updated because they are outside the circle of radius μ centered at Y . Segment S_3 does cause an update since it intersects the circle. The infinite-strip test does not reject S_1 and S_3 since they lie partly in both infinite strips, but S_2 is rejected since it is outside the vertical strip. The rectangle test rejects S_1 and S_2 since both are outside the rectangle containing the circle, but does not reject S_3 . | 195 |
| 6.5 | Closest point on a triangle to a given point: (a) $\text{Dist}(Y, \mathcal{T}) = 0$; (b) $\text{Dist}(Y, \mathcal{T}) = \text{Dist}(Y, \langle P_0, P_1 \rangle)$; (c) $\text{Dist}(Y, \mathcal{T}) = \text{Dist}(Y, P_2)$; (d) $\text{Dist}(Y, \mathcal{T}) = \text{Dist}(Y, \langle P_1, P_2 \rangle)$. | 197 |
| 6.6 | Partitioning of the parameter plane into seven regions. | 199 |
| 6.7 | Contact points of level curves of $F(t_0, t_1)$ with the triangle: (a) contact with an edge; (b) contact with a vertex. | 199 |
| 6.8 | Contact points of level curves of $F(t_0, t_1)$ with the triangle: (a) contact with an edge; (b) contact with another edge; (c) contact with a vertex. | 200 |
| 6.9 | A triangle, a bounding box of the triangle, and the regions of points closest to vertices and to edges. | 207 |
| 6.10 | Partitioning of the plane by a rectangle. | 212 |
| 6.11 | (a) An example of a single cone. (b) A frustum of a cone. (c) An orthogonal frustum. | 214 |
| 6.12 | Portion of frustum in first quadrant. | 215 |
| 6.13 | Only those edges visible to the test point must be searched for the closest point to the test point. The three visible edges are dotted. The | |

| | | |
|------|--|-----|
| | invisible edges are drawn in black. The visible edges are in a cone with vertex at the test point and whose sides are tangent to the convex polygon. | 217 |
| 6.14 | Closest point on a quadratic curve to a given point. | 218 |
| 6.15 | Closest point on a polynomial curve to a given point. | 220 |
| 6.16 | Various line-line configurations: (a) zero distance; (b) positive distance. | 222 |
| 6.17 | Various line-ray configurations: (a) zero distance; (b) positive distance. | 223 |
| 6.18 | Various line-segment configurations: (a) zero distance; (b) positive distance. | 224 |
| 6.19 | Various nonparallel ray-ray configurations: (a) zero distance; (b) positive distance from end point to interior point; (c) positive distance from end point to end point. | 224 |
| 6.20 | Relationship of level curves of F to boundary minimum at $(\hat{t}_0, 0)$ or $(0, 0)$. | 226 |
| 6.21 | Various parallel ray-ray configurations: (a) rays pointing in the same direction; (b) rays pointing in opposite directions with overlap; (c) rays pointing in opposite directions with no overlap. | 227 |
| 6.22 | The configuration for the segment S attaining current minimum distance μ that is the analogy of Figure 6.4 for the point Y attaining current minimum distance. | 230 |
| 6.23 | Segment connecting closest points is perpendicular to both objects. | 232 |
| 6.24 | (a) Triangles A and B ; (b) set $-B$; (c) set $A + B$; (d) set $A - B$, where the gray point is the closest point in $A - B$ to the origin. The black dots are the origin $(0, 0)$. | 235 |
| 6.25 | The first iteration in the GJK algorithm. | 237 |
| 6.26 | The second iteration in the GJK algorithm. | 237 |
| 6.27 | The third iteration in the GJK algorithm. | 238 |
| 6.28 | The fourth iteration in the GJK algorithm. | 238 |
| 6.29 | (a) Construction of V_{k+1} in the convex hull of $S_k \cup \{W_k\}$. (b) The new simplex \bar{S}_{k+1} generated from $M = \{W_0, W_2, W_3\}$. | 239 |
| 7.1 | An arc of a circle spanned counterclockwise from A to B . The line containing A and B separates the circle into the arc itself and the remainder of the circle. Point P is on the arc since it is on the same side of the line as the arc. Point Q is not on the arc since it is on the opposite side of the line. | 249 |
| 7.2 | Intersection of a line and a cubic curve. | 250 |
| 7.3 | Line-curve intersection testing using a hierarchy of bounding boxes. | 252 |

| | | |
|------|--|-----|
| 7.4 | A line and a curve rasterized on a grid that is initially zero. The line is rasterized by or-ing the grid with the mask 1 (light gray). The curve is rasterized by or-ing the grid with the mask 2 (dark gray). Grid cells that contain both the line and the curve have a value 3 (dotted). | 254 |
| 7.5 | Intersections of an ellipse and a parabola. | 256 |
| 7.6 | Relationship of two circles, $\vec{u} = C_1 - C_0$: (a) $\ \vec{u}\ = r_0 + r_1 $; (b) $\ \vec{u}\ = r_0 - r_1 $; (c) $ r_0 - r_1 < \ \vec{u}\ < r_0 + r_1 $. | 258 |
| 7.7 | \mathcal{E}_1 is contained in \mathcal{E}_0 . The maximum \mathcal{E}_0 level curve value λ_1 for \mathcal{E}_1 is negative. | 259 |
| 7.8 | \mathcal{E}_1 transversely intersects \mathcal{E}_0 . The minimum \mathcal{E}_0 level curve value λ_0 for \mathcal{E}_1 is negative; the maximum value λ_1 is positive. | 260 |
| 7.9 | \mathcal{E}_1 is separated from \mathcal{E}_0 . The minimum \mathcal{E}_0 level curve value λ_0 for \mathcal{E}_1 is positive. | 261 |
| 7.10 | Intersection of two ellipses. | 262 |
| 7.11 | Two curves rasterized on a grid that is initially zero. The first curve is rasterized by or-ing the grid with the mask 1 (light gray). The second curve is rasterized by or-ing the grid with the mask 2 (dark gray). Grid cells that contain both curves have a value 3 (dotted). | 264 |
| 7.12 | Nonintersecting convex objects and a separating line for them. | 266 |
| 7.13 | (a) Nonintersecting convex polygons. (b) Intersecting convex polygons. | 267 |
| 7.14 | (a) Edge-edge contact, (b) vertex-edge contact, and (c) vertex-vertex contact. | 267 |
| 7.15 | The edge normals closest to a non-edge-normal separation direction: (a) from the same triangle and (b) from different triangles. | 268 |
| 7.16 | Two polygons separated by an edge-normal direction of the first polygon. | 270 |
| 7.17 | (a) Edge-edge intersection predicted. (b) Vertex-vertex intersection predicted. (c) No intersection predicted. | 276 |
| 7.18 | Edge-edge contact for two moving triangles. | 282 |
| 8.1 | Circle through three points. | 286 |
| 8.2 | Circle tangent to three lines. | 286 |
| 8.3 | Line tangent to a circle at a given point. | 287 |
| 8.4 | Line through point, tangent to a circle. | 288 |
| 8.5 | In general, there are two tangents, but there may be one or none. | 288 |
| 8.6 | Line tangent to two circles. | 291 |
| 8.7 | Depending on the relative sizes and positions of the circles, the number of tangents between them will vary. | 292 |
| 8.8 | Circle through two points with a given radius. | 297 |
| 8.9 | Both possible circles through two points with a given radius. | 297 |

| | | |
|------|---|-----|
| 8.10 | Insight for computing circle of given radius through two points. | 298 |
| 8.11 | Circle through a point and tangent to a line with a given radius. | 299 |
| 8.12 | In general, there are two distinct circles through the given point. | 299 |
| 8.13 | If P lies on the line, the circles are mirrored across the line; if P is further from the line than the diameter of the circle, there are no solutions. | 300 |
| 8.14 | Circles tangent to two lines with a given radius. | 303 |
| 8.15 | In general, there are four circles of a given radius tangent to two lines. | 303 |
| 8.16 | Constructive approach for circle tangent to two lines. | 304 |
| 8.17 | Circles through a point and tangent to a circle with a given radius. | 305 |
| 8.18 | Depending on the relative positions and radii of the circle, there may be four, two, or no solutions. | 306 |
| 8.19 | Insight for solving problem. | 307 |
| 8.20 | Constructive approach to solving problem. | 308 |
| 8.21 | Special case for constructive approach. | 309 |
| 8.22 | Circles tangent to a line and a circle with a given radius. | 309 |
| 8.23 | The number of distinct solutions varies depending on the relative positions of the line and circle, and the circle's radius. | 310 |
| 8.24 | No solutions if given radius is too small. | 311 |
| 8.25 | Insight for finding circle of given radius. | 311 |
| 8.26 | Schematic for the solution. | 312 |
| 8.27 | Circles tangent to two circles with a given radius. | 314 |
| 8.28 | In general there are two solutions, but the number of distinct solutions varies with the relative sizes and positions of the given circles. | 315 |
| 8.29 | Construction for a circle tangent to two circles. | 316 |
| 8.30 | Line normal to a given line and through a given point. | 317 |
| 8.31 | Line between and equidistant to two points. | 318 |
| 8.32 | Line parallel to a given line at a distance d . | 319 |
| 8.33 | Line parallel to a given line at a vertical or horizontal distance d . | 321 |
| 8.34 | Lines tangent to a given circle and normal to a given line. | 322 |
| 9.1 | A plane is defined as the set of all points X satisfying $\vec{n} \cdot (X - P) = 0$. | 327 |
| 9.2 | Geometric interpretation of plane equation coefficients. | 328 |
| 9.3 | The parametric representation of a plane. | 329 |
| 9.4 | The parametric representation of a circle in 3D. | 332 |
| 9.5 | A convex polygon and its decomposition into a triangle fan. | 334 |
| 9.6 | A nonconvex polygon and its decomposition into triangles. | 335 |
| 9.7 | A triangle mesh. | 335 |

| | | |
|------|--|-----|
| 9.8 | Vertices, edges, and triangles are not a mesh since a vertex is isolated. | 336 |
| 9.9 | Vertices, edges, and triangles are not a mesh since an edge is isolated. | 336 |
| 9.10 | Vertices, edges, and triangles are not a mesh since two triangles interpenetrate. | 337 |
| 9.11 | A polyhedron that consists of a tetrahedron, but an additional vertex was added to form a depression in the centered face. | 338 |
| 9.12 | A polymesh that is not a polyhedron since it is not connected. The fact that the tetrahedron and rectangle mesh share a common vertex does not make them connected in the sense of edge-triangle connectivity. | 338 |
| 9.13 | A polymesh that is not a polyhedron since an edge is shared by three faces. | 339 |
| 9.14 | A polytope, a regular dodecahedron. | 339 |
| 9.15 | The four possible configurations for ordering of two adjacent triangles. | 343 |
| 9.16 | A rectangle has two parallel edges joined together forming (a) a cylindrical strip (orientable) or (b) a Möbius strip (nonorientable). | 344 |
| 9.17 | The five Platonic solids. Left to right: tetrahedron, hexahedron, octahedron, dodecahedron, icosahedron. | 346 |
| 9.18 | Quadrics having three nonzero eigenvalues. | 353 |
| 9.19 | Quadrics having two nonzero eigenvalues. | 354 |
| 9.20 | Quadrics having one nonzero eigenvalue. | 355 |
| 9.21 | A standard “ring” torus. | 355 |
| 9.22 | A cubic Bézier curve. | 358 |
| 9.23 | A cubic B-spline curve. | 359 |
| 9.24 | A bicubic Bézier surface. | 361 |
| 9.25 | A cubic triangular Bézier surface. | 362 |
| 9.26 | A uniform bicubic B-spline surface. | 363 |
| 10.1 | Distance between a line and a point. | 366 |
| 10.2 | The projection of Q on \mathcal{L} . | 367 |
| 10.3 | Distance between a line segment and a point. | 368 |
| 10.4 | Utilizing half-spaces to speed up point/polyline distance tests. | 370 |
| 10.5 | Rejection example for point/polyline distance. | 371 |
| 10.6 | Distance between a point and a plane. | 374 |
| 10.7 | Edge-on view of plane \mathcal{P} . | 375 |
| 10.8 | Distance between a point and a triangle. The closest point may be on the interior of the triangle (a), on an edge (b), or be one of the vertices. | 376 |
| 10.9 | Partition of the st -plane by triangle domain D . | 378 |

| | | |
|-------|--|-----|
| 10.10 | Various level curves $Q(s, t) = V$. | 379 |
| 10.11 | Alternative definition of a rectangle. | 382 |
| 10.12 | Distance between a point and a rectangle. | 383 |
| 10.13 | Partition of the plane by a rectangle. | 383 |
| 10.14 | Distance from a point to a polygon. | 385 |
| 10.15 | Solving the 3D point-polygon distance test by projecting to 2D. | 386 |
| 10.16 | Typical case, closest point to circle. | 389 |
| 10.17 | Closest point is circle center. | 390 |
| 10.18 | Closest point when P projects inside the disk. | 390 |
| 10.19 | Closest point when P projects outside the disk. | 391 |
| 10.20 | Distance from a point to a polyhedron (tetrahedron). | 392 |
| 10.21 | Distance from a point to an oriented bounding box. | 394 |
| 10.22 | Computing the distance between a point and an OBB. | 395 |
| 10.23 | The portion of the frustum in the first octant. | 398 |
| 10.24 | Six possible “closest points” on an ellipsoid’s surface. | 404 |
| 10.25 | Distance from an arbitrary point to a parametric curve. | 405 |
| 10.26 | Distance from an arbitrary point to a parametric surface. | 407 |
| 10.27 | Distance between two lines. | 410 |
| 10.28 | Domains for each possible combination of linear component distance calculation. | 413 |
| 10.29 | Definition of visibility of domain boundaries. | 414 |
| 10.30 | Cases for the four edges of the domain. | 416 |
| 10.31 | Distance between two line segments. | 416 |
| 10.32 | Distance between a line and a ray. | 419 |
| 10.33 | Distance between a line and a line segment. | 420 |
| 10.34 | Distance between two rays. | 422 |
| 10.35 | Distance between a ray and a line segment. | 424 |
| 10.36 | Partitioning of the st -plane by the unit square. | 427 |
| 10.37 | Various level curves $Q(s, t) = V$. | 428 |
| 10.38 | Distance between a line and a triangle. | 434 |
| 10.39 | Parametric representation of a triangle. | 435 |
| 10.40 | Possible partitionings of the solution space for the linear component/triangle distance problem. | 436 |
| 10.41 | Boundary strip and planes for region 3. | 440 |
| 10.42 | Distance between a line and a rectangle. | 442 |
| 10.43 | The partitioning of the solution domain for a line segment and rectangle. | 444 |
| 10.44 | Distance between a line and a tetrahedron. | 447 |

| | | |
|-------|--|-----|
| 10.45 | Projecting a tetrahedron (a) onto a plane perpendicular to \hat{d} and then (b) into 2D. | 448 |
| 10.46 | Distance between a line and an oriented bounding box. | 451 |
| 10.47 | Schematic for line-OBB distance algorithm. | 452 |
| 10.48 | Case of two zero-components. | 453 |
| 10.49 | Case of one zero-component. | 456 |
| 10.50 | Determining where to look for the closest point on the box. | 457 |
| 10.51 | Determining whether the line intersects the box. | 458 |
| 10.52 | Each “positive” face of the OBB has two edges and three vertices that may be closest to the line. | 460 |
| 11.1 | Intersection of a line and a plane. | 483 |
| 11.2 | Intersection of a line and a triangle. | 486 |
| 11.3 | Intersection of a ray and a polygon. | 489 |
| 11.4 | Intersection of a linear component and a disk. | 492 |
| 11.5 | Intersection of a ray and a polyhedron (octahedron). | 494 |
| 11.6 | Intersection of a line segment and a polygonal (triangle) mesh. | 494 |
| 11.7 | The logical intersection of half-lines defines the intersection of a line with a polyhedron. | 496 |
| 11.8 | The logical intersection of half-lines fails to exist if the line does not intersect the polyhedron. | 498 |
| 11.9 | Possible ray-sphere intersections. | 503 |
| 11.10 | Intersection of a linear component and an ellipsoid. | 505 |
| 11.11 | Parameterized standard cylinder representation. | 508 |
| 11.12 | General cylinder representation. | 509 |
| 11.13 | Parameterized standard cone representation. | 513 |
| 11.14 | General cone representation. | 514 |
| 11.15 | An acute cone. The inside region is shaded. | 514 |
| 11.16 | An acute double cone. The inside region is shaded. | 515 |
| 11.17 | Case $c_2 = 0$. (a) $c_0 \neq 0$; (b) $c_0 = 0$. | 516 |
| 11.18 | Intersection of a ray with a NURBS surface. | 520 |
| 11.19 | Failed intersection calculation due to insufficient surface tessellation (shown in cross section for clarity). | 522 |
| 11.20 | A ray represented as the intersection of two planes. | 524 |
| 11.21 | Leaf-node bounding boxes are constructed from the Bézier polygon between each pair of refined vertices. | 528 |
| 11.22 | Adjacent bounding boxes are coalesced into a single box at the next level in the hierarchy. | 529 |
| 11.23 | Intersection of two planes. | 530 |

| | | |
|-------|---|-----|
| 11.24 | Possible configurations for three planes described in Table 11.1. | 532 |
| 11.25 | Plane-triangle intersection. | 535 |
| 11.26 | Plane-triangle intersection configurations. | 536 |
| 11.27 | Triangle-triangle intersection configurations: (a) $\mathcal{P}_0 \parallel \mathcal{P}_1$, but $\mathcal{P}_0 \neq \mathcal{P}_1$; (b) $\mathcal{P}_0 = \mathcal{P}_1$; (c) \mathcal{T}_0 intersects \mathcal{T}_1 ; (d) \mathcal{T}_0 does not intersect \mathcal{T}_1 . | 540 |
| 11.28 | Triangle-triangle interval overlap configurations: (a) intersection; (b) no intersection; (c) ?. | 541 |
| 11.29 | Intersection of a trimesh and a plane. | 544 |
| 11.30 | Intersection of a polygon and a plane. | 545 |
| 11.31 | Intersection of a polygon and a triangle. | 546 |
| 11.32 | Intersection of a plane and a sphere. | 548 |
| 11.33 | Cross-sectional view of sphere-plane intersection. | 550 |
| 11.34 | Intersection of a plane and a cylinder. | 551 |
| 11.35 | Some of the ways a plane and a cylinder can intersect. | 552 |
| 11.36 | Edge-on view of plane-cylinder intersection. | 554 |
| 11.37 | Ellipse in 3D. | 556 |
| 11.38 | Circle in 3D. | 556 |
| 11.39 | Dandelin's construction. | 557 |
| 11.40 | Cross section of a plane intersecting a cylinder, with the two spheres used to define the intersecting ellipse. After Miller and Goldman (1992). | 558 |
| 11.41 | The intersection of a plane and a cylinder is a circle if the plane's normal is parallel to the cylinder's axis. | 561 |
| 11.42 | Intersection of a plane and a cone. | 564 |
| 11.43 | Some of the ways a plane and a cone can intersect. | 565 |
| 11.44 | Intersection test for a plane and an infinite cone. | 566 |
| 11.45 | Edge-on view of plane-cone intersection. | 568 |
| 11.46 | Infinite cone definition. | 569 |
| 11.47 | Geometric definitions for hyperbola and parabola. | 570 |
| 11.48 | Parabolic curve intersection of plane and cone. After Miller and Goldman (1992). | 572 |
| 11.49 | Circular curve intersection of plane and cone. After Miller and Goldman (1992). | 576 |
| 11.50 | Ellipse intersection of plane and cone. After Miller and Goldman (1992). | 577 |
| 11.51 | Hyperbola intersection of plane and cone. After Miller and Goldman (1992). | 578 |

| | | |
|-------|--|-----|
| 11.52 | Degenerate intersections of a plane and a cone. After Miller and Goldman (1992). | 581 |
| 11.53 | Intersection of a plane and a parametric surface. | 587 |
| 11.54 | Hermite basis functions (cubic). | 589 |
| 11.55 | Cubic Hermite curve, specified by end points and tangents. | 590 |
| 11.56 | A subpatch in parameter space maps to a topologically rectangular region on the patch. After Lee and Fredricks (1984). | 591 |
| 11.57 | 3-space intersection curve $R(t)$. | 592 |
| 11.58 | Parametric space intersection curve $p(t)$. | 593 |
| 11.59 | Intersection of two B-spline surfaces. | 609 |
| 11.60 | Intersection curves in one surface's parameter space. | 610 |
| 11.61 | Intersection of a linear component with an axis-aligned bounding box. | 627 |
| 11.62 | Axis-aligned box as the intersection of three "slabs." | 628 |
| 11.63 | Clipping a line against a slab. | 629 |
| 11.64 | How slab clipping correctly computes ray-AABB intersection. | 631 |
| 11.65 | Specifying an oriented bounding box. | 631 |
| 11.66 | Clipping against an "oriented slab." | 632 |
| 11.67 | Intersection of a plane and an axis-aligned bounding box. | 636 |
| 11.68 | We only need to check the corners at the end of the diagonal most closely aligned with the normal to the plane. | 636 |
| 11.69 | The intersection of a plane and an oriented bounding box. | 637 |
| 11.70 | Projecting the diagonal of an OBB onto the plane normal. | 638 |
| 11.71 | Intersection of two axis-aligned bounding boxes. | 639 |
| 11.72 | 2D schematic for OBB intersection detection. After Gottschalk, Lin, and Manocha (1996). | 640 |
| 11.73 | Intersection of an axis-aligned bounding box and a sphere. | 644 |
| 11.74 | Intersection of a linear component and a torus. | 659 |
| 11.75 | Computing the normal of a torus at a point (of intersection). | 660 |
| 11.76 | The u parameter of a point on a torus. | 661 |
| 11.77 | The v parameter of a point on a torus. | 662 |
| 12.1 | Projection of a point onto a plane. | 664 |
| 12.2 | Projection of a vector onto a plane. | 665 |
| 12.3 | Projection of one vector onto another. | 666 |
| 12.4 | Angle between a line and a plane. | 666 |
| 12.5 | Angle between two planes. | 667 |
| 12.6 | Plane normal to a line through a point. | 668 |
| 12.7 | Computing the distance coefficient for the plane. | 669 |

| | | |
|-------|---|-----|
| 12.8 | Three points defining a plane. | 670 |
| 12.9 | Angle between two lines in 3D. | 671 |
| 12.10 | Angle between two lines in 3D, with one line reversed. | 671 |
| 13.1 | BSP tree partitioning of the plane. | 674 |
| 13.2 | A partitioning line for which two coincident edges have opposite direction normals. | 676 |
| 13.3 | A sample polygon for construction of a BSP tree. | 678 |
| 13.4 | Current state after processing edge $\langle 9, 0 \rangle$. | 678 |
| 13.5 | Current state after processing edge $\langle 0, 1 \rangle$. | 679 |
| 13.6 | Current state after processing edge $\langle 1, 2 \rangle$. This edge forces a split of $\langle 4, 5 \rangle$ to $\langle 4, 10 \rangle$ and $\langle 10, 5 \rangle$. It also forces a split of $\langle 8, 9 \rangle$ to $\langle 8, 11 \rangle$ and $\langle 11, 9 \rangle$. | 679 |
| 13.7 | Current state after processing edge $\langle 10, 5 \rangle$. This edge forces a split of $\langle 7, 8 \rangle$ to $\langle 7, 12 \rangle$ and $\langle 12, 8 \rangle$. | 680 |
| 13.8 | Current state after processing edge $\langle 5, 6 \rangle$. | 680 |
| 13.9 | Final state after processing edge $\langle 13, 9 \rangle$. | 681 |
| 13.10 | Partition for a convex polygon and the corresponding BSP tree. | 682 |
| 13.11 | Partition for a convex polygon and the corresponding balanced BSP tree. | 683 |
| 13.12 | Partition of a line segment. | 687 |
| 13.13 | Point-in-convex-polygon test by determining two edges intersected by the vertical line through the test point. P is inside the polygon. Q is outside the polygon. | 700 |
| 13.14 | Point-in-polygon test by counting intersections of ray with polygon. The ray for point P_0 only crosses edges transversely. The number of crossings is odd (5), so the point is inside the polygon. The ray for point P_1 is more complex to analyze. | 701 |
| 13.15 | Points P on the “left” edges of the polygon are classified as inside. Points Q on the “right” edges of the polygon are classified as outside. | 704 |
| 13.16 | Point tags for the horizontal line containing P_1 in Figure 13.14. | 705 |
| 13.17 | Interval tags for the horizontal line containing P_1 in Figure 13.14. | 706 |
| 13.18 | Two configurations for when the test ray $P + t\hat{d}$ intersects a shared edge \vec{e} at an interior edge point. (a) The faces are on the same side of the plane formed by the edge and the ray. Parity is not changed. (b) The faces are on opposite sides. Parity is toggled. | 712 |
| 13.19 | The spherical polygon implied by the edges sharing a vertex V that the test ray intersects. If the point A corresponds to the ray direction, the ray interpenetrates the polyhedron. If the point B corresponds to the ray direction, the ray does not interpenetrate the polyhedron. | 713 |

| | | |
|-------|--|-----|
| 13.20 | Bounded and unbounded polygons that partition the plane into inside and outside regions. The inside region is gray. The unbounded polygon on the right is a half-space with a single line as the boundary of the region. | 714 |
| 13.21 | A polygon and its negation. The inside regions are gray. The edges are shown with the appropriate directions so that the inside is always to the left. | 715 |
| 13.22 | Two polygons whose inside regions are bounded. | 716 |
| 13.23 | The intersection of two polygons shown in gray. | 716 |
| 13.24 | The union of two polygons shown in gray. | 717 |
| 13.25 | The difference of two polygons: (a) The inverted L-shaped polygon minus the pentagon. (b) The pentagon minus the inverted L-shaped polygon. | 718 |
| 13.26 | The exclusive-or of two polygons shown in gray. This polygon is the union of the two differences shown in Figure 13.25. | 719 |
| 13.27 | Intersection of two triangles: (a) The two triangles, A and B . (b) Edges of A intersected with inside of B . (c) Edges of B intersected with inside of A . (d) $A \cap B$ as the collection of all intersected edges. | 721 |
| 13.28 | (a) Two polygons that are reported not to intersect by the pseudocode. (b) The actual set intersection, a line segment. | 722 |
| 13.29 | (a) Two polygons and (b) their true set of intersection. | 723 |
| 13.30 | (a) Polygon with a hole requiring two lists of vertices/edges. (b) Keyhole version to allow a single list of vertices/edges. | 723 |
| 13.31 | Intersection of a rectangle and a keyhole polygon. | 723 |
| 13.32 | (a) Convex. (b) Not convex, since the line segment connecting P and Q is not entirely inside the original set. | 730 |
| 13.33 | A point set and its convex hull. The points are in dark gray, except for those points that became hull vertices, marked in black. The hull is shown in light gray. | 730 |
| 13.34 | A convex hull H , a point V outside H , and the two tangents from V to the hull. The upper and lower tangent points are labeled as P_U and P_L , respectively. | 731 |
| 13.35 | The five possibilities for the relationship of P to a line segment with end points Q_0 and Q_1 : P is (a) to the left of the segment, (b) to the right of the segment, (c) on the line to the left of the segment, (d) on the line to the right of the segment, or (e) on the line and contained by the segment. | 734 |
| 13.36 | Two convex hulls H_L and H_R and their upper and lower tangents. | 740 |
| 13.37 | Two convex hulls H_L and H_R and the incremental search for the lower tangent. | 741 |

| | | |
|-------|--|-----|
| 13.38 | The extreme points used to initialize tangent search are on the same vertical line. The initial visibility tests both do not yield a NEGATIVE test, yet the initial segment connecting the extremes is not a tangent to the hulls. The current candidate for the tangent is shown as a dotted line. | 745 |
| 13.39 | The current hull and point to be merged. The visible faces are drawn in light gray. The hidden faces are drawn in dark gray. The polyline separating the two sets is dashed. The other edges of the visibility cone are dotted. | 746 |
| 13.40 | (a) Two icosahedrons. (b) The merged hull. The dashed lines indicate those edges that are part of faces of the original hulls. The dotted lines indicate those edges that are part of the newly added faces. | 749 |
| 13.41 | (a) A side view of the pyramid and line segment. (b) A view from behind the line segment. The line segment $\langle 0, a \rangle$ can only see triangle $\langle 2, 3, 6 \rangle$ and quadrilateral $\langle 3, 4, 5, 6 \rangle$. The line segment $\langle a, b \rangle$ can only see the quadrilateral. The line segment $\langle b, 1 \rangle$ can only see triangle $\langle 2, 4, 5 \rangle$ and the quadrilateral. The faces that are hidden in all cases are the triangles $\langle 2, 3, 4 \rangle$ and $\langle 2, 5, 6 \rangle$. The terminator consists of the boundaries of these triangles, a sequence of line segments forming two cycles, not a simple cycle. | 750 |
| 13.42 | Triangulations of finite point sets: (a) with optional requirements; (b) without. | 756 |
| 13.43 | The two triangulations for a convex quadrilateral. The angle $\alpha \doteq 0.46$ radians and the angle $\beta \doteq 1.11$ radians. (a) The minimum angle of the top triangle is α (smaller than β). (b) The minimum angle is 2α radians (smaller than β); the triangles maximize the minimum angle. | 757 |
| 13.44 | Two circumcircles for the triangles of Figure 13.43. | 757 |
| 13.45 | (a) The newly inserted point P , shown as an unlabeled black dot, is interior to a triangle, in which case the triangle is split into three subtriangles, or (b) it is on an edge of a triangle, in which case each triangle sharing the edge (if any) is split into two subtriangles. | 760 |
| 13.46 | A triangle pair $\langle T, A \rangle$ that needs an edge swap. The index tracking is necessary so that the correct objects in the vertex-edge-triangle table of the mesh are manipulated. After the edge swap, up to two new pairs of triangles occur, $\langle N_0, B_0 \rangle$ and $\langle N_1, B_1 \rangle$, each pair possibly needing an edge swap. These are pushed onto the stack of pairs that need to be processed. | 762 |
| 13.47 | Supertriangle of the input point set. | 763 |
| 13.48 | Circumcircles containing the next point to be inserted. | 764 |
| 13.49 | The insertion polygon for the next point to be inserted. | 765 |
| 13.50 | The modified insertion polygon that restores the empty circumcircle condition for the total mesh. | 765 |

| | | |
|-------|---|-----|
| 13.51 | The final mesh triangles are dark gray. The removed triangles are shown in light gray. | 766 |
| 13.52 | (a) Convex hull of 2D points lifted onto a paraboloid in 3D. (b) The corresponding Delaunay triangulation, the projection of the lower hull onto the xy -plane. | 767 |
| 13.53 | (a) Two vertices that are visible to each other. The diagonal connecting them is shown. (b) Two vertices that are not visible to each other, blocked by a vertex between them. (c) Two vertices that are not visible to each other, blocked by a single edge. (d) Two vertices that are not visible to each other, blocked by a region outside the polygon. | 768 |
| 13.54 | Illustration of why lack of visibility between V_0 and V_2 is equivalent to triangle $\langle V_0, V_1, V_2 \rangle$ containing a reflex vertex R . | 769 |
| 13.55 | Cone containment (a) for a convex vertex and (b) for a reflex vertex. | 770 |
| 13.56 | A simple polygon that is used to illustrate the horizontal decomposition into trapezoids. The edges are labeled randomly and are processed in that order in the figures that follow. | 776 |
| 13.57 | The entire plane is a single trapezoid. | 777 |
| 13.58 | Split by $s_1.y_0$. | 777 |
| 13.59 | Split by $s_1.y_1$. | 778 |
| 13.60 | Insert s_1 . | 778 |
| 13.61 | Split by $s_2.y_1$. | 779 |
| 13.62 | Split by $s_2.y_0$. | 779 |
| 13.63 | Insert s_2 . | 779 |
| 13.64 | Split by $s_3.y_1$. | 780 |
| 13.65 | Insert s_3 . | 780 |
| 13.66 | Insert s_9 . | 781 |
| 13.67 | The plane after trapezoids are merged into maximally sized ones. | 783 |
| 13.68 | The sample polygon after trapezoids are merged into maximally sized ones. | 784 |
| 13.69 | The sample polygon as a union of monotone polygons. The two polygons are drawn in light gray and dark gray. The horizontal line segments from the trapezoidal decomposition are still shown. | 784 |
| 13.70 | If the triangle at an extreme vertex is an ear, removing the ear yields another monotone polygon. | 785 |
| 13.71 | Failure of triangle $\langle V_0, V_{\min}, V_1 \rangle$ to be an ear. | 786 |
| 13.72 | (a) Not all reflex chain vertices are visible to W . (b) Removal of the triangles leads to W being the next vertex to be added to the reflex chain. | 787 |

| | | |
|-------|---|-----|
| 13.73 | (a) W occurs above the current strip, V_0 is visible to all reflex chain vertices. (b) Removal of the triangles leads to a reduced monotone polygon, so the process can be repeated. | 789 |
| 13.74 | (a) Partition using only vertices. (b) Partition using an additional point interior to the polygon. | 790 |
| 13.75 | Vertex V_0 is reflex. The diagonal $\langle V_0, V_1 \rangle$ is inessential. The diagonal $\langle V_0, V_2 \rangle$ is essential for V_0 . | 791 |
| 13.76 | Original polygon (upper left) and 11 minimum convex decompositions, with the narrowest pairs shaded in gray. A dotted line indicates that the edge of the polygon is treated instead as a diagonal. | 793 |
| 13.77 | Canonical triangulations of the convex polygons in the minimum convex decomposition of a polygon. The original polygon has edges shown in a heavy line. The diagonals used in the decomposition are dotted. The diagonals used in the triangle fans for the canonical triangulations are shown in a light line. | 794 |
| 13.78 | Circumscribed and inscribed circles for a triangle. | 799 |
| 13.79 | Purported minimum-area rectangle that has no coincident polygon edges. | 804 |
| 13.80 | (a) Current bounding circle and a point that is outside the circle, causing the circle to grow. (b) The new bounding circle, but a point inside the old circle is now outside the new circle, causing a restart of the algorithm. | 809 |
| 13.81 | Points U_1 and U_2 chosen for computing Equation 13.2. Only one edge of the triangle is visible to the first point. Two edges of the triangle are visible to the second point. | 819 |
| A.1 | The top sequence shows a nonuniform scale $(x, y) \rightarrow (2x, y)$ applied first, a counterclockwise rotation by $\pi/4$ second. The bottom sequence shows a rotation by any angle (the circle is invariant under rotations), but clearly there is no nonuniform scaling along coordinate axes that can force the circle to become the ellipse of the top sequence. | 856 |
| A.2 | Intersection of a function $f(x, y) = z$ and plane $z = 0.8$ yields a level curve, shown projected on the xy -plane. | 895 |
| A.3 | Level curves for $f(x, y) = \frac{2x^2}{3} + y^2$. | 896 |
| A.4 | Level curves for $f(x, y) = \frac{2x^2}{3} + y^2$, projected onto the xy -plane. | 897 |
| A.5 | Two functions lacking either a minimum or maximum value. | 898 |
| A.6 | Two functions that violate the assumptions of the Extreme Value Theorem. | 899 |
| A.7 | A variety of functions, showing critical points—(a) and (b) are stationary points; (c) and (d) are inflection points. | 900 |

| | | |
|------|---|-----|
| A.8 | The maximum of a function may occur at the boundary of an interval or within the interval. | 901 |
| A.9 | $f(x) = x^3 + 6x^2 - 7x + 19, \forall x \in [-8, 3]$. | 902 |
| A.10 | $f'(x) = 3x^2 + 12x - 7, \forall x \in [-8, 3]$. | 902 |
| A.11 | Relative extrema. | 903 |
| A.12 | Relative extrema of $f(x) = 3x^{\frac{5}{3}} - 15x^{\frac{2}{3}}$. | 904 |
| A.13 | Relative extrema of a function of two variables are the hills and valleys of its graph. | 906 |
| A.14 | The relative maximum of a function $z = f(x, y)$. After Anton (1980). | 907 |
| A.15 | A “saddle function”—the point $(0, 0)$ is not an extremum, in spite of the first partial derivatives being zero. | 909 |
| A.16 | Graph of $2y^2x - yx^2 + 4xy$, showing saddle points and relative minimum. | 911 |
| A.17 | Contour plot of $2y^2x - yx^2 + 4xy$. | 912 |
| A.18 | Plot of the ellipse $17x^2 + 8y^2 + 12xy = 100$. | 915 |
| A.19 | Plot of the function $x^2 + y^2$. | 915 |
| A.20 | Level curves for $x^2 + y^2$. | 916 |
| A.21 | The constraint curve and the ellipse are tangent at the minima. | 916 |
| A.22 | The closest and farthest point to $(1, 2, 2)$ on the sphere $x^2 + y^2 + z^2 = 36$. | 919 |
| A.23 | Constraint equations $g_1(x, y, z) = x - y + z = 1$ and $g_2(x, y, z) = x^2 + y^2 = 1$. | 921 |
| A.24 | Extrema of f shown as points on the constraint curve determined by the intersection of implicit surfaces defined by $g_1 = 0$ and $g_2 = 0$, and the level sets of f at those extrema. | 922 |
| B.1 | Standard terminology for angles. | 924 |
| B.2 | Definition of arc length. | 925 |
| B.3 | Definition of radians. | 925 |
| B.4 | The ratios of sides of a right triangle can be used to define trig functions. | 927 |
| B.5 | Generalized definition for trigonometric functions. | 928 |
| B.6 | Geometrical interpretation of trigonometric functions. | 930 |
| B.7 | Graphs of the fundamental trigonometric functions. | 932 |
| B.8 | The law of sines. | 937 |
| B.9 | Proof of the law of sines. | 938 |
| B.10 | Graphs of the fundamental inverse trigonometric functions. | 947 |