


Hadamard Test

Henry Lin

June 2025

1 Introduction

Hadamard test can be used to find the real or imaginary part of a quantum state. The circuit looks like the following:



1.1 Case 1: Eigenstate of U

Suppose $|\psi\rangle$ is the eigenstate of U with eigenvalue of $e^{i\lambda}$. We can write this as the following:

$$U |\psi\rangle = e^{i\lambda} |\psi\rangle.$$

Now we will analyze the circuit from left to right. At the first stage, we have a Hadamard gate, this will evolve our state as the following:

$$H(|0\rangle \otimes |\psi\rangle) = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \otimes |\psi\rangle.$$

After that we have control-unitary:

$$\begin{aligned} CU\left(\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \otimes |\psi\rangle\right) &= \frac{1}{\sqrt{2}} \cdot (|0\rangle |\psi\rangle + |1\rangle U |\psi\rangle) \\ &= \frac{1}{\sqrt{2}} \cdot (|0\rangle |\psi\rangle + e^{i\lambda} |1\rangle |\psi\rangle) \end{aligned}$$

Lastly, we have a Hadamard gate again:

$$H\left(\frac{1}{\sqrt{2}} \cdot (|0\rangle |\psi\rangle + e^{i\lambda} |1\rangle |\psi\rangle)\right) = \left(\frac{1+e^{i\lambda}}{2} |0\rangle + \frac{1-e^{i\lambda}}{2} |1\rangle\right) \otimes \psi$$

Basically, we have to measured the first ancilla qubit and from there we can estimate what is the value of λ , thus, figure out the eigenvalue of U . We can write the probability of measuring $|0\rangle$ and $|1\rangle$ as the following:

$$p_m = \frac{1 + (-1)^m \cos \lambda}{2}$$

with measurement results 0 or 1. If we want to estimate $\cos \lambda$ with a certain error ϵ , then we only need to sample about a polynomial number of times of the inverse of ϵ by using Chernoff bound.

1.2 Case 2: General Input State

First Hadamard:

$$H(|0\rangle \otimes |\psi\rangle) = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \otimes |\psi\rangle.$$

CU:

$$CU(\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \otimes |\psi\rangle) = \frac{1}{\sqrt{2}} \cdot (|0\rangle |\psi\rangle + |1\rangle U |\psi\rangle)$$

Second Hadamard:

$$\begin{aligned} H(\frac{1}{\sqrt{2}} \cdot (|0\rangle |\psi\rangle + |1\rangle U |\psi\rangle)) &= \frac{(|0\rangle + |1\rangle) |\psi\rangle}{2} + \frac{(|0\rangle - |1\rangle) U |\psi\rangle}{2} \\ &= |0\rangle \frac{I + U}{2} |\psi\rangle + |1\rangle \frac{I - U}{2} |\psi\rangle \end{aligned}$$

Now, let's calculate the probability of measuring $|0\rangle$ for the first ancilla qubit. Let define a projective operator that does this:

$$\begin{aligned} \hat{p}_0 &= |0\rangle \langle 0| \otimes I \\ \hat{p}_1 &= |1\rangle \langle 1| \otimes I \end{aligned}$$

Now, let's use this projective operator to obtain the probabilities:

$$\begin{aligned}
p_0 &= |(|0\rangle\langle 0| \otimes I)[|0\rangle \frac{I+U}{2} |\psi\rangle + |1\rangle \frac{I-U}{2} |\psi\rangle]|^2 \\
&= | |0\rangle \frac{I+U}{2} |\psi\rangle + 0|^2 \\
&= | |0\rangle|^2 | \frac{I+U}{2} |\psi\rangle|^2 \\
&= \frac{\langle \psi | (I+U)(I+U^\dagger) | \psi \rangle}{4} \\
&= \frac{\langle \psi | (2I + U + U^\dagger) | \psi \rangle}{4} \\
&= \frac{1 + \text{Re}\langle U \rangle}{2} \\
p_1 &= \frac{1 - \text{Re}\langle U \rangle}{2}
\end{aligned}$$

This means we can use Hadamard Test to find the real and imaginary part of a quantum state. To obtain the imaginary part, we just have to make sure that the input state to the CU gate is $(|0\rangle - i|1\rangle) \otimes |\psi\rangle$. This can be achieved by applying a S^\dagger after the Hadamard gate.