							Co	nce	WÙ	ty							
/.	Q.	MEGE	ve	up	2 0	dow	n.			V							
		\			,												
		C	mau	ve uy	P			con	con	e di	nun						
2.	D	efin	ition	S													
					a d	iffer	renti	ble	fcu	ctiv	n d	efin	ed	on	an.	inte	val
 	7:	s .		on	·l												
					-		_		^			· · · · ·					
																	11
																e ek	flee
	Pi	מאדנ	al	<i>b C</i>	, (Nhei	n: 7	- C	roing	res	CON	cov	y	ar	۲.		
3.	Th	LOPE	em.														
		_		l l	e c	an i	open	in	teve	al							
											abl	· 7	lunc	tion	. de	fixed	l v
		Ø	lf	V:	x & ,	ℓ , \cdot	f"C>	() >(), i	then	f	方	CÕ	nca	ve i	y o	n s
		Ø	Ц	42	ith	? - {	filx) 20	, t	he.	Ĵ.	15 0	CONC	ove	der	n o	n ,
	2. †	2. D y f 7: 1) 2) 8) P 3. Th	2. Defin Yet f is . 1) Co 2) Cor 3) Le point 3. Theore 1) Le Le Le O	concau 2. Definition Let f f is 1) Concau 2) Concau 3) Let c point at 3. Theorem. 1) Let g O lf	concave up 2. Definitions Let f be f is on 1. Concave - 2. Concave - 3. Let c b point at c 3. Theorem. 1. Let l b Let f b Olf vi	concave up 2. Definitions Let f be a d f is on l 1) Concave - up 2) Concave - don 3) Let c be a point at c, Let f be a Of the a Of	concave up 2. Definitions Let of be a difference of the concave - up: us 2) Concave - clown: 3) Let c be an expoint at c, when 3. Theorem. 1) Let l be an exposed of the concave of	Concave up & down: concave up 2. Definitions Let f be a differential f is on l: 13 Concave - up: when 23 Concave - down: when 33 Let c be an inte point at c, when: 7 3. Theorem: 13 Let l be an open Let f be a twice Of Tx & l, f"l;	Concave up & doron. concave up con 2. Definitions Let f be a differentible f is on l: 1) Concave - up: when f' 2) Concave - down: when f 3) Let c be an interior point at c, when: f c 3. Theorem. 1) Let l be an open in Let f be a twice - de o lf \(\frac{1}{2} \times 6 \), \(\frac{1}{1} \times 1 \)	Concave up & down. Concave up concave 2. Definitions Let f be a differentible four f is on l: 10 Concave - up: when f' is 20 Concave - down: when f' 30 Let c be an interior po point at c, when: f chang 3. Theorem. 10 Let l be an open interior Let f be a twice - differentible Of It x & l. f''(x) 70, in	concave up concave de 2. Definitions Let f be a differentible function f is on l: 1> Concave - up: when f' is in 2) Concave - down: when f' is a 3) Let c be an interior point point at c, when: f changes 3. Theorem. 1> Let l be an open inteval Let f be a twice-different Olf Tx & l, f''(x) 70, then	Concave up & down. 2. Definitions Let of be a differentible function of its	1. Concave up & down. concave up concave down. 2. Definitions Let f be a differentible function define f is on l: 1. Concave - up: when f' is increasing 2. Concave - down: when f' is decrease 3. Let c be an interior point to l. point at c, when: f changes concav. 3. Theorem. 1. Let l be an open interior Let f be a twice - differentiable of Of Is & l. f''(x) >0, then f is	Concave up & doron. Concave up concave down. 2. Definitions Let f be a differentible function defined of its on l: 1. Concave - up: when f' its increasing it is concave - down: when f' its obscreasing it is yet c be an interior point to l. f point at c, when: f changes concavity 3. Theorem. 1. Let l be an open interval Let f be a twice - differentiable function of the first concavity.	Concave up & down. Concave up concave clown. 2. Definitions Yet of be a differentible function defined on a fix increasing on l: 1. Concave - up: when f' is increasing on . 2. Yet c be an interior point to l. of how point at c, when f changes concavity at 3. Theorem. 1. Yet l be an open interal Yet of be a twice-differentiable function of the file of the concavity.	Concave up concave down. 2. Definitions Yet of be a differentible function defined on an of the concave of the increasing on l. 1. Concave - up: when f' is increasing on l. 2. Stoncave - down: when f' is obscreasing on l. 2. Yet c be an interior point to l. of has an point at c, when: of changes concavity at c. 3. Theorem. 1. Yet l be an open interal Yet of be a twice - differentiable function de of the l. of the concave in the left of the l	Concave up & down. 2. Definitions Let f be a differentible function defined on an inte f is on l: 1. Concave - up: when f' is increasing on l. 2. Loncave - down: when f' is obscreasing on l. 3. Let c be an interior point to l. f has an in point at c, when: f changes concavity at c. 3. Theorem.

