

# Chapter 1 Solution

January 16, 2025

## Solution to Exercise 1.1

### Part 1: Prove that $B_1, B_2, \dots$ are disjoint.

Let us consider  $B_i$  and  $B_j$  where  $i < j$ . By the definition of  $B_j$ , any  $\omega \in B_j$  satisfies  $\omega \notin A_i$ . On the other hand, by the definition of  $B_i$ , any  $\omega \in B_i$  satisfies  $\omega \in A_i$ . Therefore,  $\omega \notin B_i$  if  $\omega \in B_j$ . Similarly,  $\omega \notin B_j$  if  $\omega \in B_i$ . Hence,  $B_i \cap B_j = \emptyset$  for  $i \neq j$ , proving that  $B_1, B_2, \dots$  are disjoint.

### Part 2: Prove that $\bigcup_{i=1}^n B_i = \bigcup_{i=1}^n A_i$ .

1. **Direction 1:**  $\bigcup_{i=1}^n B_i \subseteq \bigcup_{i=1}^n A_i$  For any element  $\omega \in \bigcup_{i=1}^n B_i$ , it must belong to exactly one  $B_i$  because of the disjoint property proved above. By the definition of  $B_i$ , this implies  $\omega \in A_i$ . Therefore,  $\omega \in \bigcup_{i=1}^n A_i$ .
2. **Direction 2:**  $\bigcup_{i=1}^n A_i \subseteq \bigcup_{i=1}^n B_i$  For any  $\omega \in \bigcup_{i=1}^n A_i$ , we can find the smallest index  $j$  such that  $\omega \in A_j$ . If  $j = 1$ , then  $\omega \in B_1$  by definition. If  $j > 1$ ,  $\omega \notin A_{j-1}$  (since  $j$  is the smallest index), so  $\omega \in B_j$ . Thus,  $\omega \in \bigcup_{i=1}^n B_i$ .

By both directions, we conclude  $\bigcup_{i=1}^n B_i = \bigcup_{i=1}^n A_i$ .

### Part 3: Prove monotone decreasing behavior.

To construct disjoint sets  $B_i$  in the case of a monotone decreasing sequence of sets  $A_1 \supseteq A_2 \supseteq \dots \supseteq A_n$ , we define  $B_1 = A_1$  and  $B_i = A_i \setminus A_{i-1}$  for  $i > 1$ . These  $B_i$  are disjoint by construction.

By the same logic as in Part 2, we can prove that  $\bigcup_{i=1}^n B_i = \bigcup_{i=1}^n A_i$ . With the disjoint property and axiom 3 (countable additivity), we reach the same conclusion for this case.

## Solution to Exercise 1.11

### Problem

Suppose that A and B are independent events. Show that  $A^c$  and  $B^c$  are independent events

### Solution

We use complement rule in property 1.1 and union formula in lemma 1.6 in the proof below.

$$\begin{aligned} P(A^c)P(B^c) &= (1 - P(A))(1 - P(B)) && \text{(Property 1.1)} \\ &= 1 - (P(A) + P(B) - P(AB)) && \text{(Lemma 1.6)} \\ &= 1 - P(A \cup B) && \text{(Property 1.1)} \\ &= P((A \cup B)^c) && \text{(De Morgan's Law)} \\ &= P(A^c \cap B^c) \end{aligned}$$