

# Lecture 1 — Notation, Functions, and Limits

Henry Watson

Georgetown University

8/14/23

# Today's Agenda

# Today's Agenda

## ① Introductions

# Today's Agenda

- 1 Introductions
- 2 Math Camp Logistics and schedule

# Today's Agenda

- 1 Introductions
- 2 Math Camp Logistics and schedule
- 3 Variables and measurement

# Today's Agenda

- 1 Introductions
- 2 Math Camp Logistics and schedule
- 3 Variables and measurement
- 4 Algebra

# Today's Agenda

- 1 Introductions
- 2 Math Camp Logistics and schedule
- 3 Variables and measurement
- 4 Algebra
- 5 Functions

# Today's Agenda

- 1 Introductions
- 2 Math Camp Logistics and schedule
- 3 Variables and measurement
- 4 Algebra
- 5 Functions
- 6 Series, sequences, and limits



# Week's Agenda

# Week's Agenda

- Lecture 1: Notation, functions and limits

# Week's Agenda

- Lecture 1: Notation, functions and limits
- Lecture 2: Linear Algebra

# Week's Agenda

- Lecture 1: Notation, functions and limits
- Lecture 2: Linear Algebra
- Lecture 3: Calculus 1 - Derivatives

# Week's Agenda

- Lecture 1: Notation, functions and limits
- Lecture 2: Linear Algebra
- Lecture 3: Calculus 1 - Derivatives
- Lecture 4: Calculus 2 - Integrals and multivariate calculus

# Week's Agenda

- Lecture 1: Notation, functions and limits
- Lecture 2: Linear Algebra
- Lecture 3: Calculus 1 - Derivatives
- Lecture 4: Calculus 2 - Integrals and multivariate calculus
- Lecture 5: Probability

# Schedule

	Monday 14th	Tuesday 15th	Wednesday 16th	Thursday 17th	Friday 18th
9:30 - 12:00	Notation, functions, and limits	Linear algebra	Calculus 1	Calculus 2	Probability
12:00 - 12:10	Break	Break	Break	Break	Break
12:10 - 1:00	Problem Set	Problem Set	Problem Set	Lunch Break	LaTeX
1:00 - 2:00	Lunch Break	Lunch Break	Lunch Break	Problem Set	Lunch Break
2:00 - 2:30	Review PS	Review PS	Review PS	Review PS	Q&A with Prof Klasnja
2:30 - 3:30	Software Installation	R - Day 1	R - Day 2	R - Day 3	Q&A Continued

# Goals of Math Camp



# Goals of Math Camp

- Familiarity

# Goals of Math Camp

- Familiarity
- Recognition

# Goals of Math Camp

- Familiarity
- Recognition
- Confidence

# What is quantitative methods?

# What is quantitative methods?

- Ultimate goal is to improve our understanding of the world

# What is quantitative methods?

- Ultimate goal is to improve our understanding of the world
- We use quantitative methods to test theories regarding the relationships between concepts

# What is quantitative methods?

- Ultimate goal is to improve our understanding of the world
- We use quantitative methods to test theories regarding the relationships between concepts
- The goal of a “causal claim” is rarely achieved in full, but we can make compelling arguments

# What is quantitative methods?

- Ultimate goal is to improve our understanding of the world
- We use quantitative methods to test theories regarding the relationships between concepts
- The goal of a “causal claim” is rarely achieved in full, but we can make compelling arguments
- Qualitative methods can answer “how” and “why” questions



# Variables and Constants

# Variables and Constants

- **Theory:** A set of statements that involve concepts.

# Variables and Constants

- **Theory:** A set of statements that involve concepts.
- **Concept:** abstract ideas used to understand the world

# Variables and Constants

- **Theory:** A set of statements that involve concepts.
- **Concept:** abstract ideas used to understand the world
  - Participation, voting, democracy, war

# Variables and Constants

- **Theory:** A set of statements that involve concepts.
- **Concept:** abstract ideas used to understand the world
  - Participation, voting, democracy, war
- **Measure:** an operational indicator of a concept

# Variables and Constants

- **Theory:** A set of statements that involve concepts.
- **Concept:** abstract ideas used to understand the world
  - Participation, voting, democracy, war
- **Measure:** an operational indicator of a concept
- **Variable:** A concept or measure that takes different values in a given set.

# Variables and Constants

- **Theory:** A set of statements that involve concepts.
- **Concept:** abstract ideas used to understand the world
  - Participation, voting, democracy, war
- **Measure:** an operational indicator of a concept
- **Variable:** A concept or measure that takes different values in a given set.
- **Constant:** A concept or measure that has a single value for a given set.

# Sets and Sample Spaces



# Sets and Sample Spaces

- A *set* is a collection of elements.

# Sets and Sample Spaces

- A *set* is a collection of elements.
- **Common sets:** Natural numbers ( $\mathbb{N}$ ), Integers ( $\mathbb{Z}$ ), Rational numbers ( $\mathbb{Q}$ ), Real Numbers ( $\mathbb{R}$ ), etc.

# Sets and Sample Spaces

- A *set* is a collection of elements.
- **Common sets:** Natural numbers ( $\mathbb{N}$ ), Integers ( $\mathbb{Z}$ ), Rational numbers ( $\mathbb{Q}$ ), Real Numbers ( $\mathbb{R}$ ), etc.
- A set can be:

# Sets and Sample Spaces

- A *set* is a collection of elements.
- **Common sets:** Natural numbers ( $\mathbb{N}$ ), Integers ( $\mathbb{Z}$ ), Rational numbers ( $\mathbb{Q}$ ), Real Numbers ( $\mathbb{R}$ ), etc.
- A set can be:
  - **Finite or infinite:**  $\mathbb{Z}$  is infinite, but all the integers from 1 to 10 is finite.

# Sets and Sample Spaces

- A *set* is a collection of elements.
- **Common sets:** Natural numbers ( $\mathbb{N}$ ), Integers ( $\mathbb{Z}$ ), Rational numbers ( $\mathbb{Q}$ ), Real Numbers ( $\mathbb{R}$ ), etc.
- A set can be:
  - **Finite or infinite:**  $\mathbb{Z}$  is infinite, but all the integers from 1 to 10 is finite.
  - **Countable or uncountable:** a countable set is one whose each of its element can be associated with a natural number (or an integer).

# Sets and Sample Spaces

- A *set* is a collection of elements.
- **Common sets:** Natural numbers ( $\mathbb{N}$ ), Integers ( $\mathbb{Z}$ ), Rational numbers ( $\mathbb{Q}$ ), Real Numbers ( $\mathbb{R}$ ), etc.
- A set can be:
  - **Finite or infinite:**  $\mathbb{Z}$  is infinite, but all the integers from 1 to 10 is finite.
  - **Countable or uncountable:** a countable set is one whose each of its element can be associated with a natural number (or an integer).
  - **Bounded or unbounded:** A bounded set has finite size (but may have infinite elements).

# Sets and Sample Spaces

- A *set* is a collection of elements.
- **Common sets:** Natural numbers ( $\mathbb{N}$ ), Integers ( $\mathbb{Z}$ ), Rational numbers ( $\mathbb{Q}$ ), Real Numbers ( $\mathbb{R}$ ), etc.
- A set can be:
  - **Finite or infinite:**  $\mathbb{Z}$  is infinite, but all the integers from 1 to 10 is finite.
  - **Countable or uncountable:** a countable set is one whose each of its element can be associated with a natural number (or an integer).
  - **Bounded or unbounded:** A bounded set has finite size (but may have infinite elements).
- Some important sets that we are going to use as political scientists:

# Sets and Sample Spaces

- A *set* is a collection of elements.
- **Common sets:** Natural numbers ( $\mathbb{N}$ ), Integers ( $\mathbb{Z}$ ), Rational numbers ( $\mathbb{Q}$ ), Real Numbers ( $\mathbb{R}$ ), etc.
- A set can be:
  - **Finite or infinite:**  $\mathbb{Z}$  is infinite, but all the integers from 1 to 10 is finite.
  - **Countable or uncountable:** a countable set is one whose each of its element can be associated with a natural number (or an integer).
  - **Bounded or unbounded:** A bounded set has finite size (but may have infinite elements).
- Some important sets that we are going to use as political scientists:
  - **Solution set:** a set that contains all solutions for an equation



# Sets and Sample Spaces

- A *set* is a collection of elements.
- **Common sets:** Natural numbers ( $\mathbb{N}$ ), Integers ( $\mathbb{Z}$ ), Rational numbers ( $\mathbb{Q}$ ), Real Numbers ( $\mathbb{R}$ ), etc.
- A set can be:
  - **Finite or infinite:**  $\mathbb{Z}$  is infinite, but all the integers from 1 to 10 is finite.
  - **Countable or uncountable:** a countable set is one whose each of its element can be associated with a natural number (or an integer).
  - **Bounded or unbounded:** A bounded set has finite size (but may have infinite elements).
- Some important sets that we are going to use as political scientists:
  - **Solution set:** a set that contains all solutions for an equation
  - **Sample space:** a set that contains all values that a variable can take.

# Unions and Intersections

# Unions and Intersections

- Much as sets contain elements, they also can contain, and be contained by, other sets.

# Unions and Intersections

- Much as sets contain elements, they also can contain, and be contained by, other sets.
- Notation:

# Unions and Intersections

- Much as sets contain elements, they also can contain, and be contained by, other sets.
- Notation:
  - $A \subset B$ : “A is a **proper subset** of B” implies that set B contains all the elements in A, plus at least one more

# Unions and Intersections

- Much as sets contain elements, they also can contain, and be contained by, other sets.
- Notation:
  - $A \subset B$ : “A is a **proper subset** of B” implies that set B contains all the elements in A, plus at least one more
  - $A \subseteq B$ : “A is a **subset** of B”. In this case, it allows A and B to be the same.

# Unions and Intersections

- Much as sets contain elements, they also can contain, and be contained by, other sets.
- Notation:
  - $A \subset B$ : “A is a **proper subset** of B” implies that set B contains all the elements in A, plus at least one more
  - $A \subseteq B$ : “A is a **subset** of B”. In this case, it allows A and B to be the same.
- **Intersection:**  $A \cap B$ . The set of elements common to two sets.

# Unions and Intersections


- Much as sets contain elements, they also can contain, and be contained by, other sets.
- Notation:
  - $A \subset B$ : “A is a **proper subset** of B” implies that set B contains all the elements in A, plus at least one more
  - $A \subseteq B$ : “A is a **subset** of B”. In this case, it allows A and B to be the same.
- **Intersection:**  $A \cap B$ . The set of elements common to two sets.
- **Union:**  $A \cup B$ . The set that contains all elements in both sets.




# Unions and Intersections

- Much as sets contain elements, they also can contain, and be contained by, other sets.
- Notation:
  - $A \subset B$ : “A is a **proper subset** of B” implies that set B contains all the elements in A, plus at least one more
  - $A \subseteq B$ : “A is a **subset** of B”. In this case, it allows A and B to be the same.
- **Intersection**:  $A \cap B$ . The set of elements common to two sets.
- **Union**:  $A \cup B$ . The set that contains all elements in both sets.
- **Mutually exclusive sets**: the intersection is the empty set.


# $A \subset B$ (Proper Subset)




# $A \cap B$ (Intersection)



# $A \cup B$ (Union)



$A \cap B = \emptyset$  (Mutually Exclusive)



# Levels of Measurement

# Levels of Measurement

- Categorical

# Levels of Measurement

- Categorical
  - **Nominal**: No mathematical relationship; difference of kind.



# Levels of Measurement

- Categorical
  - **Nominal**: No mathematical relationship; difference of kind.
    - Occupation, colors, political party, gender

# Levels of Measurement

- Categorical
  - **Nominal:** No mathematical relationship; difference of kind.
    - Occupation, colors, political party, gender
  - **Ordinal:** Meaningful order, no meaningful “distance” between categories.

# Levels of Measurement

- Categorical
  - **Nominal:** No mathematical relationship; difference of kind.
    - Occupation, colors, political party, gender
  - **Ordinal:** Meaningful order, no meaningful “distance” between categories.
    - Ideology, agree/disagree, education level

# Levels of Measurement

- Categorical
  - **Nominal:** No mathematical relationship; difference of kind.
    - Occupation, colors, political party, gender
  - **Ordinal:** Meaningful order, no meaningful “distance” between categories.
    - Ideology, agree/disagree, education level
- Numerical

# Levels of Measurement

- Categorical
  - **Nominal:** No mathematical relationship; difference of kind.
    - Occupation, colors, political party, gender
  - **Ordinal:** Meaningful order, no meaningful “distance” between categories.
    - Ideology, agree/disagree, education level
- Numerical
  - **Interval:** Meaningful order and distance between values.

# Levels of Measurement

- Categorical
  - **Nominal:** No mathematical relationship; difference of kind.
    - Occupation, colors, political party, gender
  - **Ordinal:** Meaningful order, no meaningful “distance” between categories.
    - Ideology, agree/disagree, education level
- Numerical
  - **Interval:** Meaningful order and distance between values.
    - Age, budget, polity scores, vote share, feeling thermometer

# Levels of Measurement

- Categorical
  - **Nominal:** No mathematical relationship; difference of kind.
    - Occupation, colors, political party, gender
  - **Ordinal:** Meaningful order, no meaningful “distance” between categories.
    - Ideology, agree/disagree, education level
- Numerical
  - **Interval:** Meaningful order and distance between values.
    - Age, budget, polity scores, vote share, feeling thermometer
    - Discrete: countable

# Levels of Measurement

- Categorical
  - **Nominal:** No mathematical relationship; difference of kind.
    - Occupation, colors, political party, gender
  - **Ordinal:** Meaningful order, no meaningful “distance” between categories.
    - Ideology, agree/disagree, education level
- Numerical
  - **Interval:** Meaningful order and distance between values.
    - Age, budget, polity scores, vote share, feeling thermometer
    - Discrete: countable
    - Continuous: non-countable



# Levels of Measurement

- Categorical
  - **Nominal:** No mathematical relationship; difference of kind.
    - Occupation, colors, political party, gender
  - **Ordinal:** Meaningful order, no meaningful “distance” between categories.
    - Ideology, agree/disagree, education level
- Numerical
  - **Interval:** Meaningful order and distance between values.
    - Age, budget, polity scores, vote share, feeling thermometer
    - Discrete: countable
    - Continuous: non-countable
    - Ratio: meaningful zero (re-scaled thermometer)

# Why it Matters

# Why it Matters

- Visualizaton
  - Categorical: pie charts, bar charts
  - Numerical (especially Continuous): histograms, box plots

# Why it Matters

- Visualization
  - Categorical: pie charts, bar charts
  - Numerical (especially Continuous): histograms, box plots
- Analysis
  - Numerical data can use summary statistics (mean, median, standard deviation)
  - Categorical - Categorical : cross-tabulation
  - Categorical - Continuous : difference in means
  - Continuous - Continuous : scatter plots, regression

# Basic properties of arithmetic

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$



# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Commutative properties:**

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Commutative properties:**
  - $a + b = b + a$

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Commutative properties:**
  - $a + b = b + a$
  - $a \cdot b = b \cdot a$

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Commutative properties:**
  - $a + b = b + a$
  - $a \cdot b = b \cdot a$
- **Distributive properties:**

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Commutative properties:**
  - $a + b = b + a$
  - $a \cdot b = b \cdot a$
- **Distributive properties:**
  - $a \cdot (b + c) = ab + ac$

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Commutative properties:**
  - $a + b = b + a$
  - $a \cdot b = b \cdot a$
- **Distributive properties:**
  - $a \cdot (b + c) = ab + ac$
- **Identity properties:**

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Commutative properties:**
  - $a + b = b + a$
  - $a \cdot b = b \cdot a$
- **Distributive properties:**
  - $a \cdot (b + c) = ab + ac$
- **Identity properties:**
  - $a + 0 = a$



# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Commutative properties:**
  - $a + b = b + a$
  - $a \cdot b = b \cdot a$
- **Distributive properties:**
  - $a \cdot (b + c) = ab + ac$
- **Identity properties:**
  - $a + 0 = a$
  - $a \cdot 1 = a$

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Commutative properties:**
  - $a + b = b + a$
  - $a \cdot b = b \cdot a$
- **Distributive properties:**
  - $a \cdot (b + c) = ab + ac$
- **Identity properties:**
  - $a + 0 = a$
  - $a \cdot 1 = a$
- **Inverse properties (for real numbers not integers):**

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Commutative properties:**
  - $a + b = b + a$
  - $a \cdot b = b \cdot a$
- **Distributive properties:**
  - $a \cdot (b + c) = ab + ac$
- **Identity properties:**
  - $a + 0 = a$
  - $a \cdot 1 = a$
- **Inverse properties (for real numbers not integers):**
  - $(-a) + a = 0$

# Basic properties of arithmetic

- For variables that stand for real numbers or integers, these properties will always hold:
- **Associative properties:**
  - $(a + b) + c = a + (b + c)$
  - $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Commutative properties:**
  - $a + b = b + a$
  - $a \cdot b = b \cdot a$
- **Distributive properties:**
  - $a \cdot (b + c) = ab + ac$
- **Identity properties:**
  - $a + 0 = a$
  - $a \cdot 1 = a$
- **Inverse properties (for real numbers not integers):**
  - $(-a) + a = 0$
  - $a^{-1} \cdot a = 1$

# Order of operations

# Order of operations

- Order of operations - PEMDAS:

# Order of operations

- **Order of operations - PEMDAS:**
  - Parentheses ()

# Order of operations

- **Order of operations - PEMDAS:**
  - Parentheses ( $()$ )
  - Exponents ( $^x$ )



# Order of operations

- **Order of operations - PEMDAS:**
  - Parentheses ( $()$ )
  - Exponents ( $^x$ )
  - Multiplication ( $\cdot$ )

# Order of operations

- **Order of operations - PEMDAS:**
  - Parentheses ( $()$ )
  - Exponents ( $^x$ )
  - Multiplication ( $\cdot$ )
  - Division ( $\div$ )

# Order of operations

- **Order of operations - PEMDAS:**
  - Parentheses ( $()$ )
  - Exponents ( $^x$ )
  - Multiplication ( $\cdot$ )
  - Division ( $\div$ )
  - Addition ( $+$ )

# Order of operations

- **Order of operations - PEMDAS:**
  - Parentheses ( $()$ )
  - Exponents ( $^x$ )
  - Multiplication ( $\cdot$ )
  - Division ( $\div$ )
  - Addition ( $+$ )
  - Subtraction ( $-$ )

# Order of operations

- **Order of operations - PEMDAS:**
  - Parentheses ( $()$ )
  - Exponents ( $^x$ )
  - Multiplication ( $\cdot$ )
  - Division ( $\div$ )
  - Addition ( $+$ )
  - Subtraction ( $-$ )

# Inequalities

# Inequalities

- All pairs of real numbers have exactly one of the following relations:  
 $x = y$ ,  $x > y$ , or  $x < y$ .

# Inequalities

- All pairs of real numbers have exactly one of the following relations:  
 $x = y$ ,  $x > y$ , or  $x < y$ .
- Solving inequalities is similar to solving equations but there are a few extra properties



# Inequalities

- All pairs of real numbers have exactly one of the following relations:  
 $x = y$ ,  $x > y$ , or  $x < y$ .
- Solving inequalities is similar to solving equations but there are a few extra properties
- Adding any number to each side of these relations will not change them; this includes inequalities.

# Inequalities

- All pairs of real numbers have exactly one of the following relations:  
 $x = y$ ,  $x > y$ , or  $x < y$ .
- Solving inequalities is similar to solving equations but there are a few extra properties
- Adding any number to each side of these relations will not change them; this includes inequalities.
- Multiplication:

# Inequalities

- All pairs of real numbers have exactly one of the following relations:  $x = y$ ,  $x > y$ , or  $x < y$ .
- Solving inequalities is similar to solving equations but there are a few extra properties
- Adding any number to each side of these relations will not change them; this includes inequalities.
- Multiplication:
  - If  $a$  is positive and  $x > y$ , then  $ax > ay$ .

# Inequalities

- All pairs of real numbers have exactly one of the following relations:  
 $x = y$ ,  $x > y$ , or  $x < y$ .
- Solving inequalities is similar to solving equations but there are a few extra properties
- Adding any number to each side of these relations will not change them; this includes inequalities.
- Multiplication:
  - If  $a$  is positive and  $x > y$ , then  $ax > ay$ .
  - If  $a$  is negative and  $x > y$ , then  $ax < ay$ .

# Inequalities

- All pairs of real numbers have exactly one of the following relations:  
 $x = y$ ,  $x > y$ , or  $x < y$ .
- Solving inequalities is similar to solving equations but there are a few extra properties
- Adding any number to each side of these relations will not change them; this includes inequalities.
- Multiplication:
  - If  $a$  is positive and  $x > y$ , then  $ax > ay$ .
  - If  $a$  is negative and  $x > y$ , then  $ax < ay$ .
- Division:

# Inequalities

- All pairs of real numbers have exactly one of the following relations:  
 $x = y$ ,  $x > y$ , or  $x < y$ .
- Solving inequalities is similar to solving equations but there are a few extra properties
- Adding any number to each side of these relations will not change them; this includes inequalities.
- Multiplication:
  - If  $a$  is positive and  $x > y$ , then  $ax > ay$ .
  - If  $a$  is negative and  $x > y$ , then  $ax < ay$ .
- Division:
  - If  $a$  is positive and  $x > y$ , then  $\frac{x}{a} > \frac{y}{a}$ .

# Inequalities

- All pairs of real numbers have exactly one of the following relations:  
 $x = y$ ,  $x > y$ , or  $x < y$ .
- Solving inequalities is similar to solving equations but there are a few extra properties
- Adding any number to each side of these relations will not change them; this includes inequalities.
- Multiplication:
  - If  $a$  is positive and  $x > y$ , then  $ax > ay$ .
  - If  $a$  is negative and  $x > y$ , then  $ax < ay$ .
- Division:
  - If  $a$  is positive and  $x > y$ , then  $\frac{x}{a} > \frac{y}{a}$ .
  - If  $a$  is negative and  $x > y$ , then  $\frac{x}{a} < \frac{y}{a}$ .

# Exponents, logarithms, and roots



# Exponents, logarithms, and roots

- Exponential:  $x^a$

# Exponents, logarithms, and roots

- Exponential:  $x^a$
- Roots:  $\sqrt[a]{x}$ 
  - All roots can be expressed as fractional exponents
  - $\sqrt[a]{x} = x^{1/a}$
- Logarithm: inverse of exponents  $\log_a x$ 
  - To what power would you need to raise  $a$  to to get  $x$ ?

# Exponents, logarithms, and roots

- Exponential:  $x^a$
- Roots:  $\sqrt[a]{x}$ 
  - All roots can be expressed as fractional exponents
  - $\sqrt[a]{x} = x^{1/a}$
- Logarithm: inverse of exponents  $\log_a x$ 
  - To what power would you need to raise  $a$  to to get  $x$ ?
- Natural log: log base  $e$ 
  - $e \approx 2.7183$
  - Useful properties in calculus
  - Taking the natural log of a variable allows us to interpret relationships in terms of percentage changes

# Exponent, logarithms and root rules

## Exponent rules

# Exponent, logarithms and root rules

## Exponent rules

- $x^a \cdot x^b = x^{a+b}$

# Exponent, logarithms and root rules

## Exponent rules

- $x^a \cdot x^b = x^{a+b}$
- $x^a \cdot z^a = (xz)^a$

# Exponent, logarithms and root rules

## Exponent rules

- $x^a \cdot x^b = x^{a+b}$
- $x^a \cdot z^a = (xz)^a$
- $(x^a)^b = x^{ab}$

# Exponent, logarithms and root rules

## Exponent rules

- $x^a \cdot x^b = x^{a+b}$
- $x^a \cdot z^a = (xz)^a$
- $(x^a)^b = x^{ab}$
- $\frac{x^a}{x^b} = x^{a-b}$



# Exponent, logarithms and root rules

## Exponent rules

- $x^a \cdot x^b = x^{a+b}$
- $x^a \cdot z^a = (xz)^a$
- $(x^a)^b = x^{ab}$
- $\frac{x^a}{x^b} = x^{a-b}$
- $\frac{x^a}{z^a} = \left(\frac{x}{z}\right)^a$

# Exponent, logarithms and root rules

## Exponent rules

- $x^a \cdot x^b = x^{a+b}$
- $x^a \cdot z^a = (xz)^a$
- $(x^a)^b = x^{ab}$
- $\frac{x^a}{x^b} = x^{a-b}$
- $\frac{x^a}{z^a} = \left(\frac{x}{z}\right)^a$

## Logarithm rules

# Exponent, logarithms and root rules

## Exponent rules

- $x^a \cdot x^b = x^{a+b}$
- $x^a \cdot z^a = (xz)^a$
- $(x^a)^b = x^{ab}$
- $\frac{x^a}{x^b} = x^{a-b}$
- $\frac{x^a}{z^a} = \left(\frac{x}{z}\right)^a$

## Logarithm rules

- $\log(x_1 \cdot x_2) = \log(x_1) + \log(x_2)$   
for  $x_1, x_2 > 0$

# Exponent, logarithms and root rules

## Exponent rules

- $x^a \cdot x^b = x^{a+b}$
- $x^a \cdot z^a = (xz)^a$
- $(x^a)^b = x^{ab}$
- $\frac{x^a}{x^b} = x^{a-b}$
- $\frac{x^a}{z^a} = \left(\frac{x}{z}\right)^a$

## Logarithm rules

- $\log(x_1 \cdot x_2) = \log(x_1) + \log(x_2)$   
for  $x_1, x_2 > 0$
- $\log\left(\frac{x_1}{x_2}\right) = \log(x_1) - \log(x_2)$   
for  $x_1, x_2 > 0$

# Exponent, logarithms and root rules

## Exponent rules

- $x^a \cdot x^b = x^{a+b}$
- $x^a \cdot z^a = (xz)^a$
- $(x^a)^b = x^{ab}$
- $\frac{x^a}{x^b} = x^{a-b}$
- $\frac{x^a}{z^a} = \left(\frac{x}{z}\right)^a$

## Logarithm rules

- $\log(x_1 \cdot x_2) = \log(x_1) + \log(x_2)$   
for  $x_1, x_2 > 0$
- $\log\left(\frac{x_1}{x_2}\right) = \log(x_1) - \log(x_2)$   
for  $x_1, x_2 > 0$
- $\log(x^b) = b \cdot \ln(x)$   
for  $x > 0$

# Exponent, logarithms and root rules

## Root rules

- $\sqrt{x} = x^{\frac{1}{2}}$
- $\sqrt[n]{x} \cdot \sqrt[n]{z} = \sqrt[n]{xz}$   
for  $n > 1$
- $\frac{\sqrt[n]{x}}{\sqrt[n]{z}} = \sqrt[n]{\frac{x}{z}}$   
for  $n > 1$

# Functions and its characteristics

# Functions and its characteristics

- Functions provide a **specific** description of the association or relationship between two (or among several) concepts (in theoretical work) or variables (in empirical work)



# Functions and its characteristics

- Functions provide a **specific** description of the association or relationship between two (or among several) concepts (in theoretical work) or variables (in empirical work)
- Functions assign one element of the range to an element of the domain (one  $x$  is assigned to one  $y$ )

# Functions and its characteristics

- Functions provide a **specific** description of the association or relationship between two (or among several) concepts (in theoretical work) or variables (in empirical work)
- Functions assign one element of the range to an element of the domain (one  $x$  is assigned to one  $y$ )
- Noted as  $f(x) : A \rightarrow B$  or “ $f$  maps  $A$  into  $B$ ”

# Functions and its characteristics

- Functions provide a **specific** description of the association or relationship between two (or among several) concepts (in theoretical work) or variables (in empirical work)
- Functions assign one element of the range to an element of the domain (one  $x$  is assigned to one  $y$ )
- Noted as  $f(x) : A \rightarrow B$  or “ $f$  maps  $A$  into  $B$ ”
- $A$  is the *domain*, or set of possible  $x$  values.

# Functions and its characteristics

- Functions provide a **specific** description of the association or relationship between two (or among several) concepts (in theoretical work) or variables (in empirical work)
- Functions assign one element of the range to an element of the domain (one  $x$  is assigned to one  $y$ )
- Noted as  $f(x) : A \rightarrow B$  or “ $f$  maps  $A$  into  $B$ ”
- $A$  is the *domain*, or set of possible  $x$  values.
- $B$  is the *codomain*, or set of possible  $y$  values.

# Functions and its characteristics

- Functions provide a **specific** description of the association or relationship between two (or among several) concepts (in theoretical work) or variables (in empirical work)
- Functions assign one element of the range to an element of the domain (one  $x$  is assigned to one  $y$ )
- Noted as  $f(x) : A \rightarrow B$  or “ $f$  maps  $A$  into  $B$ ”
- $A$  is the *domain*, or set of possible  $x$  values.
- $B$  is the *codomain*, or set of possible  $y$  values.
- To be a function, there must only be one unique value in its range ( $y$ ) for each value in its domain ( $x$ )

# Graph examples

In which of these graphs can we observe a function?




Figure: a)






Figure: b)



# Graph examples

In which of these graphs can we observe a function?




Figure: a)






Figure: b)



# Function composition



# Function composition

- We can chain multiple functions using function composition.

# Function composition

- We can chain multiple functions using function composition.
  - This is written either as  $g \circ f(x)$  or  $g(f(x))$ .

# Function composition

- We can chain multiple functions using function composition.
  - This is written either as  $g \circ f(x)$  or  $g(f(x))$ .
  - It is read as “ $g$  composed with  $f$ ” or “ $g$  of  $f$  of  $x$ ”

# Function composition

- We can chain multiple functions using function composition.
  - This is written either as  $g \circ f(x)$  or  $g(f(x))$ .
  - It is read as “ $g$  composed with  $f$ ” or “ $g$  of  $f$  of  $x$ ”
  - Generally,  $g \circ f(x) \neq f \circ g(x)$

# Function composition

- We can chain multiple functions using function composition.
  - This is written either as  $g \circ f(x)$  or  $g(f(x))$ .
  - It is read as “ $g$  composed with  $f$ ” or “ $g$  of  $f$  of  $x$ ”
  - Generally,  $g \circ f(x) \neq f \circ g(x)$
- Example: Suppose  $f(x) = 2x$  and  $g(x) = x^3$

# Function composition

- We can chain multiple functions using function composition.
  - This is written either as  $g \circ f(x)$  or  $g(f(x))$ .
  - It is read as “ $g$  composed with  $f$ ” or “ $g$  of  $f$  of  $x$ ”
  - Generally,  $g \circ f(x) \neq f \circ g(x)$
- Example: Suppose  $f(x) = 2x$  and  $g(x) = x^3$ 
  - $g \circ f(x) = (2x)^3 = 8x^3$

# Function composition

- We can chain multiple functions using function composition.
  - This is written either as  $g \circ f(x)$  or  $g(f(x))$ .
  - It is read as “ $g$  composed with  $f$ ” or “ $g$  of  $f$  of  $x$ ”
  - Generally,  $g \circ f(x) \neq f \circ g(x)$
- Example: Suppose  $f(x) = 2x$  and  $g(x) = x^3$ 
  - $g \circ f(x) = (2x)^3 = 8x^3$
  - $f \circ g(x) = 2(x^3) = 2x^3$

# Examples of functions of one variable - linear equation




Figure:  $y=2+x$



# Examples of functions of one variable - linear equation




Figure:  $y=2+x$

- This is the classic linear equation  
 $y = a + bx$  or  $y = mx + n$

# Examples of functions of one variable - linear equation




Figure:  $y=2+x$

- This is the classic linear equation  $y = a + bx$  or  $y = mx + n$
- $a$  and  $b$  are constants and  $x$  is the variable.

# Examples of functions of one variable - linear equation




Figure:  $y=2+x$

- This is the classic linear equation  $y = a + bx$  or  $y = mx + n$
- $a$  and  $b$  are constants and  $x$  is the variable.
- $a$  is the intercept and  $b$  is the slope of the line, or the amount that  $y$  changes given a one-unit increase in  $x$ .

# Examples of functions of one variable - Quadratic function




Figure:  $y = -2x^2 + 8x + 6$

# Examples of functions of one variable - Quadratic function




Figure:  $y = -2x^2 + 8x + 6$

- This is the classical quadratic function:

$$f(x) = ax^2 + bx + c \text{ or}$$

$$f(x) = \alpha + \beta_1 x + \beta_2 x^2$$

# Examples of functions of one variable - Quadratic function




Figure:  $y = -2x^2 + 8x + 6$

- This is the classical quadratic function:  
 $f(x) = ax^2 + bx + c$  or  
 $f(x) = \alpha + \beta_1x + \beta_2x^2$
- If we set  $a > 0$  ( $\beta_2 > 0$ ) we get a curve shaped like an U (a convex parabola).

# Examples of functions of one variable - Quadratic function





Figure:  $y = -2x^2 + 8x + 6$


- This is the classical quadratic function:  
 $f(x) = ax^2 + bx + c$  or  
 $f(x) = \alpha + \beta_1x + \beta_2x^2$
- If we set  $a > 0$  ( $\beta_2 > 0$ ) we get a curve shaped like an U (a convex parabola).
- If we set  $a < 0$  ( $\beta_2 < 0$ ) we get a curve shaped like an inverse U (a concave parabola).

# Exponent, logarithms and roots - Graphs


## Exponential function



### Logarithmic function



### Root function





# Functions and Theorizing

# Functions and Theorizing

- Functions are important for theorizing about politics

# Functions and Theorizing

- Functions are important for theorizing about politics
- Most quantitative social science boils down to how two variables relate!

# Functions and Theorizing

- Functions are important for theorizing about politics
- Most quantitative social science boils down to how two variables relate!
- Our examples suggest a precise function between  $x$  and  $y$





# Functions and Theorizing

- Functions are important for theorizing about politics
- Most quantitative social science boils down to how two variables relate!
- Our examples suggest a precise function between  $x$  and  $y$
- In statistics, you will *theorize* the functional form of the relationship

# Functions and Theorizing

- Functions are important for theorizing about politics
- Most quantitative social science boils down to how two variables relate!
- Our examples suggest a precise function between  $x$  and  $y$
- In statistics, you will *theorize* the functional form of the relationship
- Statistical methods will *estimate* the most appropriate parameters; “a line of best fit”

# Anscombe's Quartet



# Solving Equations



# Solving Equations

- 1 Isolate the variable you are looking for

# Solving Equations

- 1 Isolate the variable you are looking for
- 2 Combine like terms

# Solving Equations

- 1 Isolate the variable you are looking for
- 2 Combine like terms
- 3 Factor and cancel

# Solving Equations

- 1 Isolate the variable you are looking for
- 2 Combine like terms
- 3 Factor and cancel
- 4 Operate on both sides of the equation

# Solving Equations

- 1 Isolate the variable you are looking for
- 2 Combine like terms
- 3 Factor and cancel
- 4 Operate on both sides of the equation
- 5 Check your answer

# Special products

# Special products

①  $(a + b)^2 = a^2 + 2ab + b^2$

# Special products

$$\textcircled{1} \quad (a + b)^2 = a^2 + 2ab + b^2$$

$$\textcircled{2} \quad (a - b)^2 = a^2 - 2ab + b^2$$



# Special products

- ①  $(a + b)^2 = a^2 + 2ab + b^2$
- ②  $(a - b)^2 = a^2 - 2ab + b^2$
- ③  $(x + a)(x + b) = x^2 + (a + b)x + ab$

# Special products

$$\textcircled{1} (a+b)^2 = a^2 + 2ab + b^2$$

$$\textcircled{2} (a-b)^2 = a^2 - 2ab + b^2$$

$$\textcircled{3} (x+a)(x+b) = x^2 + (a+b)x + ab$$

$$\textcircled{4} (a+b)(a-b) = a^2 - b^2$$

# Special products

$$\textcircled{1} \quad (a + b)^2 = a^2 + 2ab + b^2$$

$$\textcircled{2} \quad (a - b)^2 = a^2 - 2ab + b^2$$

$$\textcircled{3} \quad (x + a)(x + b) = x^2 + (a + b)x + ab$$

$$\textcircled{4} \quad (a + b)(a - b) = a^2 - b^2$$

$$\textcircled{5} \quad (a + b + c)^2 = a^2 + b^2 + c^2 + 2(ab + ac + bc)$$

# Special products

$$\textcircled{1} (a+b)^2 = a^2 + 2ab + b^2$$

$$\textcircled{2} (a-b)^2 = a^2 - 2ab + b^2$$

$$\textcircled{3} (x+a)(x+b) = x^2 + (a+b)x + ab$$

$$\textcircled{4} (a+b)(a-b) = a^2 - b^2$$

$$\textcircled{5} (a+b+c)^2 = a^2 + b^2 + c^2 + 2(ab+ac+bc)$$

$$\textcircled{6} (a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

# Factoring

Factoring: rearranging terms in an equation to make it simpler

$$\frac{3x^4 + 3x^3 - 6x^2}{6x^2 + 12x} \quad (1)$$

# Factoring

Factoring: rearranging terms in an equation to make it simpler

$$\frac{3x^4 + 3x^3 - 6x^2}{6x^2 + 12x} \quad (1)$$

$$\frac{3x^2(x^2 + x - 2)}{6x(x + 2)} \quad (2)$$

# Factoring

Factoring: rearranging terms in an equation to make it simpler

$$\frac{3x^4 + 3x^3 - 6x^2}{6x^2 + 12x} \quad (1)$$

$$\frac{3x^2(x^2 + x - 2)}{6x(x + 2)} \quad (2)$$

$$\frac{3x^2(x + 2)(x - 1)}{6x(x + 2)} \quad (3)$$

# Factoring

Factoring: rearranging terms in an equation to make it simpler

$$\frac{3x^4 + 3x^3 - 6x^2}{6x^2 + 12x} \quad (1)$$

$$\frac{3x^2(x^2 + x - 2)}{6x(x + 2)} \quad (2)$$

$$\frac{3x^2(x + 2)(x - 1)}{6x(x + 2)} \quad (3)$$

$$\frac{3x^2(x - 1)}{6x} \quad (4)$$



# Factoring

Factoring: rearranging terms in an equation to make it simpler

$$\frac{3x^4 + 3x^3 - 6x^2}{6x^2 + 12x} \quad (1)$$

$$\frac{3x^2(x^2 + x - 2)}{6x(x + 2)} \quad (2)$$

$$\frac{3x^2(x + 2)(x - 1)}{6x(x + 2)} \quad (3)$$

$$\frac{3x^2(x - 1)}{6x} \quad (4)$$

$$\frac{x(x - 1)}{2} \quad (5)$$

# Quadratics and Special Products

Quadratic polynomials can be factored into the product of two terms:

$$(x \pm ?) \times (x \pm ?)$$

$$x^2 - 8x + 16 = 9 \quad (6)$$

# Quadratics and Special Products

Quadratic polynomials can be factored into the product of two terms:

$$(x \pm ?) \times (x \pm ?)$$

$$x^2 - 8x + 16 = 9 \tag{6}$$

$$(x \pm ?) \times (x \pm ?) = x^2 - 8x + 16$$

# Quadratics and Special Products

Quadratic polynomials can be factored into the product of two terms:

$$(x \pm ?) \times (x \pm ?)$$

$$x^2 - 8x + 16 = 9 \quad (6)$$

$$(x \pm ?) \times (x \pm ?) = x^2 - 8x + 16$$

$$(x - 4)^2 = 9 \quad (7)$$

# Quadratics and Special Products

Quadratic polynomials can be factored into the product of two terms:

$$(x \pm ?) \times (x \pm ?)$$

$$x^2 - 8x + 16 = 9 \quad (6)$$

$$(x \pm ?) \times (x \pm ?) = x^2 - 8x + 16$$

$$(x - 4)^2 = 9 \quad (7)$$

$$(x - 4) = \pm 3 \quad (8)$$

# Quadratics and Special Products

Quadratic polynomials can be factored into the product of two terms:

$$(x \pm ?) \times (x \pm ?)$$

$$x^2 - 8x + 16 = 9 \quad (6)$$

$$(x \pm ?) \times (x \pm ?) = x^2 - 8x + 16$$

$$(x - 4)^2 = 9 \quad (7)$$

$$(x - 4) = \pm 3 \quad (8)$$

$$x = 4 \pm 3 \quad (9)$$

# Quadratics and Special Products

Quadratic polynomials can be factored into the product of two terms:

$$(x \pm ?) \times (x \pm ?)$$

$$x^2 - 8x + 16 = 9 \quad (6)$$

$$(x \pm ?) \times (x \pm ?) = x^2 - 8x + 16$$

$$(x - 4)^2 = 9 \quad (7)$$

$$(x - 4) = \pm 3 \quad (8)$$


$$x = 4 \pm 3 \quad (9)$$

$$x = 1 \text{ OR } x = 7$$

# Why two solutions?



# Why two solutions?



## Another example

$$x^2 + 7x = -12 \quad (10)$$

## Another example

$$x^2 + 7x = -12 \quad (10)$$

$$x^2 + 7x + 12 = 0 \quad (11)$$

## Another example

$$x^2 + 7x = -12 \quad (10)$$

$$x^2 + 7x + 12 = 0 \quad (11)$$

How can we use special products here? Find  $a$  and  $b$

$$(x + a)(x + b) = x^2 + (a + b)x + ab \quad (12)$$

## Another example

$$x^2 + 7x = -12 \quad (10)$$

$$x^2 + 7x + 12 = 0 \quad (11)$$

How can we use special products here? Find  $a$  and  $b$

$$(x + a)(x + b) = x^2 + (a + b)x + ab \quad (12)$$

$$(x + 3)(x + 4) = 0 \quad (13)$$

## Another example

$$x^2 + 7x = -12 \quad (10)$$

$$x^2 + 7x + 12 = 0 \quad (11)$$

How can we use special products here? Find  $a$  and  $b$

$$(x + a)(x + b) = x^2 + (a + b)x + ab \quad (12)$$

$$(x + 3)(x + 4) = 0 \quad (13)$$

Either  $x + 3 = 0$  OR  $x + 4 = 0$   
 $x = -3$  OR  $x = -4$

# Quadratic Formula

Useful if special products can't easily be found!

$$3x^2 + 8x - 13 = 0 \quad (14)$$

# Quadratic Formula

Useful if special products can't easily be found!

$$3x^2 + 8x - 13 = 0 \quad (14)$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad (15)$$



# Quadratic Formula

$$3x^2 + 8x - 13 = 0 \quad (16)$$

# Quadratic Formula

$$3x^2 + 8x - 13 = 0 \quad (16)$$

$$x = \frac{-8 \pm \sqrt{8^2 - (4 * 3 * 13)}}{2 * 3} \quad (17)$$

# Quadratic Formula

$$3x^2 + 8x - 13 = 0 \quad (16)$$

$$x = \frac{-8 \pm \sqrt{8^2 - (4 * 3 * 13)}}{2 * 3} \quad (17)$$


$$x = \frac{-8 + \sqrt{220}}{6} \quad (18)$$

$$x \approx 1.14 \quad (19)$$

$$x = \frac{-8 - \sqrt{220}}{6} \quad (20)$$

$$x \approx -3.81 \quad (21)$$

# Checking our work graphically



# Sequences and series

# Sequences and series

- A *sequence* is an ordered list of numbers

# Sequences and series

- A *sequence* is an ordered list of numbers
  - A sequence can be infinite, such as  $1, 2, 3, 4, \dots$

# Sequences and series

- A *sequence* is an ordered list of numbers
  - A sequence can be infinite, such as  $1, 2, 3, 4, \dots$
  - Or a sequence can be finite, such as  $5, 10, 15, 20, 25$



# Sequences and series

- A *sequence* is an ordered list of numbers
  - A sequence can be infinite, such as  $1, 2, 3, 4, \dots$
  - Or a sequence can be finite, such as  $5, 10, 15, 20, 25$
- A *series* is the sum of a sequence.

# Sequences and series

- A *sequence* is an ordered list of numbers
  - A sequence can be infinite, such as  $1, 2, 3, 4, \dots$
  - Or a sequence can be finite, such as  $5, 10, 15, 20, 25$
- A *series* is the sum of a sequence.
  - Typically noted as  $\sum_{i=1}^N x_i$  which means add the terms in the sequence beginning at  $x_1$  and stopping at  $x_n$ .

# Sequences and series

- A *sequence* is an ordered list of numbers
  - A sequence can be infinite, such as  $1, 2, 3, 4, \dots$
  - Or a sequence can be finite, such as  $5, 10, 15, 20, 25$
- A *series* is the sum of a sequence.
  - Typically noted as  $\sum_{i=1}^N x_i$  which means add the terms in the sequence beginning at  $x_1$  and stopping at  $x_n$ .
  - For an infinite sequence,  $N = \infty$

# Limits

# Limits

- *Limits help us describe the behavior of a sequence, series, or function as it approaches a given value.*

# Limits

- *Limits help us describe the behavior of a sequence, series, or function as it approaches a given value.*
  - A sequence/series/function *converges* if it has a finite limit.

# Limits

- *Limits help us describe the behavior of a sequence, series, or function as it approaches a given value.*
  - A sequence/series/function *converges* if it has a finite limit.
  - A sequence/series/function *diverges* if it has no limit or the limit is  $\pm\infty$

# Limits

- *Limits help us describe the behavior of a sequence, series, or function as it approaches a given value.*
  - A sequence/series/function *converges* if it has a finite limit.
  - A sequence/series/function *diverges* if it has no limit or the limit is  $\pm\infty$
- The limit of a sequence is the number  $L$  such that as we approach infinity,  $x_i$  gets arbitrarily close to  $L$ . Noted as:  $\lim_{i \rightarrow \infty} x_i = L$



# Limits

- *Limits help us describe the behavior of a sequence, series, or function as it approaches a given value.*
  - A sequence/series/function *converges* if it has a finite limit.
  - A sequence/series/function *diverges* if it has no limit or the limit is  $\pm\infty$
- The limit of a sequence is the number  $L$  such that as we approach infinity,  $x_i$  gets arbitrarily close to  $L$ . Noted as:  $\lim_{i \rightarrow \infty} x_i = L$
- Example of the limit of a sequence:

# Limits

- *Limits help us describe the behavior of a sequence, series, or function as it approaches a given value.*
  - A sequence/series/function *converges* if it has a finite limit.
  - A sequence/series/function *diverges* if it has no limit or the limit is  $\pm\infty$
- The limit of a sequence is the number  $L$  such that as we approach infinity,  $x_i$  gets arbitrarily close to  $L$ . Noted as:  $\lim_{i \rightarrow \infty} x_i = L$
- Example of the limit of a sequence:
  - The limit of the sequence  $\{i\}_{i=1}^{\infty}$  does not have an “endpoint” and approaches infinity, so it diverges.

# Limits

- *Limits help us describe the behavior of a sequence, series, or function as it approaches a given value.*
  - A sequence/series/function *converges* if it has a finite limit.
  - A sequence/series/function *diverges* if it has no limit or the limit is  $\pm\infty$
- The limit of a sequence is the number  $L$  such that as we approach infinity,  $x_i$  gets arbitrarily close to  $L$ . Noted as:  $\lim_{i \rightarrow \infty} x_i = L$
- Example of the limit of a sequence:
  - The limit of the sequence  $\{i\}_{i=1}^{\infty}$  does not have an “endpoint” and approaches infinity, so it diverges.
  - The limit of the sequence  $\{\frac{3}{10^i}\}_{i=1}^{\infty}$  approaches zero as  $i \rightarrow \infty$ , so it converges.

# Limits

# Limits

- The limit of a series is similar, but you are not looking for an “endpoint”. In this case, you are looking for the sum of all elements in an infinite sequence. For example:

# Limits

- The limit of a series is similar, but you are not looking for an “endpoint”. In this case, you are looking for the sum of all elements in an infinite sequence. For example:
  - $\lim_{n \rightarrow \infty} \sum_{i=1}^n i = \infty$  So, this series is divergent.

# Limits

- The limit of a series is similar, but you are not looking for an “endpoint”. In this case, you are looking for the sum of all elements in an infinite sequence. For example:
  - $\lim_{n \rightarrow \infty} \sum_{i=1}^n i = \infty$  So, this series is divergent.
  - $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{2^i}$ . This series converges. Where?

# Limits

- The limit of a series is similar, but you are not looking for an “endpoint”. In this case, you are looking for the sum of all elements in an infinite sequence. For example:
  - $\lim_{n \rightarrow \infty} \sum_{i=1}^n i = \infty$  So, this series is divergent.
  - $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{2^i}$ . This series converges. Where?
- For a function  $y = f(x)$ , the limit is the value of  $y$  that the function tend towards as small steps are taken towards a value  $x = c$



# Limits

- The limit of a series is similar, but you are not looking for an “endpoint”. In this case, you are looking for the sum of all elements in an infinite sequence. For example:
  - $\lim_{n \rightarrow \infty} \sum_{i=1}^n i = \infty$  So, this series is divergent.
  - $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{2^i}$ . This series converges. Where?
- For a function  $y = f(x)$ , the limit is the value of  $y$  that the function tend towards as small steps are taken towards a value  $x = c$
- Piecewise function: the function (relationship between  $x$  and  $y$ ) is different for different parts of the domain

## Piecewise Function Example

Example: Estimate the value of the following limits:  $\lim_{x \rightarrow 0^-} f(x)$  and  $\lim_{x \rightarrow 0^+} f(x)$  for the following function:



$$f(x) = \begin{cases} x+1 & x \leq 0 \\ x^2 & x > 0 \end{cases}$$

## Piecewise Function Example

Example: Estimate the value of the following limits:  $\lim_{x \rightarrow 0^-} f(x)$  and  $\lim_{x \rightarrow 0^+} f(x)$  for the following function:



$$f(x) = \begin{cases} x+1 & x \leq 0 \\ x^2 & x > 0 \end{cases}$$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} x + 1 = 1$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} x^2 = 0$$

An infinite number of mathematicians walk into a bar. The first one tells the bartender he wants a beer. The second one says he wants half a beer. The third one says she wants a fourth of a beer.

The bartender interrupts, puts two beers on the bar and says, "You people need to learn your limits."



# Limit Example

# Limit Example

- The function  $f(x) = \frac{1}{x-5}$  is undefined at which value of  $x$ ?

# Limit Example

- The function  $f(x) = \frac{1}{x-5}$  is undefined at which value of  $x$ ?
- The function  $f(x)$  has a limit at  $x = 5$

# Limit Example

- The function  $f(x) = \frac{1}{x-5}$  is undefined at which value of  $x$ ?
- The function  $f(x)$  has a limit at  $x = 5$
- What is the value of  $\lim_{x \rightarrow 5} f(x)$



# Limit Example

- The function  $f(x) = \frac{1}{x-5}$  is undefined at which value of  $x$ ?
- The function  $f(x)$  has a limit at  $x = 5$
- What is the value of  $\lim_{x \rightarrow 5} f(x)$
- Does that value change whether we approach the limit from negative or positive?

# Limit Example

- The function  $f(x) = \frac{1}{x-5}$  is undefined at which value of  $x$ ?
- The function  $f(x)$  has a limit at  $x = 5$
- What is the value of  $\lim_{x \rightarrow 5} f(x)$
- Does that value change whether we approach the limit from negative or positive?
- Functions are **continuous** if they do not have sudden breaks, and **discontinuous** otherwise

# Math Camp Exercises - Day 1

- ① M&S Chapter 1: Exercises 3, 5
- ② M&S Chapter 2: Exercises 16, 17, 19, 20, 21, 22, 26, 29
- ③ M&S Chapter 3: Exercises 2, 3
- ④ M&S Chapter 4: Exercises 3, 4, 5, 6, 9