

7-0 (A)

Recd 5/21/00 upl

SPECIES: GLAUCOUS WINGED GULL
(Vernacular Name)LARUS GLAUCESCA
(Scientific Name)

2000 - 61

DATE(S): 3/9/00 ; 4/2/00 TIME OF OBSERVATION: 3/9: 7:20 PM TO 5:30 PM
4/2: 12:00 TO 12:45EXACT LOCATION: 3/9: HORSESHOE LAKE, LOVELAND ; 4/2: LARIMER COUNTY LANDFILLNEAREST TOWN: 3/9: LOVELAND ; 4/2: FT. COLLINS COUNTY: LARIMER

OTHER OBSERVERS WHO INDEPENDENTLY IDENTIFIED THIS BIRD:

BILL LISOWSKYOPTICAL EQUIPMENT: BAUSH & LOMB ELITE BINOCULARS, BUSHNELL SPACEMASTER SCOPEWEATHER CONDITIONS: 3/9: BRIGHT WITH FEW CLOUDS ; 4/2: CLOUDY WITH SNOWBIRD/SUN ORIENTATION: BIRD IN FRONT, SUN BEHIND ON 3/9DISTANCE FROM BIRD AND HOW MEASURED: 3/9: 50-80 YDS ; 4/2: 30-40 YDSNUMBER OF BIRDS SEEN: 1 SEX: _____ PLUMAGE: 1ST BASICBEHAVIOR AND DEMEANOR: 3/9: RESTING ON SANDY SHORE AT HORSESHOE RES. WITH VARIOUS OTHER GULLS. THE BIRD WAS LIGHTER THAN THE THAYERS GULLS IN VICINITY AND WAS AS LARGE AS THE HERRING GULLS THERE ALSO. WHEN THE BIRD WOULD STRETCH ITS WING OR FLY, THE ENTIRE WING WAS A UNIFORM LIGHT TANNISH GRAY, SAME AS BACK COLOR ; 4/2 FORAGING AT LARIMER CTY LANDFILLHEAD & NECK (forehead, crown, auricular & malar regions, median line, etc.): VERY LIGHT TANNISH GRAY, DARKER THAN GLAUCOUS GULL, BUT MUCH PALER THAN THAYERS GULLS.EYES & LORES (supercilium, eye-ring, eye-stripe, color, etc.): EYE DARKLOWER & UPPER MANDIBLES (color & shape): ALL LARGISH, AND ALL BLACK ; NO PALE AREA AT BASE THAT WE COULD SEE.LEGS & FEET: BEIGISH GRAYUPPERPARTS (nape, back, rump, uppertail coverts, etc.): LIGHT TANNISH GRAY, DARKER THAN GLAUCOUS GULL BUT MUCH LIGHTER THAN THAYERS GULLS THAT WERE AT HORSESHOE LAKE. THE SHADE WAS LIGHTER THAN DEPICTED IN NAT. GEO. FOR GLAUCOUS WINGED, BUT SLIGHTLY DARKER THAN DEPICTED IN NAT. GEO FOR GLAUCOUSUNDERPARTS (chin, throat, belly, sides, flanks, crissum, etc.): LIGHT TANNISH GRAY SAME AS UPPERPARTS. THE SHADE WAS LIGHTER THAN DEPICTED IN NAT. GEO. FOR GLAUCOUS WINGED, BUT SLIGHTLY DARKER THAN DEPICTED IN NAT. GEO. FOR GLAUCOUS.

RETRICES (inner, middle & outer tail feathers; color, pattern, shape, etc.):
SAME COLOR AS UPPERPARTS, SOME LIGHT BARRING ON
UNDERTAIL COVERTS

WINGS (color, length, shape, bars, coverts, etc.): SAME COLOR AS UPPERPARTS.
THERE WAS NO DARKENING IN THE PRIMARIES, COLOR WAS
UNIFORM TO WING TIAS, UNDERWING PRIMARIES LOOKED SLIGHTLY
LIGHTER TAN COLOR THAN UPPERWING

VOCALIZATION (call/song, method of delivery, i.e., from perch, in flight):
NONE

HABITAT: LAKESHORE AND LANDFILL

ELIMINATION OF SIMILARLY APPEARING SPECIES:

GLAUCOUS GULL SLIGHTLY LARGER AND MUCH LIGHTER WITH
WHITISH PRIMARIES. BILL TWO-TONED ON 1ST BASIC

THAYERS GULL SMALLER WITH CONSIDERABLY DARKER PRIMARIES
THAN INNER WING, DARKER TAIL

NEJONS GULL (HYBRID) MUCH DARKER ON PRIMARIES AND
TAIL

PRIOR EXPERIENCE WITH THIS AND SIMILARLY APPEARING SPECIES:

I HAVE SEEN DOZENS OF THAYERS GULLS AND HAVE SEEN
15-25 GLAUCOUS GULLS.

THIS WOULD BE MY SECOND GLAUCOUS-WINGED GULL IN
COLORADO, BUT I HAVE SEEN SEVERAL OTHER SPECIMENS
ON THE WEST COAST.

BOOKS, ILLUSTRATIONS AND ADVICE CONSULTED THAT INFLUENCED THIS REPORT:

NATIONAL GEOGRAPHIC

THIS REPORT WRITTEN FROM NOTES MADE DURING ☒ AFTER ☒ OBSERVATION; ☐ FROM MEMORY

NAME: JOE MAMMOSER SIGNATURE: *Joe Mammoser* DATE: 4/15/00