

0.1 Todo: [The Kuznetsov Trace Formula]

The Kuznetsov trace formula is an analog of the Petersson trace formula for weight zero Maass forms. From ??, $\mathcal{L}(N, \chi)$ admits an orthonormal basis of Maass forms for the point spectrum (these forms are generally not Hecke-Maass eigenforms because they need not be Hecke normalized or even cuspidal in the case of the discrete spectrum). However, by ?? and ?? we make take this orthonormal basis to consist of Hecke-Maass eigenforms and the constant function. Denote this basis by $\{u_j\}_{j \geq 0}$ with $u_0(z) = 1$ and let u_j be of type ν_j for $j \geq 1$. In particular, $\{u_j\}_{j \geq 1}$ is an orthonormal basis of Hecke-Maass eigenforms and each such form admits a Fourier series at the \mathfrak{a} cusp given by

$$(u_j | \sigma_{\mathfrak{a}})(z) = \sum_{n \neq 0} a_{j,\mathfrak{a}}(n) \sqrt{y} K_{\nu_j}(2\pi n y) e^{2\pi i n x}.$$

The Kuznetsov trace formula is an equation relating the Fourier coefficients $a_{j,\mathfrak{a}}(n)$ and $a_{j,\mathfrak{b}}(n)$ of the basis $\{u_j\}_{j \geq 1}$ for two cusps \mathfrak{a} and \mathfrak{b} of $\Gamma_0(N) \backslash \mathbb{H}$ to a sum of integral transforms involving test functions and Salié sums. Similar to the Petersson trace formula, we will compute the inner product of two Poincaré series $P_{n,\chi,\mathfrak{a}}(z, \psi)(z)$ and $P_{m,\chi,\mathfrak{b}}(z, \varphi)(z)$ in two different ways. The first will be geometric in nature while the second will be spectral. We first need to compute the Fourier series of such a Poincaré series. Although we will not need it explicitly, we will work over any congruence subgroup:

Proposition 0.1.1. *Let $m \geq 1$, χ be Dirichlet character with conductor dividing the level, \mathfrak{a} and \mathfrak{b} be cusps of $\Gamma \backslash \mathbb{H}$, and $\psi(y)$ be a smooth function such that $\psi(y) \ll_{\varepsilon} y^{1+\varepsilon}$ as $y \rightarrow 0$. The Fourier series of $P_{m,\chi,\mathfrak{a}}(z, \psi)$ on $\Gamma \backslash \mathbb{H}$ at the \mathfrak{b} cusp is given by*

$$(P_{m,\chi,\mathfrak{a}} | \sigma_{\mathfrak{b}})(z, \psi) = \sum_{t \in \mathbb{Z}} \left(\delta_{\mathfrak{a},\mathfrak{b}} \delta_{m,t} \psi(\text{Im}(z)) + \sum_{c \in \mathcal{C}_{\mathfrak{a},\mathfrak{b}}} \psi(y, m, t, c) S_{\chi,\mathfrak{a},\mathfrak{b}}(m, t, c) \right) e^{2\pi i t z},$$

where $\psi(y, m, t, c)$ is the integral transform given by

$$\psi(y, m, t, c) = \int_{\text{Im}(z)=y} \psi \left(\frac{y}{|cz|^2} \right) e^{-\frac{2\pi i m}{c^2 z} - 2\pi i t z} dz.$$

Proof. From the cocycle condition and ??, we have

$$(P_{m,\chi,\mathfrak{a}} | \sigma_{\mathfrak{b}})(z, \psi) = \delta_{\mathfrak{a},\mathfrak{b}} \psi(\text{Im}(z)) e^{2\pi i m z} + \sum_{\substack{c \in \mathcal{C}_{\mathfrak{a},\mathfrak{b}}, d \in \mathbb{Z} \\ d \pmod{c} \in \mathcal{D}_{\mathfrak{a},\mathfrak{b}}(c)}} \bar{\chi}(d) \psi \left(\frac{\text{Im}(z)}{|cz + d|^2} \right) e^{2\pi i m \left(\frac{a}{c} - \frac{1}{c^2 z + cd} \right)},$$

where a and b are chosen such that $\det \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) = 1$ and we have used the fact that

$$\frac{a}{c} - \frac{1}{c^2 z + cd} = \frac{az + b}{cz + d}.$$

Summing over all pairs (c, d) with $c \in \mathcal{C}_{\mathfrak{a},\mathfrak{b}}$, $d \in \mathbb{Z}$, and $d \pmod{c} \in \mathcal{D}_{\mathfrak{a},\mathfrak{b}}(c)$ is the same as summing over all triples (c, ℓ, r) with $c \in \mathcal{C}_{\mathfrak{a},\mathfrak{b}}$, $\ell \in \mathbb{Z}$, and r taken modulo c with $r \in \mathcal{D}_{\mathfrak{a},\mathfrak{b}}(c)$. Indeed, this is seen by writing $d = c\ell + r$. Moreover, since $ad - bc = 1$ we have $a(c\ell + r) - bc = 1$ which further implies that

$ar \equiv 1 \pmod{c}$. So we may take a to be the inverse for r modulo c . Then

$$\begin{aligned}
\sum_{\substack{c \in \mathcal{C}_{a,b}, d \in \mathbb{Z} \\ d \pmod{c} \in \mathcal{D}_{a,b}(c)}} \bar{\chi}(d) \psi \left(\frac{\text{Im}(z)}{|cz + d|^2} \right) e^{2\pi i m \left(\frac{a}{c} - \frac{1}{c^2 z + cd} \right)} &= \sum_{(c, \ell, r)} \bar{\chi}(c\ell + r) \psi \left(\frac{\text{Im}(z)}{|cz + c\ell + r|^2} \right) e^{2\pi i m \left(\frac{a}{c} - \frac{1}{c^2 z + c^2 \ell + cr} \right)} \\
&= \sum_{(c, \ell, r)} \bar{\chi}(r) \psi \left(\frac{\text{Im}(z)}{|cz + c\ell + r|^2} \right) e^{2\pi i m \left(\frac{a}{c} - \frac{1}{c^2 z + c^2 \ell + cr} \right)} \\
&= \sum_{\substack{c \in \mathcal{C}_{a,b} \\ r \in \mathcal{D}_{a,b}(c)}} \sum_{\ell \in \mathbb{Z}} \bar{\chi}(r) \psi \left(\frac{\text{Im}(z)}{|cz + c\ell + r|^2} \right) e^{2\pi i m \left(\frac{a}{c} - \frac{1}{c^2 z + c^2 \ell + cr} \right)} \\
&= \sum_{\substack{c \in \mathcal{C}_{a,b} \\ r \in \mathcal{D}_{a,b}(c)}} \bar{\chi}(r) \sum_{\ell \in \mathbb{Z}} \psi \left(\frac{\text{Im}(z)}{|cz + c\ell + r|^2} \right) e^{2\pi i m \left(\frac{a}{c} - \frac{1}{c^2 z + c^2 \ell + cr} \right)},
\end{aligned}$$

where on the right-hand side it is understood that we are summing over all triples (c, ℓ, r) with the prescribed properties and the second line holds since χ has conductor dividing the level and $d \in \mathcal{D}_{a,b}(c)$ is determined modulo c . Now let

$$I_{c,r}(z, \psi) = \sum_{\ell \in \mathbb{Z}} \psi \left(\frac{\text{Im}(z)}{|cz + c\ell + r|^2} \right) e^{2\pi i m \left(\frac{a}{c} - \frac{1}{c^2 z + c^2 \ell + cr} \right)}.$$

We apply the Poisson summation formula to $I_{c,r}(z, \psi)$. This is allowed since the summands are absolutely integrable by ??, as they exhibit polynomial decay of order $\sigma > 1$ because $\psi(y) \ll_{\varepsilon} y^{1+\varepsilon}$ as $y \rightarrow 0$, and $I_{c,r}(z, \psi)$ is holomorphic because $(P_{m,\chi,a}|\sigma_b)(z, \psi)$ is. By the identity theorem it suffices to apply the Poisson summation formula for $z = iy$ with $y > 0$. So let $f(x)$ be given by

$$f(x) = \psi \left(\frac{y}{|cx + r + icy|^2} \right) e^{2\pi i m \left(\frac{a}{c} - \frac{1}{c^2 x + cr + ic^2 y} \right)}.$$

As we have just noted, $f(x)$ is absolutely integrable on \mathbb{R} . We compute the Fourier transform:

$$\hat{f}(t) = \int_{-\infty}^{\infty} f(x) e^{-2\pi i t x} dx = \int_{-\infty}^{\infty} \psi \left(\frac{y}{|cx + r + icy|^2} \right) e^{2\pi i m \left(\frac{a}{c} - \frac{1}{c^2 x + cr + ic^2 y} \right)} e^{-2\pi i t x} dx.$$

Complexify the integral to get

$$\int_{\text{Im}(z)=0} \psi \left(\frac{y}{|cz + r + icy|^2} \right) e^{2\pi i m \left(\frac{a}{c} - \frac{1}{c^2 z + cr + ic^2 y} \right)} e^{-2\pi i t z} dz.$$

Now make the change of variables $z \rightarrow z - \frac{r}{c} - iy$ to obtain

$$e^{2\pi i m \frac{a}{c} + 2\pi i t \frac{r}{c} - 2\pi t y} \int_{\text{Im}(z)=y} \psi \left(\frac{y}{|cz|^2} \right) e^{-\frac{2\pi i m}{c^2 z} - 2\pi i t z} dz.$$

As the remaining integral is $\psi(y, m, t, c)$, it follows that

$$\hat{f}(t) = \psi(y, m, t, c) e^{2\pi i m \frac{a}{c} + 2\pi i t \frac{r}{c} - 2\pi t y}.$$

By the Poisson summation formula and the identity theorem, we have

$$I_{c,r}(z, \psi) = \sum_{t \in \mathbb{Z}} (\psi(y, m, t, c) e^{2\pi i m \frac{a}{c} + 2\pi i t \frac{r}{c}}) e^{2\pi i t z},$$

for all $z \in \mathbb{H}$. Substituting this back into the Eisenstein series gives a form of the Fourier series:

$$\begin{aligned}
(P_{m,\chi,a}|\sigma_b)(z, \psi) &= \delta_{a,b} \psi(\text{Im}(z)) e^{2\pi i m z} + \sum_{\substack{c \in \mathcal{C}_{a,b} \\ r \in \mathcal{D}_{a,b}}} \bar{\chi}(r) \sum_{t \in \mathbb{Z}} \psi(y, m, t, c) e^{2\pi i m \frac{a}{c} + 2\pi i t \frac{r}{c}} e^{2\pi i t z} \\
&= \sum_{t \in \mathbb{Z}} \left(\delta_{a,b} \delta_{m,t} \psi(\text{Im}(z)) + \sum_{\substack{c \in \mathcal{C}_{a,b} \\ r \in \mathcal{D}_{a,b}}} \bar{\chi}(r) \psi(y, m, t, c) e^{2\pi i m \frac{a}{c} + 2\pi i t \frac{r}{c}} \right) e^{2\pi i t z} \\
&= \sum_{t \in \mathbb{Z}} \left(\delta_{a,b} \delta_{m,t} \psi(\text{Im}(z)) + \sum_{c \in \mathcal{C}_{a,b}} \psi(y, m, t, c) \sum_{r \in \mathcal{D}_{a,b}} \bar{\chi}(r) e^{2\pi i m \frac{a}{c} + 2\pi i t \frac{r}{c}} \right) e^{2\pi i t z}.
\end{aligned}$$

We will simplify the innermost sum. Since a is the inverse for r modulo c , the innermost sum above becomes

$$\sum_{r \in \mathcal{D}_{a,b}} \bar{\chi}(r) e^{2\pi i m \frac{a}{c} + 2\pi i t \frac{r}{c}} = \sum_{r \in \mathcal{D}_{a,b}} \bar{\chi}(\bar{a}) e^{2\pi i m \frac{a}{c} + 2\pi i t \frac{\bar{a}}{c}} = \sum_{r \in \mathcal{D}_{a,b}} \chi(a) e^{\frac{2\pi i (am + \bar{a}t)}{c}} = S_{\chi,a,b}(m, t, c).$$

So at last, we obtain our desired Fourier series:

$$(P_{m,\chi,a}|\sigma_b)(z) = \sum_{t \in \mathbb{Z}} \left(\delta_{a,b} \delta_{m,t} \psi(\text{Im}(z)) + \sum_{c \in \mathcal{C}_{a,b}} \psi(y, m, t, c) S_{\chi,a,b}(m, t, c) \right) e^{2\pi i t z}.$$

□

We can now derive the first half of the Kuznetsov trace formula by computing the inner product between $P_{n,\chi,a}(z, \psi)$ and $P_{m,\chi,b}(z, \varphi)$:

$$\begin{aligned}
\langle P_{n,\chi,a}(\cdot, \psi), P_{m,\chi,b}(\cdot, \varphi) \rangle &= \frac{1}{V_{\Gamma_0(N)}} \int_{\mathcal{F}_{\Gamma_0(N)}} P_{n,\chi,a}(z, \psi) \overline{P_{m,\chi,b}(z, \varphi)} d\mu \\
&= \frac{1}{V_{\Gamma_0(N)}} \int_{\mathcal{F}_{\Gamma_0(N)}} \sum_{\gamma \in \Gamma_b \backslash \Gamma_0(N)} \chi(\gamma) P_{n,\chi,a}(z, \psi) \overline{\varphi(\text{Im}(\sigma_b^{-1} \gamma z))} e^{-2\pi i m \overline{\sigma_b^{-1} \gamma z}} d\mu \\
&= \frac{1}{V_{\Gamma_0(N)}} \int_{\mathcal{F}_{\Gamma_0(N)}} \sum_{\gamma \in \Gamma_b \backslash \Gamma_0(N)} P_{n,\chi,a}(\gamma z, \psi) \overline{\varphi(\text{Im}(\sigma_b^{-1} \gamma z))} e^{-2\pi i m \overline{\sigma_b^{-1} \gamma z}} d\mu \\
&= \frac{1}{V_{\Gamma_0(N)}} \int_{\mathcal{F}_{\sigma_b^{-1} \Gamma_0(N) \sigma_b}} \sum_{\gamma \in \Gamma_b \backslash \Gamma_0(N)} P_{n,\chi,a}(\gamma \sigma_b z, \psi) \overline{\varphi(\text{Im}(\sigma_b^{-1} \gamma \sigma_b z))} e^{-2\pi i m \overline{\sigma_b^{-1} \gamma \sigma_b z}} d\mu \\
&= \frac{1}{V_{\Gamma_0(N)}} \int_{\mathcal{F}_{\sigma_b^{-1} \Gamma_0(N) \sigma_b}} \sum_{\gamma \in \Gamma_\infty \backslash \sigma_b^{-1} \Gamma_0(N) \sigma_b^{-1}} P_{n,\chi,a}(\sigma_b \gamma z, \psi) \overline{\varphi(\text{Im}(\gamma z))} e^{-2\pi i m \overline{\gamma z}} d\mu \\
&= \frac{1}{V_{\Gamma_0(N)}} \int_{\Gamma_\infty \backslash \mathbb{H}} (P_{n,\chi,a}|\sigma_b)(z, \psi) \overline{\varphi(\text{Im}(z))} e^{-2\pi i m \bar{z}} d\mu,
\end{aligned}$$

where in the third line we have used the automorphy of $P_{n,\chi,a}(z, \psi)$, in the forth and fifth lines we have made the change of variables $z \rightarrow \sigma_b z$ and $\gamma \rightarrow \sigma_b \gamma \sigma_b^{-1}$ respectively, and in the sixth line we have unfolded. Now substitute in the Fourier series of $P_{n,\chi,a}(z, \psi)$ at the b cusp to obtain

$$\frac{1}{V_{\Gamma_0(N)}} \int_{\Gamma_\infty \backslash \mathbb{H}} \sum_{t \in \mathbb{Z}} \left(\delta_{a,b} \delta_{n,t} \psi(\text{Im}(z)) + \sum_{c \in \mathcal{C}_{a,b}} \psi(y, n, t, c) S_{\chi,a,b}(n, t, c) \right) \overline{\varphi(\text{Im}(z))} e^{2\pi i t z - 2\pi i m \bar{z}} d\mu,$$

which is equivalent to

$$\frac{1}{V_{\Gamma_0(N)}} \int_0^\infty \int_0^1 \sum_{t \geq 1} \left(\delta_{a,b} \delta_{n,t} \psi(y) + \sum_{c \in \mathcal{C}_{a,b}} \psi(y, n, t, c) S_{\chi,a,b}(n, t, c) \right) \overline{\varphi(y)} e^{2\pi i(t-m)x} e^{-2\pi(t+m)y} \frac{dx dy}{y^2}.$$

By the dominated convergence theorem, we can interchange the sum and the two integrals. Then ?? implies that the inner integral cuts off all of the terms except the diagonal $t = m$. This leaves

$$\frac{1}{V_{\Gamma_0(N)}} \int_0^\infty \left(\delta_{a,b} \delta_{n,m} \psi(y) + \sum_{c \in \mathcal{C}_{a,b}} \psi(y, n, m, c) S_{\chi,a,b}(n, m, c) \right) \overline{\varphi(y)} e^{-4\pi m y} \frac{dy}{y^2}.$$

Interchanging the integral and the remaining sum by the dominated convergence theorem again, we arrive at

$$\langle P_{n,\chi,a}(\cdot, \psi), P_{m,\chi,b}(\cdot, \varphi) \rangle = \delta_{a,b} \delta_{n,m} (\psi, \varphi)_{n,m} + \sum_{c \in \mathcal{C}_{a,b}} S_{\chi,a,b}(n, m, c) V(n, m, c, \psi, \varphi),$$

where we have set

$$(\psi, \varphi)_{n,m} = \frac{1}{V_{\Gamma_0(N)}} \int_0^\infty \psi(y) \overline{\varphi(y)} e^{-2\pi(n+m)y} \frac{dy}{y^2},$$

and

$$V(n, m, c; \psi, \varphi) = \frac{1}{V_{\Gamma_0(N)}} \int_0^\infty \int_{\text{Im}(z)=y} \psi\left(\frac{y}{|cz|^2}\right) \overline{\varphi(y)} e^{-\frac{2\pi i m}{c^2 z} - 2\pi i n z - 4\pi m y} \frac{dz dy}{y^2}.$$

This is the first half of the Kuznetsov trace formula. For the second half, ?? gives

$$P_{n,\chi,a}(\cdot, \psi) = \sum_{j \geq 0} \langle P_{n,\chi,a}(\cdot, \psi), u_j \rangle u_j(z) + \sum_a \frac{1}{4\pi} \int_{-\infty}^\infty \left\langle P_{n,\chi,a}(\cdot, \psi), E_a\left(\cdot, \frac{1}{2} + ir\right) \right\rangle E_a\left(z, \frac{1}{2} + ir\right) dr,$$

and

$$P_{m,\chi,a}(\cdot, \varphi) = \sum_{j \geq 0} \langle P_{m,\chi,a}(\cdot, \varphi), u_j \rangle u_j(z) + \sum_a \frac{1}{4\pi} \int_{-\infty}^\infty \left\langle P_{m,\chi,a}(\cdot, \varphi), E_a\left(\cdot, \frac{1}{2} + ir\right) \right\rangle E_a\left(z, \frac{1}{2} + ir\right) dr.$$

By orthonormality, it follows that

$$\begin{aligned} \langle P_{n,\chi,a}(\cdot, \psi), P_{m,\chi,a}(\cdot, \varphi) \rangle &= \sum_j \langle P_{n,\chi,a}(\cdot, \psi), u_j \rangle \overline{\langle P_{m,\chi,a}(\cdot, \varphi), u_j \rangle} \\ &\quad + \sum_a \frac{1}{4\pi} \int_{-\infty}^\infty \left\langle P_{n,\chi,a}(\cdot, \psi), E_a\left(\cdot, \frac{1}{2} + ir\right) \right\rangle \overline{\left\langle P_{m,\chi,a}(\cdot, \varphi), E_a\left(\cdot, \frac{1}{2} + ir\right) \right\rangle} dr. \end{aligned}$$

Now we must simplify the remaining inner products. Let $f \in \mathcal{L}(N, \chi)$ with Fourier series

$$f(z) = a^+(0) y^{\frac{1}{2}+\nu} + a^-(0) y^{\frac{1}{2}-\nu} + \sum_{n \neq 0} a(n) \sqrt{y} K_\nu(2\pi|n|y) e^{2\pi i n x}.$$

By unfolding the integral in the Petersson inner product and cutting off everything except the diagonal using ?? exactly as in the case for $\langle P_{n,\chi,a}(\cdot, \psi), P_{m,\chi,a}(\cdot, \varphi) \rangle$, we see that

$$\langle P_{n,\chi,a}(\cdot, \psi), f \rangle = \frac{1}{V_\Gamma} \int_0^\infty \overline{a(n) \sqrt{y} K_\nu(2\pi n y)} \psi(y) e^{-4\pi m y} \frac{dy}{y^2}.$$

Now set

$$\omega_\nu(n, \psi) = \frac{1}{V_\Gamma} \int_0^\infty \sqrt{y} K_\nu(2\pi|n|y) \overline{\psi(y)} e^{-4\pi my} \frac{dy}{y^2}.$$

Then it follows from the Fourier series of cusp forms and Eisenstein series that

$$\langle P_{n,\chi,a}(\cdot, \psi), u_j \rangle = \overline{a_j(n)} \omega_{\nu_j}(n, \psi),$$

for $j \geq 1$ and

$$\left\langle P_{n,\chi,a}(\cdot, \psi), E_a\left(\cdot, \frac{1}{2} + ir\right) \right\rangle = \overline{\tau_a\left(n, \frac{1}{2} + ir\right)} \omega_{ir}(n, \psi).$$

In particular, $\langle P_{n,\chi,a}(\cdot, \psi), u_0 \rangle = 0$. So we obtain

$$\begin{aligned} \langle P_{n,\chi,a}(\cdot, \psi), P_{m,\chi,a}(\cdot, \varphi) \rangle &= \sum_{j \geq 1} \overline{a_j(n)} a_j(m) \overline{\omega(n, \psi)} \omega(m, \varphi) \\ &\quad + \sum_a \frac{1}{4\pi} \int_{-\infty}^\infty \overline{\tau_a\left(n, \frac{1}{2} + ir\right)} \tau_a\left(m, \frac{1}{2} + ir\right) \overline{\omega(n, \psi)} \omega(m, \varphi) dr. \end{aligned}$$

This is the second half of the Kuznetsov trace formula. Equating the first and second halves we get the **Kuznetsov trace formula**:

$$\begin{aligned} \delta_{n,m}(\psi, \varphi) + \sum_{\substack{c \geq 1 \\ c \equiv 0 \pmod{N}}} \frac{1}{c} S_\chi(n, m, c) V(n, m, c, \psi, \varphi) &= \sum_{j \geq 1} \overline{a_j(n)} a_j(m) \overline{\omega(n, \psi)} \omega(m, \varphi) \\ &\quad + \sum_a \frac{1}{4\pi} \int_{-\infty}^\infty \overline{\tau_a\left(n, \frac{1}{2} + ir\right)} \tau_a\left(m, \frac{1}{2} + ir\right) \overline{\omega(n, \psi)} \omega(m, \varphi) dr. \end{aligned}$$

The left-hand side is called the **geometric side** and the right-hand side is called the **spectral side**. We collect our work as a theorem:

Theorem 0.1.1 (Kuznetsov trace formula). *Let $\{u_j\}_{j \geq 1}$ be an orthonormal basis of Hecke-Maass eigenforms for $\mathcal{L}(N, \chi)$ of types ν_j with Fourier coefficients $a_j(n)$. Then for any positive integers $n, m \geq 1$, we have*

$$\begin{aligned} \delta_{n,m}(\psi, \varphi) + \sum_{\substack{c \geq 1 \\ c \equiv 0 \pmod{N}}} \frac{1}{c} S_\chi(n, m, c) V(n, m, c, \psi, \varphi) &= \sum_{j \geq 1} \overline{a_j(n)} a_j(m) \overline{\omega(n, \psi)} \omega(m, \varphi) \\ &\quad + \sum_a \frac{1}{4\pi} \int_{-\infty}^\infty \overline{\tau_a\left(n, \frac{1}{2} + ir\right)} \tau_a\left(m, \frac{1}{2} + ir\right) \overline{\omega(n, \psi)} \omega(m, \varphi) dr. \end{aligned}$$

Extra

0.2 Todo: [Dedekind Zeta Functions]

The Definition & Euler Product

We can associate an L -function to every number field. Let K be a number field of degree d and signature (r_1, r_2) . The **Dedekind zeta function** $\zeta_K(s)$ is defined by the following Dirichlet series

$$\zeta_K(s) = \sum_{n \geq 1} \frac{a_K(n)}{n^s}.$$

We will show that $\zeta_K(s)$ is a Selberg class L -function. First we show that the coefficients are polynomially bounded. By ??, $a_K(n) \leq \sigma_0(n)^d \ll_\varepsilon n^\varepsilon$ (recall ??). Therefore $\zeta_K(s)$ is locally absolutely uniformly convergent for $\sigma > 1$. To determine the Euler product, we need to express $\zeta_K(s)$ in another form. Since $\zeta_K(s)$ converges absolutely, we can write

$$\zeta_K(s) = \sum_{\alpha} \frac{1}{N(\alpha)^s},$$

where the sum is over all integral ideals of K . Since the norm is completely multiplicative by ??, the unique factorization of integral ideals implies that we can express $\zeta_K(s)$ as the infinite product

$$\zeta_K(s) = \prod_{\mathfrak{p}} (1 - N(\mathfrak{p})p^{-s})^{-1},$$

for $\sigma > 1$, and where the product is over all prime integral ideals \mathfrak{p} of K . Since each such ideal is above a unique prime p and this product is absolutely convergent for $\sigma > 1$, it can be expressed as

$$\zeta_K(s) = \prod_p \prod_{\mathfrak{p} \text{ above } p} (1 - N(\mathfrak{p})p^{-s})^{-1} = \prod_p \prod_{1 \leq i \leq r_p} (1 - p^{-f_{\mathfrak{p}_i}s})^{-1},$$

where r_p is the number of prime integral ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_{r_p}$ above p . As finitely many primes ramify by ??, ?? implies that all but finitely many primes p satisfy

$$d = \sum_{1 \leq i \leq r_p} f_{\mathfrak{p}_i}.$$

In particular, at least one prime does. Therefore, letting $\omega_{p,i}$ be a primitive $f_{\mathfrak{p}_i}$ -th root of unity, we have the degree d Euler product:

$$\zeta_K(s) = \prod_p \prod_{\substack{1 \leq i \leq r_p \\ 1 \leq k \leq f_{\mathfrak{p}_i}}} (1 - \omega_{p,i}^k p^{-s})^{-1},$$

in this region as well. The local factor at p is

$$\zeta_{K,p}(s) = \prod_{\substack{1 \leq i \leq r_p \\ 1 \leq k \leq f_{\mathfrak{p}_i}}} (1 - \omega_{p,i}^k p^{-s})^{-1},$$

with local roots $\omega_{p,i}^k$.

Todo: [The Integral Representation: Part I]

To find an integral representation for $\zeta_K(s)$, consider the function

$$\omega_K(\mathbf{z}) = \sum_{\mathbf{n} \in \mathbb{Z}_{\geq 0}^d - \{\mathbf{0}\}} \text{Todo: [xxx]} e^{\pi i |\mathbf{n}|^2 z},$$

defined for $\mathbf{z} \in \mathbb{H}^d$.

Todo: [The Hecke Theta Function]

Todo: [The Integral Representation: Part II]

Todo: [The Functional Equation]

Todo: [Imprimitivity]