

# SUBCONVEXITY FOR SECOND MOMENTS OF $L$ -FUNCTIONS TWISTED BY CHARACTERS

NIKOS DIAMANTIS, JEFF HOFFSTEIN, MIN LEE, AND HENRY TWISS

ABSTRACT.

## CONTENTS

1. Introduction	1
2. Preliminaries	1
Sums	1
Forms	2
$L$ -functions	2
3. Shifted Dirichlet Series	2
The Shifted Dirichlet Series $D_{f,g}(s; h, \ell_1, \ell_2)$	2
The Shifted Dirichlet Series $D_{f,v}(w; n, \ell_1, \ell_2)$	2
The Multiple Dirichlet Series $Z_{f,g}(s, v, u, \ell_1, \ell_2)$	2
4. An Amplified Second Moment	2
5. Triple Shifted Sums	2
A Triple Shifted Sum	2
The Diagonal Contribution	2
The Off-diagonal Contribution	2
Balancing	2

## 1. INTRODUCTION

Our aim is to prove the following subconvexity result:

**Theorem 1.1.** *Let  $f$  be a cusp form on  $\mathrm{PSL}_2(\mathbb{Z}) \backslash \mathbb{H}$  of trivial character and let  $\chi$  be a Dirichlet character modulo  $Q$ . Then*

$$L\left(\frac{1}{2}, f \otimes \chi\right) \ll_{\epsilon} Q^{\frac{1}{3} + \frac{\theta}{3} + \epsilon},$$

where  $\theta$  is the best bound towards the Ramanujan-Petersson conjecture for Maass forms.

## 2. PRELIMINARIES

**Sums.** Let  $q \geq 1$  and let  $\chi$  be a Dirichlet character modulo  $q$ . For  $n \in \mathbb{Z}$ , the Ramanujan sum associated to  $\chi$  is

$$c_{\chi}(n) = \sum_{a \bmod q} \chi(a) e^{2\pi i n \frac{a}{q}}.$$

Then  $\tau(\chi) = c_{\chi}(1)$  is the usual Gauss sum associated to  $\chi$ . When  $\chi$  is primitive, we have

$$c_{\chi}(m) = \overline{\chi}(m) \tau(\chi),$$

and  $|\tau(\chi)| = \sqrt{q}$ .

**Forms.**

**$L$ -functions.**

### 3. SHIFTED DIRICHLET SERIES

**The Shifted Dirichlet Series  $D_{f,g}(s; h, \ell_1, \ell_2)$ .**

**The Shifted Dirichlet Series  $D_{f,v}(w; n, \ell_1, \ell_2)$ .**

**The Multiple Dirichlet Series  $Z_{f,g}(s, v, u, \ell_1, \ell_2)$ .**

### 4. AN AMPLIFIED SECOND MOMENT

### 5. TRIPLE SHIFTED SUMS

**A Triple Shifted Sum.**

**The Diagonal Contribution.**

**The Off-diagonal Contribution.**

**Balancing.**