## SUBCONVEXITY FOR SECOND MOMENTS OF L-FUNCTIONS TWISTED BY CHARACTERS

NIKOS DIAMANTIS, JEFF HOFFSTEIN, MIN LEE, AND HENRY TWISS

Abstract.

## Contents

1. Introduction	1
2. Preliminaries	1
Sums	1
Forms	2
L-functions	2
3. Shifted Dirichlet Series	2
The Shifted Dirichlet Series $D_{f,q}(s;h,\ell_1,\ell_2)$	2
The Shifted Dirichlet Series $D_{f,v}(w;n,\ell_1,\ell_2)$	2
The Multiple Dirichlet Series $Z_{f,q}(s,v,u,\ell_1,\ell_2)$	2
4. An Amplified Second Moment	2
5. Triple Shifted Sums	2
A Triple Shifted Sum	2
The Diagonal Contribution	2
The Off-diagonal Contribution	2
Ralancing	9

## 1. Introduction

Our aim is to prove the following subconvexity result:

**Theorem 1.1.** Let f be a cusp form on  $\mathrm{PSL}_2(\mathbb{Z})\backslash\mathbb{H}$  of trivial character and let  $\chi$  be a Dirichlet character modulo Q. Then

$$L\left(\frac{1}{2}, f \otimes \chi\right) \ll_{\epsilon} Q^{\frac{1}{3} + \frac{\theta}{3} + \epsilon},$$

where  $\theta$  is the best bound towards the Ramanujan-Petersson conjecture for Maass forms.

## 2. Preliminaries

**Sums.** Let  $q \ge 1$  and let  $\chi$  be a Dirichlet character modulo q. For  $n \in \mathbb{Z}$ , the Ramanujan sum associated to  $\chi$  is

$$c_{\chi}(n) = \sum_{a \mod q} \chi(a) e^{2\pi i n \frac{a}{q}}.$$

Then  $\tau(\chi) = c_{\chi}(1)$  is the usual Gauss sum associated to  $\chi$ . When  $\chi$  is primitive, we have

$$c_{\chi}(m) = \overline{\chi}(m)\tau(\chi),$$

and  $|\tau(\chi)| = \sqrt{q}$ . When  $\chi$  is the trivial character, we obtain the usual Ramanujan sum

$$c(n) = \sum_{a \mod q} e^{2\pi i n \frac{a}{q}}.$$

Forms.

L-functions.

3. Shifted Dirichlet Series

The Shifted Dirichlet Series  $D_{f,g}(s; h, \ell_1, \ell_2)$ .

The Shifted Dirichlet Series  $D_{f,v}(w;n,\ell_1,\ell_2)$ .

The Multiple Dirichlet Series  $Z_{f,g}(s,v,u,\ell_1,\ell_2)$ .

4. An Amplified Second Moment

5. Triple Shifted Sums

A Triple Shifted Sum.

The Diagonal Contribution.

The Off-diagonal Contribution.

Balancing.