

Arithmetic Functions and Exponential Sums

Henry Twiss

Contents

1	Arithmetic Functions	1
1.1	Arithmetic Functions	1
1.2	Dirichlet Convolution and Möbius Inversion	4
1.3	Todo: [Generating Series]	5
2	Asymptotics	5
2.1	Asymptotics of the Divisor Function	5
3	Dirichlet Characters	6
3.1	Dirichlet Characters	6
3.2	Quadratic Dirichlet Characters	10
4	Exponential Sums	13
4.1	Exponential Sums	13
4.2	Gauss Sums	14
4.3	Quadratic Gauss Sums	18
4.4	Character Sums and the Pólya-Vinogradov Inequality	23

1 Arithmetic Functions

1.1 Arithmetic Functions

Any function

$$f : \mathbb{Z}_+ \rightarrow \mathbb{C},$$

is said to be **arithmetic**. That is, arithmetic functions are maps from the positive integers to the complex numbers. If an arithmetic function f satisfies

$$f(nm) = f(n)f(m) \quad \text{or} \quad f(nm) = f(n) + f(m),$$

whenever n and m are relatively prime, we say f is **multiplicative** or **additive** respectively. If either condition holds regardless of the relative primality of n and m then we say f is **completely multiplicative** or **completely additive**. If f is additive or multiplicative then f is uniquely determined by its values on prime powers and if f is completely additive or

completely multiplicative then it is uniquely determined by its values on primes. Moreover, we have

$$f(1) = 1 \quad \text{or} \quad f(1) = 0,$$

according to if f is multiplicative or additive. Most important arithmetic functions are either multiplicative or additive. Below is a list of the most important arithmetic functions (some of these functions are restrictions of common functions but we define them here as arithmetic functions for completeness):

- (i) The **constant function**: The function $\mathbf{1}(n)$. This function is neither additive nor multiplicative.
- (ii) The **indicator function**: The function $\varepsilon(n)$ defined by

$$\varepsilon(n) = \begin{cases} 1 & \text{if } n = 1, \\ 0 & \text{if } n \geq 2. \end{cases}$$

This function is completely multiplicative.

- (iii) The **identity function**: The function $\text{id}(n)$. This function is completely multiplicative.
- (iv) The **logarithm**: The function $\log(n)$. This function is completely additive.
- (v) The **Möbius function**: The function $\mu(n)$ defined by

$$\mu(n) = \begin{cases} 1 & \text{if } n \text{ is square-free with an even number of prime factors,} \\ -1 & \text{if } n \text{ is square-free with an odd number of prime factors,} \\ 0 & \text{if } n \text{ is not square-free.} \end{cases}$$

This function is multiplicative.

- (vi) The **characteristic function of square-free integers**: The square of the Möbius function $\mu^2(n)$. This function is multiplicative.
- (vii) **Liouville's function**: The function $\lambda(n)$ defined by

$$\lambda(n) = \begin{cases} 1 & \text{if } n = 1, \\ (-1)^k & \text{if } n \text{ is composed of } k \text{ not necessarily distinct prime factors.} \end{cases}$$

This function is completely multiplicative.

- (viii) **Euler's totient function**: The function $\varphi(n)$ defined by

$$\varphi(n) = \sum_{\substack{a \pmod n \\ (a,n)=1}} 1.$$

This function is multiplicative.

(ix) The **divisor function**: The function $\sigma_0(n)$ defined by

$$\sigma_0(n) = \sum_{d|n} 1.$$

This function is multiplicative.

(x) The **sum of divisors function**: The function $\sigma_1(n)$ defined by

$$\sigma_1(n) = \sum_{d|n} d.$$

This function is multiplicative.

(xi) The **generalized sum of divisors function**: The function $\sigma_s(n)$ defined by

$$\sigma_s(n) = \sum_{d|n} d^s,$$

for any $s \in \mathbb{C}$. This function is multiplicative.

(xii) The **number of distinct prime factors function**: The function $\omega(n)$ defined by

$$\omega(n) = \sum_{p|n} 1.$$

This function is additive.

(xiii) The **total number of prime divisors function**: The function $\Omega(n)$ defined by

$$\Omega(n) = \sum_{p^m|n} 1.$$

This function is completely additive.

(xiv) The **von Mangoldt function**: The function $\Lambda(n)$ defined by

$$\Lambda(n) = \begin{cases} 0 & \text{if } n \text{ is not a prime power,} \\ \log(p) & \text{if } n = p^m \text{ for some prime } p \text{ and positive integer } m. \end{cases}$$

This function is neither additive or multiplicative.

Much of number theory is concerned with estimating these functions, or sums thereof, in terms of elementary functions.

1.2 Dirichlet Convolution and Möbius Inversion

If f and g are two arithmetic functions then we can define a new arithmetic function $f * g$ called their **Dirichlet convolution** defined by

$$(f * g)(n) = \sum_{d|n} f(d)g\left(\frac{n}{d}\right).$$

As the sum is over all divisors, Dirichlet convolution is a symmetric operation. Dirichlet convolution preserves multiplicative functions.

Proposition 1.1. *If f and g are multiplicative arithmetic functions then their Dirichlet convolution $f * g$ is as well.*

Proof. Let n and m be relatively prime positive integers. Every divisor d of nm is of the form $d = d'd''$ with d' a divisor of n , d'' a divisor of m , and d' and d'' relatively prime. A short computation shows

$$\sum_{d|nm} f(d)g\left(\frac{nm}{d}\right) = \left(\sum_{d'|n} f(d')g\left(\frac{n}{d'}\right)\right) \left(\sum_{d''|m} f(d'')g\left(\frac{m}{d''}\right)\right),$$

which is equivalent to the claim. □

This result makes the set of multiplicative functions into a commutative semigroup under Dirichlet convolution. It is actually a commutative monoid since the indicator function ε acts as an identity. This means

$$f * \varepsilon = f.$$

A certain case of interest for Dirichlet convolution is when the Möbius function is convolved with the constant function.

Proposition 1.2. *We have*

$$\sum_{d|n} \mu(d) = \begin{cases} 1 & \text{if } n = 1, \\ 0 & \text{if } n \geq 2. \end{cases}$$

In particular,

$$\mu * \mathbf{1} = \varepsilon.$$

Proof. The first statement is equivalent to the second, so it suffices to prove the first. The sum $\sum_{d|n} \mu(d)$ is multiplicative by Proposition 1.1 so we may assume $n = p^r$ is a nonnegative power of a prime. When $r = 0$, $d = 1$ and the sum is 1. When $r \geq 1$, d runs over $1, p, \dots, p^r$. Every value is zero except $\mu(1) = 1$ and $\mu(p) = -1$. This proves that the sum is zero. □

With this result we can prove **Möbius inversion**:

Theorem (Möbius inversion). *If f and g are arithmetic functions, then*

$$g(n) = \sum_{d|n} f(d) \quad \text{if and only if} \quad f(n) = \sum_{d|n} g(d) \mu\left(\frac{n}{d}\right).$$

In particular,

$$g = f * \mathbf{1} \quad \text{if and only if} \quad f = g * \mu.$$

Proof. As the first statement is equivalent to the second, we will prove the second statement. Convolving $g = f * \mathbf{1}$ with μ gives $f = g * \mu$ in view of Proposition 1.2. This proves the forward implication. The reverse implication follows by convolving $f = g * \mu$ with $\mathbf{1}$ and arguing analogously. $\square$

1.3 Todo: [Generating Series]

2 Asymptotics

2.1 Asymptotics of the Divisor Function

It is essential to have a bound for the number of divisors $\sigma_0(n)$ of a positive integer n . A sufficient estimate is that $\sigma_0(n)$ grows slower than any polynomial.

Proposition 2.1.

$$\sigma_0(n) \ll_{\varepsilon} n^{\varepsilon}.$$

Proof. Let $n = p_1^{r_1} \cdots p_k^{r_k}$ be the prime factorization of n so that $\sigma_0(n) = (r_1 + 1) \cdots (r_k + 1)$. Then

$$\frac{\sigma_0(n)}{n^{\varepsilon}} = \prod_i \frac{r_i + 1}{p_i^{r_i \varepsilon}}.$$

It suffices to show that the right-hand side is $O_{\varepsilon}(1)$. For a prime p , consider the nonnegative continuous function $f_{p,\varepsilon}(x)$ defined by

$$f_{p,\varepsilon}(x) = \frac{x + 1}{p^{x\varepsilon}},$$

for $x \geq -1$. The derivative is easily seen to be

$$f'_{p,\varepsilon}(x) = \frac{1 - \log(p^{\varepsilon})(x + 1)}{p^{x\varepsilon}},$$

and is negative for $\frac{1}{\log(p^{\varepsilon})} - 1 < x$. Therefore $f_{p,\varepsilon}(x)$ is eventually decreasing and so attains a maximum positive value. In particular, it attains some maximal positive value at a positive integer a_p amongst all positive integers because $f_{p,\varepsilon}(-1) = 0$ and $f_{p,\varepsilon}(x)$ is positive for $x > -1$. Then the inequalities

$$f_{p,\varepsilon}(a_p - 1) \leq f_{p,\varepsilon}(a_p) \quad \text{and} \quad f_{p,\varepsilon}(a_p + 1) \leq f_{p,\varepsilon}(a_p),$$

are

$$\frac{a_p}{p^{(a_p-1)\varepsilon}} \leq \frac{a_p+1}{p^{a_p\varepsilon}} \quad \text{and} \quad \frac{a_p+2}{p^{(a_p+1)\varepsilon}} \leq \frac{a_p+1}{p^{a_p\varepsilon}},$$

respectively. Upon isolating a_p to obtain an upper bound in the first inequality and a lower bound in the second inequality, a short computation shows

$$\frac{1}{p^\varepsilon - 1} - 1 \leq a_p \leq \frac{1}{p^\varepsilon - 1}.$$

Therefore we may take $a_p = \left\lfloor \frac{1}{p^\varepsilon - 1} - 1 \right\rfloor$. Moreover,

$$\frac{\frac{1}{p^\varepsilon - 1}}{p^{\varepsilon(\frac{1}{p^\varepsilon - 1} - 1)}} \leq f_p(a_p) \leq \frac{\frac{1}{p^\varepsilon - 1} + 1}{p^{\varepsilon(\frac{1}{p^\varepsilon - 1}})}.$$

Taking the logarithm yields

$$\log\left(\frac{1}{p^\varepsilon - 1}\right) - \log(p^\varepsilon) \left(\frac{1}{p^\varepsilon - 1} - 1\right) \leq \log(f_p(a_p)) \leq \log\left(\frac{1}{p^\varepsilon - 1} + 1\right) - \log(p^\varepsilon) \left(\frac{1}{p^\varepsilon - 1}\right).$$

Taking the limit as $p \rightarrow \infty$, a short computation shows that both the left-hand and right-hand sides tend to zero. Therefore $\log(f_p(a_p)) \rightarrow 0$ and hence $f_p(a_p) \rightarrow 1$. This guarantees that the infinite product $\prod_p f_{p,\varepsilon}(a_p)$ converges to some positive value whence is $O_\varepsilon(1)$. As

$$\prod_i \frac{r_i + 1}{p_i^{r_i\varepsilon}} \leq \prod_p f_{p,\varepsilon}(a_p),$$

the claim follows. □

3 Dirichlet Characters

Throughout, for any integers a and m the symbol $\left(\frac{a}{m}\right)$ will stand for the Kronecker symbol. In particular, it is the Jacobi symbol when m is a positive odd integer, and the Legendre symbol when $m = p$ is a prime. If a is an integer modulo m with $(a, m) = 1$ we write a^{-1} for the inverse of a modulo m . Also, for any complex z we set

$$e(z) = e^{2\pi iz}.$$

3.1 Dirichlet Characters

One of the most important types of multiplicative functions are those that are periodic. These are the Dirichlet characters. Let m be a positive integer. A multiplicative homomorphism

$$\chi : \mathbb{Z} \rightarrow \mathbb{C},$$

is said to be a **Dirichlet character** modulo m if it is m -periodic and such that $\chi(a) = 0$ if and only if $(a, m) > 1$. We call m the **modulus** of χ . A Dirichlet character is necessarily a completely multiplicative arithmetic function when restricted to the positive integers.

We say a Dirichlet character χ is **principal** if it only takes values 0 or 1. There is always a unique principal Dirichlet character modulo m , denoted $\chi_{m,0}$, defined by

$$\chi_{m,0}(a) = \begin{cases} 1 & (a, m) = 1, \\ 0 & (a, m) > 1. \end{cases}$$

When the modulus is 1, the principal Dirichlet character is identically 1 and we call this the **trivial Dirichlet character**. This is the only Dirichlet character modulo 1.

By Euler's little theorem, $a^{\varphi(m)} \equiv 1 \pmod{m}$ whenever $(a, m) = 1$. Multiplicativity forces $\chi(a)^{\varphi(m)} = 1$ and so the nonzero values of any Dirichlet character modulo m are $\varphi(m)$ -th roots of unity. This implies that there are only finitely many Dirichlet characters of any fixed modulus. Given two Dirichlet character χ and ψ modulo m , the functions

$$\chi\psi : \mathbb{Z} \rightarrow \mathbb{C} \quad \text{and} \quad \bar{\chi} : \mathbb{Z} \rightarrow \mathbb{C},$$

are also Dirichlet characters modulo m . This turns the set of such Dirichlet characters into an abelian group denote by X_m where the identity is the principal Dirichlet character modulo m and the inverse is given by the conjugate as the nonzero values of Dirichlet characters are roots of unity.

This is all strikingly similar to characters on $(\mathbb{Z}/m\mathbb{Z})^*$ and there is indeed a connection. By the multiplicativity and m -periodicity of χ , it induces a character of $(\mathbb{Z}/m\mathbb{Z})^*$. Conversely, if we are given a character on $(\mathbb{Z}/m\mathbb{Z})^*$ we can extend it to a Dirichlet character χ by defining it to be m -periodic with $\chi(a) = 0$ if $(a, m) > 1$. We call this extension a **zero extension**. This argument shows that Dirichlet characters modulo m are exactly the zero extensions of characters on $(\mathbb{Z}/m\mathbb{Z})^*$. As abelian groups are isomorphic to their character groups, we deduce that the group of Dirichlet characters modulo m is isomorphic to $(\mathbb{Z}/m\mathbb{Z})^*$. Therefore there are $\varphi(m)$ Dirichlet characters modulo m and we identify them with the characters on $(\mathbb{Z}/m\mathbb{Z})^*$. Just as for characters of abelian groups, we have orthogonality relations called the **Dirichlet orthogonality relations**.

Proposition (Dirichlet orthogonality relations). *Let χ and ψ be Dirichlet characters modulo m and let $a, b \in (\mathbb{Z}/m\mathbb{Z})^*$. Then*

$$\sum_{a \pmod{m}} (\chi\bar{\psi})(a) = \varphi(m)\delta_{\chi,\psi} \quad \text{and} \quad \sum_{\chi \pmod{m}} \chi(ab^{-1}) = \varphi(m)\delta_{a,b}.$$

In particular,

$$\sum_{a \pmod{m}} \chi(a) = \varphi(m)\delta_{\chi,\chi_{m,0}} \quad \text{and} \quad \sum_{\chi \pmod{m}} \chi(a) = \varphi(m)\delta_{a,1}.$$

Proof. If $\chi = \psi$ then the first sum is clearly $\phi(m)$. If not, let $b \in (\mathbb{Z}/m\mathbb{Z})^*$ be such that $(\chi\bar{\psi})(b) \neq 1$. A short computation shows

$$(\chi\bar{\psi})(b) \sum_{a \pmod{m}} (\chi\bar{\psi})(a) = \sum_{a \pmod{m}} (\chi\bar{\psi})(a),$$

in which case the sum vanishes. This proves the first identity. For the second, if $a = b$ then the second sum is clearly $\varphi(m)$. If $a \neq b$, we claim that there exists a Dirichlet character ψ modulo m with $\psi(ab^{-1}) \neq 1$. Set $g = ab^{-1}$. The cyclic subgroup $\langle g \rangle$ of $(\mathbb{Z}/m\mathbb{Z})^*$ has some order $d > 1$. Consider the homomorphism

$$\psi_d : \langle g \rangle \rightarrow \mathbb{C} \quad g^k \mapsto e\left(\frac{k}{d}\right).$$

By the structure theorem for finite abelian groups, $(\mathbb{Z}/m\mathbb{Z})^* \cong \langle g \rangle \times H$ for some subgroup H . Whence we define a Dirichlet character ψ modulo m by

$$\psi : (\mathbb{Z}/m\mathbb{Z})^* \rightarrow \mathbb{C} \quad g^k h \mapsto e\left(\frac{k}{d}\right).$$

This ψ is the desired Dirichlet character. A short computation shows

$$\psi(ab^{-1}) \sum_{\chi \pmod{m}} \chi(ab^{-1}) = \sum_{\chi \pmod{m}} \chi(ab^{-1}),$$

in which case the sum vanishes. This proves the second identity and the first statement in its entirety. The second statement follows from the first upon taking $\psi = \chi_{m,0}$ and $b = 1$ respectively. $\square$

It is possible for Dirichlet characters of a fixed modulus to arise from Dirichlet characters of a smaller modulus. Suppose χ and χ^* are Dirichlet characters modulo m and d respectively with $d \mid m$. We say χ is induced **induced** from χ^* if

$$\chi(a) = \begin{cases} \chi^*(a) & \text{if } (a, m) = 1, \\ 0 & \text{if } (a, m) > 1. \end{cases}$$

This means χ is a Dirichlet character whose values are given by those of χ^* . Necessarily χ is d -periodic and its nonzero values are $\varphi(d)$ -th roots of unity. We say a Dirichlet character is **primitive** if it is not induced by any Dirichlet character other than itself and **imprimitive** otherwise. The principal Dirichlet characters are precisely those induced from the trivial Dirichlet character and the only primitive one is the trivial Dirichlet character itself. Moreover, a Dirichlet character is primitive if and only if its conjugate is. Our primary aim will be to show that every Dirichlet character is induced from a unique primitive Dirichlet character.

Theorem 3.1. *Suppose χ is a Dirichlet character modulo m . There exists a unique primitive Dirichlet character $\tilde{\chi}$ such that χ is induced from $\tilde{\chi}$.*

Proof. Let q be the positive integer given by

$$q = \min\{d \mid m : \chi(a) = \chi(b) \text{ for } a \equiv b \pmod{d} \text{ with } (ab, m) = 1\},$$

and let $\tilde{\chi} : \mathbb{Z} \rightarrow \mathbb{C}$ be defined by

$$\tilde{\chi}(a) = \begin{cases} \chi(a) & \text{if } (a, q) = 1, \\ 0 & \text{if } (a, q) > 1. \end{cases}$$

The definition of q implies $\tilde{\chi}$ is well-defined and hence q -periodic. In fact, $\tilde{\chi}$ is a Dirichlet character modulo q and minimality forces $\tilde{\chi}$ to be primitive. By construction χ is induced from $\tilde{\chi}$ and this proves existence. Now suppose $\tilde{\chi}_1$ and $\tilde{\chi}_2$ are two primitive Dirichlet characters modulo q_1 and q_2 respectively both of which induce χ . Then $\tilde{\chi}_1(a) = \tilde{\chi}_2(a)$ whenever $(a, m) = 1$. Setting $q = (q_1, q_2)$, we also have $\tilde{\chi}_1(a) = \tilde{\chi}_2(a)$ whenever $(a, q) = 1$. Hence $\tilde{\chi}_1$ and $\tilde{\chi}_2$ are both induced from the same Dirichlet character modulo q . Primitivity implies $q_1 = q_2$ and $\tilde{\chi}_1 = \tilde{\chi}_2$. This proves uniqueness. $\square$

In light of this result, we define the **conductor** q of a Dirichlet character χ modulo m to be the modulus of the unique primitive Dirichlet character $\tilde{\chi}$ inducing χ . By the proof, the conductor is given by

$$q = \min\{d \mid m : \chi(a) = \chi(b) \text{ for } a \equiv b \pmod{d} \text{ with } (ab, m) = 1\}.$$

Also observe that χ is q -periodic, q is the minimal period of χ , and the nonzero values of χ are $\varphi(q)$ -th roots of unity. Moreover,

$$\chi = \tilde{\chi} \chi_{\frac{m}{q}, 0},$$

and χ is primitive if and only if its conductor and modulus are equal.

As not every Dirichlet character of a fixed modulus is primitive, it is natural to ask how many primitive Dirichlet characters there are for a given modulus. Let $N(m)$ be the number of primitive Dirichlet characters modulo m . Then $N(m)$ is easily determined via Möbius inversion.

Proposition 3.2. *For any positive integer m , we have*

$$\phi(m) = \sum_{d|m} N(d),$$

where $N(d)$ be the number of primitive Dirichlet characters modulo d . In particular, $N(m)$ is given by

$$N(m) = \sum_{d|m} \phi(d) \mu\left(\frac{m}{d}\right).$$

Proof. To prove the first formula, the right-hand side counts the number of Dirichlet characters modulo m since every such Dirichlet character is induced from a unique primitive Dirichlet character whose modulus divides m by Theorem 3.1. The left hand side also counts the number of Dirichlet characters modulo m as are $\phi(m)$ many. This proves the first formula. The second follows by Möbius inversion. $\square$

Primitive Dirichlet characters also behave well with respect to multiplication if the conductors are relatively prime as the following proposition shows:

Proposition 3.3. *Suppose χ_1 and χ_2 are Dirichlet characters modulo m_1 and m_2 respectively with $(m_1, m_2) = 1$. Set $\chi = \chi_1 \chi_2$. Then χ is a primitive if and only if χ_1 and χ_2 both are.*

Proof. By construction, χ is a Dirichlet character modulo m_1m_2 . Let q_1 and q_2 be the conductors of χ_1 and χ_2 respectively and let q be the conductor of χ . Then χ is q_1q_2 -periodic and $q \mid q_1q_2$.

For the forward implication, suppose χ is primitive so that $q = m_1m_2$ whence $m_1m_2 \mid q_1q_2$. This forces $m_1 = q_1$ and $m_2 = q_2$ proving χ_1 and χ_2 are both primitive. For the reverse implication, suppose χ_1 and χ_2 are both primitive so that $q_1 = m_1$ and $q_2 = m_2$. Then $(q_1, q_1) = 1$ and the Chinese remainder theorem implies that χ is q_1 -periodic on those integers that are congruent to 1 modulo q_2 and q_2 -periodic on those integers that are congruent to 1 modulo q_1 . This forces $q_1 \mid q$ and $q_2 \mid q$ which together imply $q = q_1q_2$ and thus χ is primitive. $\square$

We would now like to distinguish Dirichlet characters based on their nonzero values. We say χ is **real** if it is real-valued. This means the nonzero values of χ are 1 or -1 since they are the only real roots of unity. We say χ is **complex** if it is not real. More commonly, we distinguish Dirichlet characters by the roots of unity that their nonzero values take. We say χ is of **order** n if the nonzero values of χ are all n -th roots of unity. In the cases of small order we will often use the latin derived names **quadratic**, **cubic**, etc. To connect these two naming conventions observe that a nontrivial Dirichlet character is quadratic if and only if it is real and an other nontrivial Dirichlet character is necessarily complex. Also note that quadratic Dirichlet characters are their own conjugates.

We will also distinguish Dirichlet characters by their parity. By multiplicativity, we must have $\chi(-1) = \pm 1$. Accordingly, we say χ is **even** if $\chi(-1) = 1$ and **odd** if $\chi(-1) = -1$. Then even Dirichlet characters are even functions while odd Dirichlet characters are odd functions. Note that conjugate and induced Dirichlet characters necessarily have the same parity. The parity is also expressed via the formula

$$\frac{\chi(1) - \chi(-1)}{2} = \begin{cases} 0 & \text{if } \chi \text{ is even,} \\ 1 & \text{if } \chi \text{ is odd.} \end{cases}$$

3.2 Quadratic Dirichlet Characters

Quadratic Dirichlet characters deserve special attention as it is possible to classify all of them explicitly. This is due to the fact that they arise from Jacobi and Kronecker symbols. For a positive odd integer m , define

$$\chi_m(a) = \left(\frac{a}{m} \right).$$

By definition of the Jacobi symbol, χ_m becomes a quadratic Dirichlet character modulo m . Unfortunately, the quadratic Dirichlet characters constructed in this manner do not exhaust all possible examples. To accomplish this we need to use Kronecker symbols. An integer D is said to be a **fundamental discriminant** if it is of the form

$$D = \begin{cases} d & \text{if } D \equiv 1 \pmod{4}, \\ 4d & \text{if } D \equiv 8, 12 \pmod{16}, \end{cases}$$

for some square-free integer d . Necessarily $d \equiv 1 \pmod{4}$ or $d \equiv 2, 3 \pmod{4}$ respectively and thus is nonzero. We define $\chi_D : \mathbb{Z} \rightarrow \mathbb{C}$ by

$$\chi_D(a) = \left(\frac{D}{a} \right).$$

It turns out that χ_D defines a primitive quadratic Dirichlet character modulo $|D|$, provided $D \neq 1$, and exhausts all such possibilities.

Theorem 3.4. *If D is a fundamental discriminant and $D \neq 1$ then χ_D is a primitive quadratic Dirichlet character of conductor $|D|$. Moreover, all primitive quadratic Dirichlet characters are of this form.*

Proof. We first show that χ_D is a primitive quadratic Dirichlet character modulo $|D|$. If $D \equiv 1 \pmod{4}$, the sign in quadratic reciprocity is always 1. Then

$$\chi_D(a) = \left(\frac{a}{|D|} \right),$$

which is a Dirichlet character modulo $|D|$. If $D \equiv 12 \pmod{16}$ then $\frac{D}{4} \equiv 3 \pmod{4}$ and the sign in quadratic reciprocity is $\left(\frac{-1}{a} \right)$ which is the primitive quadratic Dirichlet character modulo 4 as there are only two such Dirichlet characters and $\left(\frac{-1}{a} \right)$ is non-principal. Whence

$$\chi_D(a) = \left(\frac{-1}{a} \right) \left(\frac{a}{\left| \frac{D}{4} \right|} \right),$$

which is a Dirichlet character modulo $|D|$. If $D \equiv 8 \pmod{16}$ first observe that $\left(\frac{D}{a} \right) = \left(\frac{8}{a} \right) \left(\frac{\frac{D}{8}}{a} \right)$ where $\left(\frac{8}{a} \right)$ is one of the two primitive quadratic Dirichlet character modulo 8 (the other is $\left(\frac{-8}{a} \right)$). As $\frac{D}{8} \equiv 1, 3 \pmod{4}$, the sign in quadratic reciprocity is either 1 or $\left(\frac{-1}{a} \right)$ according to these two cases. Thus

$$\chi_D(a) = \left(\frac{8}{a} \right) \left(\frac{a}{\left| \frac{D}{8} \right|} \right) \quad \text{or} \quad \chi_D(a) = \left(\frac{-8}{a} \right) \left(\frac{a}{\left| \frac{D}{8} \right|} \right),$$

according to if $\frac{D}{8} \equiv 1, 3 \pmod{4}$ respectively, and in either case is a Dirichlet character modulo $|D|$. We can compactly express all of these cases as follows:

$$\chi_D(a) = \begin{cases} \left(\frac{a}{|D|} \right) & \text{if } D \equiv 1 \pmod{4}, \\ \left(\frac{-1}{a} \right) \left(\frac{a}{\left| \frac{D}{4} \right|} \right) & \text{if } \frac{D}{4} \equiv 3 \pmod{4}, \\ \left(\frac{8}{a} \right) \left(\frac{a}{\left| \frac{D}{8} \right|} \right) & \text{if } \frac{D}{8} \equiv 1 \pmod{4}, \\ \left(\frac{-8}{a} \right) \left(\frac{a}{\left| \frac{D}{8} \right|} \right) & \text{if } \frac{D}{8} \equiv 3 \pmod{4}. \end{cases}$$

This proves χ_D is a Dirichlet character modulo $|D|$. It is not hard to see that χ_D is primitive. Indeed, we have already mentioned that the characters $\left(\frac{-1}{a} \right)$, $\left(\frac{8}{a} \right)$, and $\left(\frac{-8}{a} \right)$ are all primitive.

Since D , $\frac{D}{4}$, and $\frac{D}{8}$ are square-free according to their equivalences modulo 4 and $D \neq 1$, it suffices to show that χ_p is primitive for primes p with $p \neq 2$ by Proposition 3.3. This is immediate since p is prime and χ_p is non-principal.

We now show that every primitive quadratic Dirichlet character is of the form χ_D for some fundamental discriminant D . By Proposition 3.3 again, it suffices to consider primitive quadratic Dirichlet character modulo a prime power p^m .

First suppose $p \neq 2$. Then $(\mathbb{Z}/p^m\mathbb{Z})^*$ is cyclic, generated by say g , and every $a \in (\mathbb{Z}/p^m\mathbb{Z})^*$ is of the form $a = g^\nu$ for some $\nu \in (\mathbb{Z}/\varphi(p^m)\mathbb{Z})$. It follows that every corresponding Dirichlet character χ is of the form

$$\chi(a) = e\left(\frac{k\nu}{\varphi(p^m)}\right),$$

for an integer k modulo $\varphi(p^m)$. Moreover, χ is primitive if and only if $k \not\equiv 0 \pmod{p}$ for otherwise χ is a Dirichlet character modulo p^{m-1} . Similarly, χ is quadratic if and only if $k \equiv \frac{\varphi(p^m)}{2} \pmod{\varphi(p^m)}$. Such a k exists and is unique because $p \neq 2$. We also see that if χ is quadratic then it is imprimitive unless $m = 1$ for then $\varphi(p) = p - 1$ which is not a multiple of p . To summarize, there is a unique quadratic Dirichlet character modulo p^m and it is primitive if and only if $m = 1$. Necessarily, this unique primitive quadratic Dirichlet character modulo p is given by χ_D for the fundamental discriminant $D = p$ if $p \equiv 1 \pmod{4}$ and $D = -p$ if $p \equiv 3 \pmod{4}$.

Now suppose $p = 2$ so that $p^m = 2^m$. If $m = 1$ then $\varphi(2) = 1$ and there are no primitive quadratic Dirichlet characters as the only Dirichlet character is principal. If $m = 2$ then $\varphi(4) = 2$ so that there are two Dirichlet characters. They are both quadratic but only one is primitive namely the aforementioned $\left(\frac{-1}{a}\right)$. This primitive quadratic Dirichlet character is given by χ_D for the fundamental discriminant $D = -4$. For $m \geq 3$ there is an isomorphism $(\mathbb{Z}/2^m\mathbb{Z})^* \cong C_2 \times C_{2^{m-2}}$ where C_2 and $C_{2^{m-2}}$ are the cyclic groups of order 2 and 2^{m-2} respectively. Therefore every $a \in (\mathbb{Z}/2^m\mathbb{Z})^*$ is of the form $a = (-1)^\mu 5^\nu$ for some $\mu \in \mathbb{Z}/2\mathbb{Z}$ and $\nu \in \mathbb{Z}/2^{m-2}\mathbb{Z}$. Then every corresponding Dirichlet character χ is of the form

$$\chi(a) = e\left(\frac{j\mu}{2}\right) e\left(\frac{k\nu}{2^{m-2}}\right),$$

for integers j modulo 2 and k modulo 2^{m-2} . Similarly to the case for $p \neq 2$, χ is primitive if and only if $k \not\equiv 0 \pmod{2^{m-2}}$. Moreover, χ is quadratic if and only if $k \equiv 0 \pmod{2^{m-3}}$. These congruences together imply that a primitive quadratic Dirichlet character exists if and only if $m = 3$. In this case there are four Dirichlet characters. They are all quadratic but only two are primitive, namely the aforementioned $\left(\frac{8}{a}\right)$ and $\left(\frac{-8}{a}\right)$. These two primitive quadratic Dirichlet characters are given by χ_D for the fundamental discriminants $D = 8$ and $D = -8$ respectively. $\square$

It follows from Theorem 3.4 that all quadratic Dirichlet characters are induced from some χ_D including $D = 1$ since this corresponds to the trivial Dirichlet character. In particular, so too are the quadratic Dirichlet characters given by Jacobi symbols.

4 Exponential Sums

Throughout, for any integers a and m the symbol $\left(\frac{a}{m}\right)$ will stand for the Kronecker symbol. In particular, it is the Jacobi symbol when m is a positive odd integer, and the Legendre symbol when $m = p$ is a prime. If a is an integer modulo m with $(a, m) = 1$ we write a^{-1} for the inverse of a modulo m . Also, for any complex z we set

$$e(z) = e^{2\pi iz}.$$

4.1 Exponential Sums

An **exponential sum** S_f is a sum of the form

$$S_f = \sum_{a \in A} e(f(a)),$$

where A is a finite set of integers and f is a complex function on A . In many instances, f is an arithmetic function so that A may be any finite set of positive integers. In any case, it is fruitful to view S_f as a geometric walk in $\mathbb{C}$ where the a -th step is in the direction $e(f(a))$. Taking absolute values, we always have the trivial bound

$$S_f \leq |A|.$$

However, since the summands all occur on the unit circle, we should expect lots of oscillatory cancellation if the phases are essentially random. In the optimal case, suppose the $e(f(a))$ model independent and identically distributed random variables with mean zero. Heuristically speaking, this walk should exhibit square-root cancellation and we expect a bound of the form

$$S_f \ll |A|^{\frac{1}{2} + \varepsilon}.$$

Obtaining such estimates is highly dependent upon the particular choices of f and A . Often we will have A depend upon various parameters and will want to obtain some type of cancellation in one or more of these parameters.

A particular type of exponential sum, where we expect close to square-root cancellation, are those of complete exponential sums. To define them, fix a integer m and let f be a complex function on $(\mathbb{Z}/m\mathbb{Z})$ or $(\mathbb{Z}/m\mathbb{Z})^*$. Then the corresponding exponential sum

$$S_{f,m} = \sum_{a \pmod{m}} e(f(a)) \quad \text{or} \quad S_{f,m} = \sum_{\substack{a \pmod{m} \\ (a,m)=1}} e(f(a)),$$

is said to be **complete**. In contrast, an exponential sum not of this form is said to be **incomplete**. The upshot of these exponential sums is that they are over a finite additive or multiplicative abelian group respectively. As is often demonstrated by example, many complete exponential sums exhibit close to square-root cancellation. We will see examples of this shortly. For a common example of an incomplete exponential sum, let M and N be nonnegative integers and consider

$$S_f(M, N) = \sum_{M \leq n \leq M+N} e(f(n)).$$

We will see an example of an incomplete sub exhibiting square-root cancellation, but often this is far from true.

4.2 Gauss Sums

Let χ be a Dirichlet character modulo m . For any integer b , the **Gauss sum** $G(b, \chi)$ associated to χ is given by

$$G(b, \chi) = \sum_{a \pmod{m}} \chi(a) e\left(\frac{ab}{m}\right).$$

This is a complete exponential sum. When $b \mid m$ the summands are all 1 and the Dirichlet orthogonality relations imply

$$G(b, \chi) = \varphi(m) \delta_{\chi, \chi_{m,0}}.$$

When $b = 1$ we simply write $G(\chi) = G(1, \chi)$. The simplest case of a Gauss sum occurs when $\chi = \chi_{m,0}$ in which case we write $R(b, m) = G(b, \chi_{m,0})$ and call this the **Ramanujan sum** modulo m . It can be expressed as

$$R(b, m) = \sum_{\substack{a \pmod{m} \\ (a, m) = 1}} e\left(\frac{ab}{m}\right).$$

When $b \mid m$ the summands are all 1 and the Ramanujan sum has the simple evaluation

$$R(b, m) = \varphi(m).$$

For a general index the Ramanujan sums can be computed explicitly by means of the Möbius function.

Proposition 4.1. *For integers b and m with m positive, we have*

$$R(b, m) = \sum_{d \mid (b, m)} d \mu\left(\frac{m}{d}\right).$$

Proof. The identity is obvious if $m = 1$ since the Ramanujan sum is 1. So assume $m > 1$. Every a modulo m is of the form $a = a'd$ for some divisor d of m and a' modulo $\frac{m}{d}$ with $(a', \frac{m}{d}) = 1$. So summing $R(b, d)$ over the divisors d of m gives

$$\sum_{d \mid m} R(b, d) = \sum_{a \pmod{m}} e\left(\frac{ab}{m}\right).$$

If $m \mid b$ the latter sum is m while if $m \nmid b$ the sum vanishes as it is the sum of all m -th roots of unity. Thus

$$\sum_{d \mid m} R(b, d) = \begin{cases} m & \text{if } m \mid b, \\ 0 & \text{if } m \nmid b. \end{cases}$$

Now apply Möbius inversion. □

This explicit evaluation is not typical of a larger reality. Gauss sums, in general, are extremely difficult to compute exactly. Nevertheless, we can develop many useful properties which help to reduce their evaluation.

Proposition 4.2. *Let χ and ψ be nontrivial Dirichlet characters modulo m and n respectively and let b be an integer. Then the following properties hold:*

- (i) $\overline{G(b, \bar{\chi})} = \chi(-1)G(b, \chi)$.
- (ii) If $(b, m) = 1$ then $G(b, \chi) = \bar{\chi}(b)G(\chi)$.
- (iii) If $(b, m) > 1$ and χ is primitive then $G(b, \chi) = 0$.
- (iv) If $(m, n) = 1$ then $G(b, \chi\psi) = \chi(n)\psi(m)G(b, \chi)G(b, \psi)$.
- (v) If $\tilde{\chi}$ is the primitive Dirichlet character of conductor q inducing χ , then

$$G(\chi) = \mu\left(\frac{m}{q}\right) \tilde{\chi}\left(\frac{m}{q}\right) G(\tilde{\chi}).$$

Proof. We will prove the properties separately.

- (i) This is equivalent to the identity

$$\sum_{a \pmod{m}} \chi(a) e\left(-\frac{ab}{m}\right) = \chi(-1) \sum_{a \pmod{m}} \chi(a) e\left(\frac{ab}{m}\right),$$

which is clear upon making the change of variables $a \mapsto -a$.

- (ii) This is equivalent to the identity

$$\sum_{a \pmod{m}} \chi(a) e\left(\frac{ab}{m}\right) = \bar{\chi}(b) \sum_{a \pmod{m}} e\left(\frac{a}{m}\right),$$

which is clear under the change of variables $a \mapsto ab^{-1}$.

- (iii) Observe that if $(b, m) > 1$ then $m > 1$. We will separate the proof into two cases depending on if m is prime or not. In the first case, $m = p$ for a prime p . Then $p \mid b$ whence

$$G(b, \chi) = \sum_{a \pmod{m}} \chi(a).$$

Therefore $G(b, \chi) = 0$ by the orthogonality relations because χ is primitive. In the second case, we have $p \mid (b, m)$ for a prime p . Then there exists an integer c satisfying $c \equiv 1 \pmod{\frac{m}{p}}$ and $\chi(c) \neq 1$. Such a c exists because otherwise χ is induced from a Dirichlet character modulo $\frac{m}{p}$ and thus is imprimitive. A short computation then shows

$$G(b, \chi) = \chi(c)G(b, \chi),$$

which implies $G(b, \chi) = 0$.

- (iv) Since $(m, n) = 1$, the Chinese remainder theorem implies that any a modulo mn is of the form $a = a'n + a''m$ with a' modulo m and a'' modulo n . Whence

$$(\chi\psi)(a) = \chi(a'n)\psi(a''m).$$

Using this fact, a short computation shows

$$\begin{aligned} \sum_{a \pmod{mn}} (\chi\psi)(a) e\left(\frac{ab}{mn}\right) &= \chi(n)\psi(m) \\ &\cdot \left(\sum_{a' \pmod{m}} \chi(a') e\left(\frac{ab}{m}\right) \right) \cdot \left(\sum_{a'' \pmod{n}} \psi(a'') e\left(\frac{a''b}{n}\right) \right), \end{aligned}$$

which is equivalent to the claim.

- (v) First consider the case when $\left(\frac{m}{q}, q\right) = 1$. In view of $\chi = \tilde{\chi}\chi_{\frac{m}{q},0}$, we use (iv) to obtain

$$G(\chi) = G(\chi_{\frac{m}{q},0})\tilde{\chi}\left(\frac{m}{q}\right)G(\tilde{\chi}).$$

As $G(\chi_{\frac{m}{q},0}) = r\left(1, \frac{m}{q}\right)$, we use Proposition 4.1 to compute $G(\chi_{\frac{m}{q},0}) = \mu\left(\frac{m}{q}\right)$. Whence

$$G(\chi) = \mu\left(\frac{m}{q}\right)\tilde{\chi}\left(\frac{m}{q}\right)G(\tilde{\chi}).$$

Now suppose $\left(\frac{m}{q}, q\right) > 1$. In this case the right-hand side is zero because $\tilde{\chi}\left(\frac{m}{q}\right) = 0$. So we must show $G(\chi) = 0$. Now there exists a prime p with $p \mid \frac{m}{q}$ and $p \mid q$. For any a modulo m write $a = a'\frac{m}{p} + a''$ with a' modulo p and a'' modulo $\frac{m}{p}$. Moreover, as $p \mid \frac{m}{q}$ we know $q \mid \frac{m}{p}$. These two facts and a short computation together show

$$G(\chi) = \left(\sum_{a' \pmod{p}} e\left(\frac{a'}{p}\right) \right) \left(\sum_{a'' \pmod{\frac{m}{p}}} \tilde{\chi}(a'') e\left(\frac{a''}{m}\right) \right).$$

The first sum vanishes since it is the sum of all p -th roots of unity. This proves $G(\chi) = 0$. $\square$

Via properties (i) and (ii), the evaluation of the Gauss sum is reduced to that of $G(\chi)$ at least when χ is primitive. It is also not difficult to show that $G(\chi)$ exhibits square-root cancellation. In fact, we can determine the modulus exactly.

Theorem 4.3. *Let χ be a primitive Dirichlet character of conductor q . Then*

$$|G(\chi)| = \sqrt{q}.$$

Proof. If χ is the trivial Dirichlet character the claim is obvious since the Gauss sum is 1. So assume χ is nontrivial whence $q > 1$. Consider instead $|G(\chi)|^2 = G(\chi)\overline{G(\chi)}$. Expanding the Gauss sum $\overline{G(\chi)}$ and invoking Proposition 4.2 (ii), a short computation shows

$$|G(\chi)|^2 = \sum_{a \pmod{q}} G(a, \chi) e\left(-\frac{a}{q}\right).$$

Upon expanding the Gauss sum $G(a, \chi)$, another short computation gives

$$|G(\chi)|^2 = \sum_{a' \pmod{q}} \chi(a') \left(\sum_{a \pmod{q}} e\left(\frac{a(a' - 1)}{q}\right) \right).$$

If $a' \equiv 1 \pmod{q}$ the inner sum is q and otherwise vanishes as it is the sum of all q -th roots of unity. It follows that the double sum is q whence $|G(\chi)|^2 = q$. This is equivalent to the claim. $\square$

As an almost immediate corollary, we deduce a useful expression for primitive Dirichlet characters as exponential sums.

Corollary 4.4. *Let χ be a primitive Dirichlet character of conductor q . Then for any integer b , we have*

$$G(b, \chi) = \overline{\chi}(b)G(\chi).$$

In particular,

$$\chi(b) = \frac{1}{G(\overline{\chi})} \sum_{a \pmod{q}} \overline{\chi}(a) e\left(\frac{ab}{q}\right).$$

Proof. For the first statement, the identity is obvious if χ is the trivial character as the Gauss sum is 1. So assume χ is nontrivial. If $(b, q) = 1$ then this is exactly Proposition 4.2 (ii). If $(b, q) > 1$ then the identity follows from Proposition 4.2 (iii) and that $\overline{\chi}(b) = 0$. This proves the first statement in full. For the second statement, observe that $G(\chi) \neq 0$ by Theorem 4.3. The second identity follows upon replacing χ with $\overline{\chi}$, dividing by $G(\chi)$, and expanding the Gauss sum. $\square$

For a Dirichlet character χ modulo m , we define the **epsilon factor** ε_χ by

$$\varepsilon_\chi = \frac{G(\chi)}{\sqrt{m}}.$$

When χ is primitive, the epsilon factor lies on the unit circle by Theorem 4.3. The question of the evaluation of Gauss sums boils down to determining what value the epsilon factor is. This is the real difficulty in evaluating Gauss sums. However, when χ is primitive there is a simple relationship between the epsilon factors ε_χ and $\varepsilon_{\overline{\chi}}$:

Proposition 4.5. *Let χ be a primitive Dirichlet character of conductor q . Then*

$$\varepsilon_\chi \varepsilon_{\overline{\chi}} = \chi(-1).$$

Proof. If χ is trivial this is obvious since both epsilon factors are 1. So assume χ is nontrivial. On the one hand, $\varepsilon_{\bar{\chi}}$ lies on the unit circle so that

$$\varepsilon_{\bar{\chi}}^{-1} = \frac{\overline{G(\chi)}}{\sqrt{q}}.$$

On the other hand, Proposition 4.2 (i) implies

$$\varepsilon_{\chi} = \chi(-1) \frac{\overline{G(\chi)}}{\sqrt{q}}.$$

Combining these identities gives the result. □

4.3 Quadratic Gauss Sums

Our primary aim will be to evaluate the epsilon factor of the Gauss sum for quadratic Dirichlet characters defined by Jacobi symbols. To accomplish this we will study an auxiliary exponential sum. For integers b and m with m positive, the **quadratic Gauss sum** $g(b, m)$ is defined by

$$g(b, m) = \sum_{a \pmod{m}} e\left(\frac{a^2 b}{m}\right).$$

This is a complete exponential sum. When $b \mid m$ the summands are all 1 and the quadratic Gauss sum evaluates to

$$g(b, m) = m.$$

If $b = 1$ we write $g(m) = g(1, m)$. It turns out that for square-free m the Gauss sum attached to the quadratic Dirichlet character modulo m given by the Jacobi symbol is precisely the quadratic Gauss sum. This takes some work to prove. The first step is to reduce to the case when $(b, m) = 1$.

Proposition 4.6. *Let b and m be integers with m positive. Then*

$$g(b, m) = (b, m) g\left(\frac{b}{(b, m)}, \frac{m}{(b, m)}\right).$$

Proof. Any a modulo m is of the form $a = a' \frac{m}{(b, m)} + a''$ with a' modulo (b, m) and a'' modulo $\frac{m}{(b, m)}$. A short computation shows

$$\sum_{a \pmod{m}} e\left(\frac{a^2 b}{m}\right) = (b, m) \sum_{a'' \pmod{\frac{m}{(b, m)}}} e\left(\frac{(a'')^2 b}{m}\right).$$

The remaining sum is exactly $g\left(\frac{b}{(b, m)}, \frac{m}{(b, m)}\right)$ and the desired identity follows. □

The second step is to deduce an equivalent formulation of the Gauss sum associated to quadratic Dirichlet characters given by Jacobi symbols. This will imply equivalence between the aforementioned exponential sums when the modulus is an odd prime.

Proposition 4.7. *Let b and m be integers with m positive, odd, and such that $(b, m) = 1$. Let χ_m be the quadratic Dirichlet character modulo m given by the Jacobi symbol. Then*

$$G(b, \chi_m) = \sum_{a \pmod{m}} \left(1 + \left(\frac{a}{m}\right)\right) e\left(\frac{ab}{m}\right).$$

When $m = p$ is prime,

$$G(b, \chi_p) = g(b, p).$$

Proof. The first statement is obvious when $m = 1$ since the Gauss sum is 1. So assume $m > 1$. Now write the sum as

$$\sum_{a \pmod{m}} e\left(\frac{ab}{m}\right) + \sum_{a \pmod{m}} \left(\frac{a}{m}\right) e\left(\frac{ab}{m}\right).$$

The first sum vanishes as it is the sum of all m -th roots of unity. This proves the first identity. Now let $m = p$ be an odd prime. Then

$$1 + \left(\frac{a}{p}\right) = \begin{cases} 2 & \text{if } a \text{ is a quadratic residue modulo } p, \\ 0 & \text{if } a \text{ is not quadratic residue modulo } p, \\ 1 & \text{if } p \mid a. \end{cases}$$

Moreover, when a is a quadratic residue modulo p there exists an a' modulo p with $a \equiv (a')^2 \pmod{p}$. As there are $\frac{p-1}{2}$ such residues, the first statement implies

$$G(b, \chi_p) = 1 + \sum_{\substack{a' \pmod{p} \\ a' \neq 0}} e\left(\frac{(a')^2 b}{p}\right).$$

This is exactly $g(b, p)$ which proves the second statement. □

The third step is to generalize the second statement in Proposition 4.7 for square-free m . To accomplish this we will need to develop some properties of quadratic Gauss sums.

Proposition 4.8. *Let b , m , and n be integers with m and n positive and let p be an odd prime. Then the following properties hold:*

- (i) *If $(b, p) = 1$ then $g(b, p^r) = pg(b, p^{r-2})$ provided $r \geq 2$.*
- (ii) *If $(m, n) = 1$ and $(b, mn) = 1$ then $g(b, mn) = g(bn, m)g(bm, n)$.*
- (iii) *If m is odd and $(b, m) = 1$ then $g(b, m) = \left(\frac{b}{m}\right)g(m)$ where $\left(\frac{b}{m}\right)$ is the Jacobi symbol.*

Proof. We will prove the properties separately.

(i) Every a modulo p satisfies $(a, p) = 1$ or is a multiple of p . Whence

$$g(b, p^r) = \sum_{\substack{a \pmod{p^r} \\ (a, p)=1}} e\left(\frac{a^2 b}{p^r}\right) + \sum_{a \pmod{p^{r-1}}} e\left(\frac{a^2 b}{p^{r-2}}\right),$$

since every a modulo p satisfies $(a, p) = 1$ or not. Every a modulo p^{r-1} is of the form $a = a'p^{r-2} + a''$ with a' modulo p and a'' modulo p^{r-2} . A short computation shows

$$\sum_{a \pmod{p^{r-1}}} e\left(\frac{a^2 b}{p^{r-2}}\right) = p \sum_{a'' \pmod{p}} e\left(\frac{(a'')^2 b}{p^{r-2}}\right).$$

The remaining sum is exactly $g(b, p^{r-2})$. So it suffices to show that the first sum vanishes. This sum is exactly $R(b, p^r)$ and Proposition 4.1 implies $R(b, p^r) = \mu(p^r)$ which is zero as $r \geq 2$.

(ii) Since $(m, n) = 1$, the Chinese remainder theorem implies that any a modulo mn is of the form $a = a'n + a''m$ with a' modulo m and a'' modulo n . Whence

$$\left(\sum_{a' \pmod{m}} e\left(\frac{(a')^2 bn}{m}\right) \right) \left(\sum_{a'' \pmod{n}} e\left(\frac{(a'')^2 bm}{n}\right) \right) = \sum_{a \pmod{mn}} e\left(\frac{a^2 b}{mn}\right).$$

This is equivalent to the claim.

(iii) The claim is obvious if $m = 1$ because the quadratic Gauss sum is 1. So assume $m > 1$. By multiplicativity of the Jacobi symbol and (ii), it suffices to prove the claim when $m = p^r$ is an odd prime power. The case when $r = 1$ follows from Proposition 4.7, Proposition 4.2 (ii), and that quadratic Dirichlet characters are their own conjugates. The case when $r \geq 2$ follows by induction using (i). $\square$

At last we can prove our Gauss and quadratic Gauss sums agree.

Theorem 4.9. *Suppose m is a square-free positive odd integer and let χ_m be the quadratic Dirichlet character modulo m given by the Jacobi symbol. Then for any integer b with $(b, m) = 1$, we have*

$$G(b, \chi_m) = g(b, m).$$

Proof. The claim is obvious if $m = 1$ because the Gauss and quadratic Gauss sums are both 1. So suppose $m > 1$. It suffices to assume $b = 1$ by Proposition 4.2 (ii), Proposition 4.8 (iii), and that quadratic Dirichlet characters are their own conjugates. Let $m = p_1 p_2 \cdots p_k$ be the prime decomposition of m . Repeated application of Proposition 4.2 (iv) shows

$$G(\chi) = \left(\prod_{i < j} \chi_{p_j}(p_i) \chi_{p_i}(p_j) \right) \left(\prod_i G(\chi_{p_i}) \right).$$

By Proposition 4.7, we may write

$$\left(\prod_{i < j} \chi_{p_j}(p_i) \chi_{p_i}(p_j) \right) \left(\prod_i G(\chi_{p_i}) \right) = \left(\prod_{i < j} \left(\frac{p_i}{p_j} \right) \left(\frac{p_j}{p_i} \right) \right) \left(\prod_i g(p_i) \right).$$

Repeated application of Proposition 4.8 (ii) gives

$$\left(\prod_{i < j} \left(\frac{p_i}{p_j} \right) \left(\frac{p_j}{p_i} \right) \right) \left(\prod_i g(p_i) \right) = g(m).$$

This completes the proof. $\square$

The properties in Proposition 4.8 help to reduce the evaluation of quadratic Gauss sums. Thankfully, it is possible to completely evaluate quadratic Gauss sums when $b = 1$. As with the Gauss sum, we first deduce a fact about the modulus.

Theorem 4.10. *Let m be a positive odd integer. Then*

$$|g(m)| = \sqrt{m}.$$

Proof. The claim is obvious if $m = 1$ because the quadratic Gauss sum is 1. So assume $m > 1$. By Proposition 4.8 (ii) and (iii), it suffices to prove the claim when $m = p^r$ is an odd prime power. The case when $r = 1$ follows from Theorems 4.3 and 4.9. The case when $r \geq 2$ is proved by induction and Proposition 4.8 (i). $\square$

For any integer m , we define the **epsilon factor** ε_m by

$$\varepsilon_m = \frac{g(m)}{\sqrt{m}}.$$

Theorem 4.10 says that this value lies on the unit circle when m is odd. This evaluation of these epsilon factors was completely resolved by Gauss in 1808. Modern proofs use analytic techniques by expressing $g(m)$ in a form where Poisson summation can be applied.

Theorem 4.11. *Let $m \geq 1$. Then*

$$\varepsilon_m = \begin{cases} (1+i) & \text{if } m \equiv 0 \pmod{4}, \\ 1 & \text{if } m \equiv 1 \pmod{4}, \\ 0 & \text{if } m \equiv 2 \pmod{4}, \\ i & \text{if } m \equiv 3 \pmod{4}. \end{cases}$$

Proof. We will write the quadratic Gauss in a form to which Poisson summation can be applied. For then we can evaluate the quadratic Gauss sum by means of Fourier transforms. To this end, consider the function

$$f(x) = \begin{cases} e\left(\frac{x^2}{m}\right) & \text{if } x \in [0, m], \\ 0 & \text{if } x \notin [0, m], \end{cases}$$

which is piecewise smooth and compactly supported on $\mathbb{R}$. Therefore Poisson summation applies. On the one hand, this means

$$\sum_{n \in \mathbb{Z}} f(n) = g(m).$$

On the other hand, we have

$$\sum_{n \in \mathbb{Z}} f(n) = \sum_{t \in \mathbb{Z}} \sqrt{m} e\left(-\frac{t^2 m}{4}\right) \int_{\frac{t\sqrt{m}}{2}}^{\sqrt{m} + \frac{t\sqrt{m}}{2}} e(x^2) dx.$$

As $t \equiv 0, 1 \pmod{4}$ according to if t is even or odd, the subsums according to this parity are

$$\sqrt{m} \int_{-\infty}^{\infty} e(x^2) dx \quad \text{and} \quad \sqrt{m} e\left(-\frac{m}{4}\right) \int_{-\infty}^{\infty} e(x^2) dx,$$

respectively. As the sum is ordered with respect to this parity, we have

$$\sum_{n \in \mathbb{Z}} f(n) = \sqrt{m} \left(1 + e\left(-\frac{m}{4}\right)\right) \int_{-\infty}^{\infty} e(x^2) dx.$$

Equating our two expressions gives

$$g(m) = \sqrt{m} \left(1 + e\left(-\frac{m}{4}\right)\right) \int_{-\infty}^{\infty} e(x^2) dx.$$

To compute the remaining integral we take $m = 1$. As the Gauss sum is 1 and $e\left(-\frac{1}{4}\right) = -i$, we find that

$$\int_{-\infty}^{\infty} e(x^2) dx = \frac{1}{1-i}.$$

Therefore

$$\varepsilon_m = \frac{1 + e\left(-\frac{m}{4}\right)}{1-i} = \begin{cases} (1+i) & \text{if } m \equiv 0 \pmod{4}, \\ 1 & \text{if } m \equiv 1 \pmod{4}, \\ 0 & \text{if } m \equiv 2 \pmod{4}, \\ i & \text{if } m \equiv 3 \pmod{4}, \end{cases}$$

as desired. □

As an immediate corollary, we can evaluate the epsilon factor ε_{χ_p} for the quadratic Dirichlet character χ_p modulo p given by the Jacobi symbol when p is an odd prime.

Corollary 4.12. *Let p be an odd prime and χ_p be the quadratic Dirichlet character modulo p given by the Jacobi symbol. Then*

$$\varepsilon_{\chi_p} = \begin{cases} 1 & \text{if } p \equiv 1 \pmod{4}, \\ i & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Proof. The claim follows immediately from Theorem 4.11 and Proposition 4.7. □

4.4 Character Sums and the Pólya-Vinogradov Inequality

Let χ be a non-principal Dirichlet character modulo m . For nonnegative integers M and N , the **character sum** $S_\chi(M, N)$ associated to χ is given by

$$S_\chi(M, N) = \sum_{M \leq n \leq M+N} \chi(n).$$

This is an incomplete exponential sum unless $N = m - 1$. If $M = 0$, we write $S_\chi(N) = S_\chi(0, N)$ for simplicity. An important question is how effectively can $S_\chi(M, N)$ be estimated for various Dirichlet characters χ and regimes of M and N relative to the modulus of χ . Since this is an exponential sum, we have the trivial bound

$$S_\chi(M, N) \leq N - 1.$$

On the other hand, by the Dirichlet orthogonality relations and that χ is non-principal we have

$$S_\chi(M, N) \leq m.$$

This second estimate is only an improvement upon the first in the regime $m \ll N$. Surprisingly, we can obtain square-root cancellation in this regime with very little work. We will require a small lemma.

Lemma 4.13. *Let $f(x)$ be a integrable convex function on $(a - \frac{b}{2}, a + \frac{b}{2})$ for some real a and positive b . Then*

$$f(a) \leq \frac{1}{b} \int_{a-\frac{b}{2}}^{a+\frac{b}{2}} f(x) dx.$$

Proof. By the definition of convexity, write

$$f(a) \leq \frac{1}{2}f(x) + \frac{1}{2}f(2a - x).$$

Integrate the left-hand side over $(a - \frac{b}{2}, a + \frac{b}{2})$ yields

$$\int_{a-\frac{b}{2}}^{a+\frac{b}{2}} f(a) dx = bf(a),$$

while integrating the right-hand side over $(a - \frac{b}{2}, a + \frac{b}{2})$ gives

$$\int_{a-\frac{b}{2}}^{a+\frac{b}{2}} \left(\frac{1}{2}f(x) + \frac{1}{2}f(2a - x) \right) dx = \int_{a-\frac{b}{2}}^{a+\frac{b}{2}} f(x) dx,$$

upon making the change of variables $x \mapsto 2a - x$ in the second term. Hence

$$bf(a) \leq \int_{a-\frac{b}{2}}^{a+\frac{b}{2}} f(x) dx,$$

which is equivalent to the claim. □

The aforementioned improvement is known as the **Pólya-Vinogradov inequality** as it was proved independently by Pólya and Vinogradov in 1918.

Theorem (Pólya-Vinogradov inequality). *Let χ be a non-principal Dirichlet character modulo m . Then for any nonnegative integers M and N , we have*

$$S_\chi(M, N) \ll \sqrt{m} \log(m).$$

Proof. First suppose χ is primitive of conductor $q > 1$. Using Corollary 4.4, write

$$S_\chi(M, N) = \frac{1}{G(\bar{\chi})} \sum_{a \pmod{q}} \bar{\chi}(a) \sum_{M \leq n \leq M+N} e\left(\frac{an}{q}\right).$$

The inner sum is geometric and a short computation shows

$$\sum_{M \leq n \leq M+N} e\left(\frac{an}{q}\right) = e\left(\frac{a\left(M + \frac{N-1}{2}\right)}{q}\right) \frac{\sin\left(\frac{\pi Na}{q}\right)}{\sin\left(\frac{\pi a}{q}\right)},$$

upon using the identity $\sin(x) = \frac{e^{ix} - e^{-ix}}{2i}$. This result together with Theorem 4.3 gives

$$S_\chi(M, N) \ll \frac{1}{\sqrt{q}} \sum_{1 \leq a \leq q-1} \frac{1}{\sin\left(\frac{\pi a}{q}\right)}.$$

The function $\frac{1}{\sin(\pi x)}$ is easily checked to be convex and integrable on $(0, 1)$ (from checking its second derivative is positive). Invoking Lemma 4.13 gives

$$\frac{1}{\sqrt{q}} \sum_{1 \leq a \leq q-1} \frac{1}{\sin\left(\frac{\pi a}{q}\right)} \leq \sqrt{q} \int_{\frac{1}{2q}}^{1 - \frac{1}{2q}} \frac{1}{\sin(\pi x)} dx.$$

The integrand is symmetric over $\left(\frac{1}{2q}, \frac{1}{2}\right)$ and $\left(\frac{1}{2}, 1 - \frac{1}{2q}\right)$. Whence the latter expression is equivalent to

$$2\sqrt{q} \int_{\frac{1}{2q}}^{\frac{1}{2}} \frac{1}{\sin(\pi x)} dx.$$

Now $\sin(\pi x) \geq 2x$ on the interval $[0, \frac{1}{2}]$. Applying this estimate shows

$$2\sqrt{q} \int_{\frac{1}{2q}}^{\frac{1}{2}} \frac{1}{\sin(\pi x)} dx \ll \sqrt{q} \log(q).$$

Therefore

$$S_\chi(M, N) \ll \sqrt{q} \log(q).$$

This proves the bound in the case χ is primitive. Now suppose χ is induced by the primitive character $\tilde{\chi}$ of conductor $q > 1$. Then $q \mid m$, and so we may write $m = kq$ for some positive integer k . Whence

$$S_\chi(M, N) = \sum_{\substack{M \leq n \leq M+N \\ (n, k)=1}} \tilde{\chi}(n).$$

Invoke Proposition 1.2 to remove the relatively prime condition obtaining

$$\sum_{d|k} \mu(d) \sum_{\substack{M \leq n \leq M+N \\ d|n}} \tilde{\chi}(n).$$

Now make the change of variables $n \mapsto dn$ to get

$$\sum_{d|k} \mu(d) \tilde{\chi}(d) \sum_{\lfloor \frac{M}{d} \rfloor \leq n \leq \lfloor \frac{M+N}{d} \rfloor} \tilde{\chi}(n).$$

The inner sum is $O(\sqrt{q} \log(q))$ by the primitive case and so

$$S_{\chi}(M, N) \ll \sigma_0(k) \sqrt{q} \log(q),$$

since there are $\sigma_0(k)$ divisors of k . The bound $\sigma_0(k) \ll \sqrt{k}$ is implied by Proposition 2.1. This estimate together with $\log(q) \leq \log(m)$ gives

$$S_{\chi}(M, N) \ll \sqrt{m} \log(m),$$

as claimed. □

The Pólya-Vinogradov inequality is particularly useful when $N \gg m$. The following slight improvement was made in 1977 by Montgomery and Vaughn under the Riemann hypothesis for Dirichlet L -functions:

Theorem 4.14. *Let χ be a non-principal Dirichlet character modulo m . Then for any $M \geq 0$ and $N \geq 1$,*

$$S_{\chi}(M, N) \ll \sqrt{m} \log \log(m),$$

provided the Riemann hypothesis for Dirichlet L -functions holds.

While this result is not much of an improvement from the Pólya-Vinogradov inequality, it is sharp due to a result of Paley in 1932. This means that, quite remarkably, the Pólya-Vinogradov inequality is almost optimal. Still, it is useful to have sharper estimates when $N \ll m$. In 1963, Burgess made progress in this direction by proving the following result:

Theorem 4.15. *Let χ be a non-principal Dirichlet character modulo $m > 1$, let M and N be nonnegative integers, and let r be a positive integer. Then*

$$S_{\chi}(M, N) \ll_{\varepsilon} N^{1-\frac{1}{r}} m^{\frac{r+1}{4r^2}+\varepsilon},$$

provided m is cube-free or $r = 2$.

Burgess's result can be thought of as a blend between the aforementioned trivial bound and the Pólya-Vinogradov inequality. Despite the proof being reasonably short, it requires machinery coming from finite fields.