


Data & Qualitative Data

Exploring Data | Data & Graphical Summaries

© University of Sydney DATA1001/1901

Unit Overview





Module1 Exploring Data

Design of Experiments

Where did the data come from & can we make reliable conclusions?

Data & Graphical Summaries

What type of data do we have & how can we visualise it?

Numerical Summaries

What are the main features of the data?



Data & Qualitative Data

Data Story | What causes Australian Road Fatalities?

Initial Data Analysis

Structure of the Data

Graphical Summaries

Qualitative Data

Extra Example

Summary

Data Story

What causes Australian Road Fatalities?

This is every road death since 1989

By Inga Ting , Alex Palmer , Ri Liu , Nathanael Scott and Mark Doman

Updated 26 Jan 2018, 5:15pm
Published 25 Jan 2018, 5:58am

Three men, aged 23, 29 and 30, killed while travelling in a 50km/h speed zone

Six youths killed in Tasmania when a car collides with a log truck. It is 8:20pm on a Wednesday

Road deaths are largely predictable and preventable — a fact public health experts have sought to underscore by discouraging use of the word “accident” when it comes to road crashes.

Monday, 12:20pm: A 63-year-old cyclist
in a 50 km/h speed zone is killed

NATIONAL NSW

No longer such a safe zone, especially in the afternoon

By Tim Barlass, Rachel Browne
7 October 2012 — 4:00am


f t e | A A A

CHILDREN are at risk of being hit by speeding drivers with new research showing more than 70 per cent of drivers entering school zones break the 40 km/h speed limit.

As schools return from holidays this week, a Macquarie University study shows the afternoon school pick-up is more dangerous for children than the morning drop-off.

The research showed that in the afternoon, 78.8 per cent of vehicles were exceeding the 40 km/h limit as they entered the zone, while 75 per cent were speeding as they left.

In the morning, 70 per cent of cars entered school zones above the speed limit while 47 per



Australian Road Fatality Data

- Despite preventative measures such as [compulsory seat belts \(from 1970\)](#) and [school zones \(2001\)](#), the number of road fatalities in Australia continues to rise.
 - In 2015, 1,209 died in road fatalities.
 - Why?
- We are going to investigate data from the Australian Bureau of Statistics (ABS) for January-April 2016.

```
# Read in data
data = read.csv("data/2016Fatalities.csv",header=T)
# Names of Variables
names(data)
```

```
## [1] "Crash.ID"          "State"
## [3] "Date"             "Day"
## [5] "Month"            "Year"
## [7] "Dayweek"          "Time"
## [9] "Hour"             "Min"
## [11] "Crash.Type"        "BusInvolvement"
## [13] "RigidTruck..Involvement" "Articulated.Truck..Involvement."
## [15] "SpeedLimit"        "RoadUser"
## [17] "Gender"            "Age"
```

 Find Data Dictionary



Statistical Thinking

What questions do you have?

- How many road fatalities have there been so far this year, and how does it compare to last year?
- What is the most common day and time for a crash?
- Does biological sex affect the type of road fatality?
- What is the chance that a motorcycle rider is involved in a road fatality?
- How many people wear seatbelts?


Initial Data Analysis

What is data?



Data is **information** about the set of **subjects** being studied (like road fatalities).

- Most commonly, data refers to the sample, not the population.



Different types of data

There are many different types of data, in different formats.

For example:

- survey data
- spreadsheet type data
- MRI image data




Initial Data Analysis (IDA)



Initial Data Analysis (IDA)

Initial Data Analysis is a first general look at the data, without formally answering the research questions.

- IDA helps you to see whether the data can answer your research questions.
- IDA may pose other research questions.
- IDA can
 - identify the data's main qualities;
 - suggest the population from which a sample derives.



“The purposes of IDA are to ensure that the later statistical analysis can be performed efficiently and to minimize the risk of incorrect or misleading results.”

Huebner et al. 2016

What's involved in IDA?

Initial Data Analysis commonly involves:

- data background: checking the quality and integrity of the data
- data structure: what information has been collected?
- data wrangling: scraping, cleaning, tidying, reshaping, splitting, combining
- data summaries: graphical and numerical

Here we focus on **structure & graphical summaries** for qualitative and quantitative data.

Structure of the Data


Variables

Variable

A **variable** measures or describes some attribute of the subjects.

- Data with p variables is said to have **dimension** p .

Number of variables





Statistical Thinking

How many variables does the road fatality data have?


- The road fatality data has dimension $p = 17$, as the CrashID serves as an anonymous identifier.

Size of Data

```
dim(data)
```

```
## [1] 442 18
```

Types of variables





Statistical Thinking

Classify the variable **Age** in the Road Fatality Data.

- Technically Age is a quantitative, continuous variable, but here the ages have been reported as discrete 'integer' (by rounding down to the nearest year).
- Age may also be recorded as a qualitative variable in a survey, as respondents may be more willing to give their age category. However, it is optimal to record quantitative data if possible.

Suggest a similar variable.

- Income.

Structure of Data

str(data)

```
## 'data.frame':  442 obs. of  18 variables:
## $ Crash.ID      : num  3.2e+12 3.2e+12 2.2e+12 3.2e+12 1.2e+12 ...
## $ State         : Factor w/  8 levels "ACT","NSW","NT",...: 4 4 7 4 2 8 2 7 8 5 ...
## $ Date          : Factor w/ 113 levels "1-Apr-16","1-Feb-16",...: 29 62 73 57 102 106 70 58 103 32 ...
## $ Day           : int   16 24 26 22 7 8 26 23 7 17 ...
## $ Month         : Factor w/  4 levels "April","February",...: 2 1 4 4 1 1 1 1 2 1 ...
## $ Year          : int   2016 2016 2016 2016 2016 2016 2016 2016 2016 2016 ...
## $ Dayweek       : Factor w/  7 levels "Friday","Monday",...: 6 4 3 6 5 1 6 3 4 4 ...
## $ Time          : Factor w/ 225 levels "0:00","0:12",...: 53 187 90 73 219 106 108 67 44 46 ...
## $ Hour          : int    13  5 16 15  9 17 17 14 12 12 ...
## $ Min           : int     0  0  0  0 20 10 19 32 26 35 ...
## $ Crash.Type    : Factor w/  3 levels "Multiple vehicle",...: 1 3 2 2 2 1 2 3 1 1 ...
## $ BusInvolvement : Factor w/  2 levels "No","Yes": 1 1 1 1 1 1 1 1 1 1 ...
## $ RigidTruck..Involvement : Factor w/  2 levels "No","Yes": 1 1 1 1 1 1 1 1 1 2 ...
## $ Articulated.Truck..Involvement.: Factor w/  2 levels "No","Yes": 1 1 1 1 1 1 1 1 1 1 ...
## $ SpeedLimit    : int   100 70 100 50 50 50 50 -9 60 110 ...
## $ RoadUser      : Factor w/  6 levels "Bicyclist (includes pillion passengers)",...: 2 2 6 6 6 6 6 5 5 5 ...
## $ Gender        : Factor w/  2 levels "Female","Male": 1 2 1 2 2 2 2 1 1 1 ...
## $ Age           : int    -9 -9  0  1  1  1  2  4  5  5 ...
```


Graphical Summaries

Choosing a graphical summary

The aim of a graphical summary is to best highlight features of this data.

- To some extent we use trial and error.
- While the pie chart may be popular, it is usually not informative.
- The relatively new 'Shiny Apps' present data in accessible ways. [See examples.](#)

Twitter debate about pie charts




Max Roser
@MaxCRoser

Follow



Why you should not use pie charts.



5:22 PM - 26 Apr 2017

6,216 Retweets 8,851 Likes






William Lundstedt @healthire_ · 28 Apr 2017

Replying to @MaxCRoser

Why you should not make blanket statements.



4 8 83



Max Roser @MaxCRoser · 28 Apr 2017

Agreed. There are cases, like datasets with two categories as in your example, where they make sense. Just shared a more complete article.


Shiny Apps

Visualizing SNAP

Enter your household monthly SNAP allotment (in \$) to receive a recommended weekly expenditure on groceries:

125

We recommend spending about \$ 78 per week on food. Enter the range below that you believe reflects this budget:




What would you like your daily caloric intake to be?:

I am primarily trying to minimize:

Cost

Hover over points to find out how many servings of each ingredient you need to prepare dishes each week. Toggle the bars on the left to adjust to your dietary needs.

What Foods Should I Buy?



Qualitative Data

Simple bar plot


The bar plot (or bar chart or bar graph) is a simple summary of qualitative data.

```
# Select the DayWeek variable from the whole data frame
Dayweek = data$Dayweek


# Produce a frequency table of fatalities per day of the week
table(Dayweek)
```

```
## Dayweek
##   Friday   Monday  Saturday   Sunday  Thursday  Tuesday  Wednesday
##      68      50      85      56      58      67      58
```

```
# Produce a bar chart  
barplot(table(Dayweek))
```




```
# A simpler version  
plot(table(Dayweek))
```





Statistical Thinking

What was the most common day of road fatality?

- Saturday

Why might that be the case?

- More volume of cars on the road, or people driving faster?

What data would you need to check your hypotheses?

- Data on volume and speed of cars on the road each day.

Double bar plot

Things get more interesting when we consider 2 qualitative variables.

```
# Select DayWeek and Gender (Biological Sex) variables
Dayweek = data$Dayweek
Gender = data$Gender


# Produce a double frequency table (contingency table)
data1 = table(Gender, Dayweek)
data1
```

```
##          Dayweek
## Gender  Friday Monday Saturday Sunday Thursday Tuesday Wednesday
##  Female    15    17      25    13      16      17      13
##   Male     53    33     60    43     42     50     45
```

Note: Here we have called the variables by the names in the data set. Here “Gender” refers to biological sex, as it was historically recorded in this dataset. [Read more](#).


Stacked Bar Plot

```
barplot(data1, main="Fatalities by Day of the Week and Biological Sex",  
        xlab="Day of the week", col=c("lightblue","lightgreen"),  
        legend = rownames(data1))
```



Side-by-Side Bar Plot

```
barplot(data1, main="Fatalities by Day of the Week and Biological Sex",  
        xlab="Day of the week", col=c("lightblue","lightgreen"),  
        legend = rownames(data1), beside=TRUE)
```





Statistical Thinking


Do you have a preference for stacked or side-by-side? Why?

Are these plots telling us anything useful? How could they be misread?

- There seems to be a similar proportion of fatalities across each day, for biological sex.
- We could posit that men are more likely to be involved in fatal accidents than women. However, perhaps there are more men on the road than women. More data is needed.

Graphical summaries of big data

- **Big data** is the massive amounts of data being collected in fields such as genomics, astrophysics, marketing and sociology.
- Big data is commonly **high dimensional**, which means that there are more variables p than subjects n .
 - For example, genomics data can have 3 billion variables, as a person's DNA sequence is 3 billion basepairs long.
- Big data can be described by many “V”s: high volume, high velocity, high variety, high variability, low veracity/validity, high vulnerability, high volatility and high value.
- Big data requires more complex visualisations.



Sources: McKinsey Global Institute, Twitter, Cisco, Gartner, EMC, SAS, IBM, MEPTec, QAS

IBM

Extra Example

Investigating Speed Limit


Speed Limit could be considered as a discrete, quantitative variable, but it is best classified as an (ordinal) qualitative variable. Why?

```
SpeedLimit = data$SpeedLimit  
table(SpeedLimit)
```

```
## SpeedLimit  
##  -9  40  50  60  70  80  90 100 110 130 888  
##  28   4  53  70  21  53  10 128  71   3   1
```

Simple Bar Plot

```
barplot(table(SpeedLimit))
```





Statistical Thinking

What is curious about this data? Why?


- -9 indicates a missing value. Why speed of 888? We could 'clean' the data.

What was the most common speed at which a road fatality occurred? How might this affect public policy?

- Notice how many fatalities happen at high speeds (100km/hr+).
- This might lead to change in speed limits or investigation of road conditions for high speeds.
- Can we assume that a vehicle travelling in that speed zone was travelling at that speed limit?

Double Bar Plot

```
#Gender = data$Gender  
data2 = table(Gender,SpeedLimit)  
barplot(data2, main="Fatalities by Speed Limit and Biological Sex",  
  xlab="Speed Limit", col=c("lightblue","lightgreen"),  
  legend = rownames(data1), beside=TRUE)
```





Statistical Thinking

Are there any interesting patterns?

Summary

The type of variables determines what type of graphical summary.

1 Qual	2 Qual	1 Quant	2 Quant	1 Quant + 1 Qual
Simple Bar Plot	Double Bar Plot	Histogram	Scatter Plot	Comparative Box Plot
		Box Plot		

Key Words

data, subjects, population, sample, Initial Data Analysis, data background, data structure, data wrangling, data summaries, variable, dimension, multivariate, bivariate, univariate, qualitative/categorical, quantitative/numerical, ordinal, nominal, discrete, continuous, binary, bar plot (chart)

Further Thinking

Big data in the humanitarian sector

Data is the new seatbelt

Data is the new seat belt | Yiem Sunbhanich | TEDxColumbus

