

Module-wise Data

Module: Post Abortion Care

Icon: /icon/module/post_abortion_care

Module Key: post-abortion-care_1487676484604

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Action Cards:

- definitions_1486843717068
- emergency-management_1486843944185
- following-management_1486845338095
- emergency-referral_1490716065855

Procedures:

- cleaning-manual-vacuum-aspirator_1487245553833
- urine-catheter_1487245590353
- intravenous-access-equipment_1487245827833

Videos:

- /India english/Post abortion care/intro
- /India english/Post abortion care/how_to_diagnose
- /India english/Post abortion care/how_to_manage
- /India english/Post abortion care/evacuation_of_the_uterus
- /India english/Post abortion care/manual_vacuum_aspiration

Drugs:

- ampicillin_1486975748548
- diazepam_1487102080825
- gentamicin_1487679704293
- ibuprofen_1490364323757
- metronidazole_1487112262252

- morphine_1487151518002

Key Learning Points:

- test-version_1498639439941
- septic-abortion-_1504773307717
- complication-to-unsafe-abortion_1504773499553
- urine-test_1504773825842
- complication-to-abortion_1504774087137
- fever-and-septic-abortion_1504774293226
- what-to-measure-_1504774413976
- signs-of-septic-abortion_1504887365737
- suspect-lower-abdominal-tenderness_1504887830839
- septic-abortion-can-be-fatal_1504888032303
- most-important-action-at-septic-abortion_1504889099044
- action-at-septic-abortion_1504889244498
- ab-and-evacuation-of-the-uterus_1504889778909
- which-ab-to-give_1504889919542
- urgent-management-_1504890132226
- correct-order-of-actions_1504890487022
- peritonitis_1504890766676
- evacuation-of-the-uterus-_1504891033708
- manual-vacuum-aspiration_1504891220598
- uterus-size-in-week-12_1504891361270
- preparation-for-mva_1504891579618
- analgesic-for-mva_1504891713356
- valves-closed_1504891839206
- vacuum-to-mva-syringe_1504891916767
- tenaculum_1504892022670
- dilate-the-cervix_1504892212948
- no-touch-technique_1504892332161
- size-of-cannula-for-mva_1504893593707
- insert-of-the-cannula_1504893750649
- attachment-of-the-syringe_1504893864011
- rotate-and-move-cannula-back-and-forth_1504893996349

- end-of-suctioning_1504894096498
- use-the-cannula-again_1504894223366
- ab-after-mva_1504894299772
- counselling_1504894423681
- advise-before-dismissal_1504894568322

Module: Infection Prevention

Icon: /icon/module/infection_prevention

Module Key: infection-prevention_1487676001934

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Procedures:

- mixing-alcohol-based-handrub_1487245431108
- mixing-chlorine-solution_1487245480781

Videos:

- /India english/Infection prevention/intro
- /India english/Infection prevention/hand_hygiene
- /India english/Infection prevention/procedure_for_handwash
- /India english/Infection prevention/procedure_for_handrub
- /India english/Infection prevention/
personal_protective_equipment
- /India english/Infection prevention/cleaning_decontamination
- /India english/Infection prevention/storage
- /India english/Infection prevention/housekeeping
- /India english/Infection prevention/hiv_hepatitis_b_prevention

Drugs:

- post-exposure-prophylaxis---pep_1490708337530

Key Learning Points:

- prevent-infections_1497356424386
- hand-hygiene-wash-and-alcohol_1497356912190
- use-alcohol-based-handrub_1497357082394
- soap-for-handwashing_1497357310306
- personal-protective-equipment_1497357545527
- gloves-for-different-patients_1497357682494
- soap-solution_1497357821040
- mix-chlorine-solution_1497357928324
- package-of-sterile-items_1497358040371
- storage-of-sterile-items_1497358161852
- cleaning-and-delivery-bed_1497358695244
- recommended-vaccination_1497358858644
- test_1497612420029
- test-1_1497612450477
- test-2_1497612458572
- test-3_1497612477070
- testing-order_1497875988625
- infections-and-maternal-death_1498119404097
- infection-and-neonatal-death_1498119889065
- infection-and-germs_1498122575522
- healthcare-workers-hands_1498122780668
- alcohol-most-effective_1498123474560
- no-jewellery-on-hands_1498129645819
- towel_1498130033393
- when-hand-hygiene_1498130470830
- soak-items-in-soap-solution_1498130755471
- order-of-cleaning-and-storage_1498131043643
- disinfection-solution_1498131845883
- spread-of-hiv-and-hep-b_1498132068449
- when-hand-hygiene_1498206602602
- wash-hands_1498466635307
- procedure-for-handwashing_1498466869003
- amount-of-handrub_1498468033321
- product-in-alcohol-based-handrub_1498468332125
- label-of-alcohol-based-handrub_1498468847400

- removing-gloves_1498469162217
- gloves-for-birth_1498469312963
- cleaning-and-decontamination_1498469480918
- high-risk-area_1498469624522
- transmission-most-likely_1498469768452
- action-if-exposed_1498469888180
- soap-wet-container_1504268064997
- air-dry-hands_1504268482403
- product-in-alcohol-based-handrub-2_1511791111025

Module: Hypertension

Icon: /icon/module/hypertension

Module Key: hypertension_1487677992452

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Action Cards:

- definitions_1486814136250
- management-hypertension_1486815930026
- management-eclampsia_1486815158568
- emergency-referral_1490716065855

Procedures:

- urine-catheter_1487245590353
- intravenous-access-equipment_1487245827833

Videos:

- /India english/Hypertension/intro
- /India english/Hypertension/definitions
- /India english/Hypertension/causes
- /India english/Hypertension/how_to_diagnose
- /India english/Hypertension/management_of_hypertension
- /India english/Hypertension/severe_preeclampsia_eclampsia

Drugs:

- calcium-gluconate_1490273124941
- hydralazine_1487109388353
- labetalol_1490625535572
- magnesium-sulphate_1487111473832
- methyldopa_1487111904464
- nifedipine_1487151847555

Key Learning Points:

- severe-preeclampsia_1498640805283
- gestational-hypertension_1498725687229
- bp-gestational-hypertension_1498725828989
- preeclampsia_1498726056310
- severely-elevated-blood-pressure_1498726396275
- severe-headache_1498726747222
- visual-disturbances_1498727185096
- upper-abdominal-pain_1498727830882
- respiratory-distress_1498728011726
- oliguria_1498728193444
- eclampsia_1498728592426
- eclampsia-normal-blood-pressure_1498728833764
- differential-diagnosis_1498728983703
- causes-of-pre-eclampsia_1498729147494
- anc-blood-pressure_1498729463504
- confirm-bp_1498732684396
- confirmed-elevated-bp_1498733012381
- eclampsia-maternal-and-perinatal-death_1498733194577
- keep-systolic-bp-below-160_1498733307152
- keep-diastolic-bp-above-80_1498733486432
- labetalol_1498733591134
- methyldopa_1498733669777
- iv-with-bp-above-160-_1498733752579
- hydralazine_1498733943552
- eclampsia-seizures_1498734086062

- magnesium-sulphate-24-hours_1498734258534
- severe-pre-eclampsia-or-eclampsia_1498734362115
- hypertension-and-urine-catheter_1498735254202
- eclampsia---monitor-12-hour_1498735362891
- give-magnesium-sulphate_1498735437004
- mix-magnesium-sulphate_1498735538053
- toxicity_1498735622169
- antidote-to-magnesium-sulphate_1498735735523
- maintain-dose-magnesium-sulphate_1498735869673
- refer-and-eclampsia_1498735994297
- lung-maturity_1498736081492
- proteinuria_1505054086730
- cure-of-pre-eclampsia_1505055063577
- maternal-death-and-pre-eclampsia_1505055560037
- action-eclampsia_1505059069497
- fluid-overload_1505059297071
- calcium-gluconate_1505059639547

Module: Normal Labour and Birth

Icon: /icon/module/infection_prevention

Module Key: normal-labour-and-birth_1592462784361

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Module: Active Management of Third Stage Labour

Icon: /icon/module/active_management_of_third_stage_labour

Module Key: active-management-of-third-stage-labour_1487678773861

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Action Cards:

- emergency-referral_1490716065855

Videos:

- /India english/Active management of third stage labour/intro
- /India english/Active management of third stage labour/
give_a_uterotonic_drug
- /India english/Active management of third stage labour/
assess_the_baby
- /India english/Active management of third stage labour/
clamp_and_cut_the_umbilical_cord
- /India english/Active management of third stage labour/
controlled_cord_traction
- /India english/Active management of third stage labour/
examine_the_placenta
- /India english/Active management of third stage labour/
monitoring
- /India english/Active management of third stage labour/
early_breastfeeding

Drugs:

- antibiotic-eye-drops_1490261954893
- misoprostol_1487112562173
- oxytocin_1487156988152
- vitamin-k_1487160613336

Key Learning Points:

- test-1_1497203846089
- active-management_1499780658036
- uterotonic-drug_1499780861769
- oxytocin_1499780939312
- exclude-second-baby_1499781002066
- dose-of-oxytocin_1499781138851
- when-to-give-oxytocin_1499781203804
- storage-of-oxytocin_1499781280757
- misoprostol_1499781426287
- administration-of-misoprostol_1499781506837

- when-to-assess-the-newborn_1499781574406
- thick-meconium_1499782019890
- suction-thick-meconium_1499782064819
- dry-of-the-baby_1499782139883
- wipe-of-baby_1499782205916
- warm-baby_1499782287277
- newborn-not-breathing-well_1499782350880
- stable-newborn_1499782411878
- clamp-umbilical-cord_1499782532294
- cord-traction_1499782592747
- counter-pressure-pubic-bone_1499782667213
- rounded-uterus_1499782790928
- push-for-the-placenta_1499782842943
- check-uterus_1499782915737
- measure-blood-loss_1499783000460
- examine-the-placenta_1499783053169
- maternal-surface-of-placenta_1499783139366
- no-unattended-mother-and-newborn_1499783196382
- skin-to-skin-contact_1499783293686
- check-mother-after-birth_1499783355207
- how-often-check-mother_1499783434630
- how-often-check-newborn_1499783488277
- check-the-newborn-after-birth_1499783546639
- cord-monitor_1499783616183
- eye-drop-and-vit-k_1499783708999
- early-breastfeeding-newborn_1499783782397
- early-breastfeeding-mother_1499783921892
- active-management_1504529794145
- delivery-of-membranes_1504531639638
- estimation-of-blood-loss_1504531689289

Module: Prolonged Labour

Icon: /icon/module/prolonged_labour

Module Key: prolonged-labour_1487678855904

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Action Cards:

- indications_1486847771915
- prerequisites_1486847909849
- procedures-by-also_1486848037447
- emergency-referral_1490716065855

Procedures:

- urine-catheter_1487245590353
- foetal-heart-rate_1487245616159
- vacuum-extractor_1487245649303
- intravenous-access-equipment_1487245827833

Videos:

- /India english/Prolonged labour/intro
- /India english/Prolonged labour/causes
- /India english/Prolonged labour/1st_stage_latent_phase
- /India english/Prolonged labour/1st_stage_active_phase
- /India english/Prolonged labour/2nd_stage_prolonged
- /India english/Prolonged labour/2nd_stage_vacuum_extraction

Drugs:

- ampicillin_1486975748548
- diazepam_1487102080825
- ergometrine_1487103846814
- iron_1487110835869
- misoprostol_1487112562173
- morphine_1487151518002
- oxytocin_1487156988152

Key Learning Points:

- complication-prolonged-mother_1500303341759
- complication-prolonged-infant_1500303472241
- meaning-of-prolonged_1500303544246
- four-ps_1500303686918
- power_1500303769050
- power-seconds_1500303835797
- pelvis-passage_1500303897179
- small-pelvis_1500303985877
- small-pelvis-refer_1500304101224
- passenger_1500304149296
- psyche_1500304254893
- normal-progression-first-stage_1500304315481
- false-labour_1500304449630
- false-labour-and-urinary-tract-infections_1500304515323
- start-of-active-phase_1500304605377
- start-partograph_1500304739350
- vaginal-examination-interval_1500304824357
- when-prolonged-first-labour_1500304876769
- action-for-prolonged-first-labour_1500304936911
- reassess-in-two-hours_1500305041698
- cervix-fully-dilated_1500305095706
- push_1500305543117
- push-for-60-min_1500305592951
- vacuum-after-60-min_1500305658715
- indication-for-vacuum-extraction-_1500305701962
- assess-of-fhr_1500305759343
- critical-fhr_1500305923791
- condition-for-vacuum_1500305971761
- ga-less-37-at-a-hospital_1500306017747
- risk-of-vacuum-extraction_1500306057945
- preparation-of-vacuum_1500306101811
- address-the-patient_1500306177213
- flexion-point_1500306221884
- steady-traction_1500306294118

- give-up-vacuum-delivery_1500306346908
- incision-rarely_1500306426261
- performing-vacuum_1500306468033
- vacuum-extraction-fails_1500306519814
- cephalic-occiput-posterior_1504777759514
- more-comfortable_1504777788729
- prolonged-first-stage-latent-phase_1504777971422
- contractions-causes-if-not-prolonged-latent-phase_1504777990862
- prolonged-first-stage-active-phase_1504778019268
- progress-of-labour_1504778029127
- partograph-with-dilatation-between-alert-and-action-line-_1504778050950
- cervical-dilatation_1504778060957
- immediate-delivery_1504778233462
- head-above-the-pubic-bone_1504778275952
- posterior-fontanelle-where_1504778354078
- posterior-fontanelle_1504778363034
- placing-the-cup_1504778371941
- check-for-maternal-tissue_1504778388031
- flexion-point_1504778425046
- intact-perineum_1504778470373
- prolonged-second-stage_1504778541913

Module: Post Partum Hemorrhage

Icon: /icon/module/post_partum_hemorrhage

Module Key: post-partum-hemorrhage_1487678905876

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Action Cards:

- 1-uterus_1486846813345
- 2-legs_1486846790935
- 3-arms_1486846769297
- emergency-referral_1490716065855

Procedures:

- urine-catheter_1487245590353
- intravenous-access-equipment_1487245827833
- condom-balloon-tamponade_1490714664807

Videos:

- /India english/Post partum hemorrhage/intro
- /India english/Post partum hemorrhage/uterus_massage
- /India english/Post partum hemorrhage/empty_the_bladder
- /India english/Post partum hemorrhage/bimanual_compression
- /India english/Post partum hemorrhage/resuscitation
- /India english/Post partum hemorrhage/give_uterotonic_drugs
- /India english/Post partum hemorrhage/trauma
- /India english/Post partum hemorrhage/tissue
- /India english/Post partum hemorrhage/monitoring_after_bleeding

Drugs:

- oxytocin_1487156988152
- ergometrine_1487103846814
- misoprostol_1487112562173
- morphine_1487151518002
- diazepam_1487102080825
- ampicillin_1486975748548
- iron_1487110835869

Key Learning Points:

- maternal-death-_1504532310198
- loss-of-tone_1504532426966
- call-for-help-then-stop-the-bleeding-_1504532471069
- symptoms-for-pph_1504532547937
- occult-bleeding_1504532588274
- moderate-bleeding---characteristics-_1504532614848

- moderate-bleeding-actions-act-differently-depending-on-the-character-of-the-bleeding_1504532653096
- uterine-massage-_1504532689921
- empty-the-bladder-_1504532728261
- vital-signs_1504532772329
- strong-bleeding--characteristics_1504532942008
- bimanual-compression_1504532979962
- bimanual-compression-procedure_1504533043410
- bimanual-compression-type-of-gloves-_1504533076462
- pph-drugs_1504533111438
- misoprostol-_1504533151987
- ergometrine-_1504533241147
- prioritize-manual-maneuvers_1504533285341
- bleeding-from-a-vaginal-tear-_1504533312898
- always-do-a-rectal-exploration-_1504533348068
- placental-tissue-missing-_1504533398776
- treatment-if-there-is-placental-tissue-missing-_1504533431739
- removal-tissue-1_1504533465688
- removal-tissue-2_1504533503753
- when-to-measure-hemoglobin-_1504533536483
- observation-after-treatment-pph-_1504533565253
- cbt-is-indicated-_1504533594043
- the-causes-of-vaginal-bleeding-_1504533620719
- positioning-pph_1504533661330
- positioning---vital-organs-_1504533702197
- actions-helper-at-the-head_1504533732416
- actions-helper-at-the-arm_1504533786043
- referral-pph_1504533836750
- iv-fluids-pph_1504533868169
- oxytocin-storage_1504533913864
- cardiovascular-symptoms-pph_1504533944971
- blood-inside-uterus_1504534828505

Module: Manual Removal of Placenta

Icon: /icon/module/manual_removal_of_placenta

Module Key: manual-removal-of-placenta_1487678949720

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Action Cards:

- emergency-referral_1490716065855

Procedures:

- urine-catheter_1487245590353
- intravenous-access-equipment_1487245827833
- condom-balloon-tamponade_1490714664807

Videos:

- /India english/Manual removal of placenta/intro
- /India english/Manual removal of placenta/aorta_compression
- /India english/Manual removal of placenta/resuscitation
- /India english/Manual removal of placenta/drugs_before_removal
- /India english/Manual removal of placenta/gloves
- /India english/Manual removal of placenta/remove_the_placenta
- /India english/Manual removal of placenta/drugs_after_removal
- /India english/Manual removal of placenta/monitoring

Drugs:

- ampicillin_1486975748548
- diazepam_1487102080825
- ergometrine_1487103846814
- iron_1487110835869
- morphine_1487151518002
- misoprostol_1487112562173
- oxytocin_1487156988152

Key Learning Points:

- time-before-placenta-be-born_1500550057787
- remove-manually_1500550146760
- act-fast_1500550214224
- life-threatening_1500550548020
- overall-action_1500550609958
- when-aorta-compression_1500550713329
- mother-in-shock_1500550773682
- how-aorta-compression_1500550835318
- adequate-aorta-compression_1500550901996
- maintain-compression_1500550952103
- woman-lie-down_1500551007297
- vital-signs_1500551043524
- iv-fluid_1500551093153
- fluid-gives-fast-_1500551155005
- blood-transfusion_1500551204516
- pain-relief_1500551251327
- pain-relief-and-aorta-compression_1500551299297
- gloves-and-manual-removal-of-placenta_1500551362316
- womans-consent_1500551424020
- first-hand_1500551465462
- second-hand_1500551522653
- right-order-for-removal-of-placenta_1500551561159
- withdrawing-and-second-hand_1500551610790
- fragile-uterine-wall_1500551652340
- placenta-detached_1500551743553
- palpate-the-uterine-cavity_1500551802100
- the-overall-action-to-control-pph_1500551855322
- condom-tamponade_1500551920106
- bolus-drug-after-removal-of-placenta_1500551982282
- substituted-drug-for-oxytocin_1500552127586
- oxytocin-drip-after-removal-of-placenta_1500552209102
- how-to-mix-oxytocin-drip_1500552277806
- massage-the-fundus_1500552326288
- day-after-recommendations_1500552382430

- refer-manual-removal_1500552464602
- ab-to-newborn_1500553942631
- manual-removal-bemonc_1504534513078
- when-manual-removal_1504534643086
- heavy-bleeding_1504534768047
- excessive-bleeding_1504535064090
- stop-bleeding-retained-placenta_1504535188380
- resuscitation_1504535401875
- stop-compression-at-resuscitation_1504682726957
- how-to-manually-remove-the-placenta_1504686597840
- aorta-compression-and-manual-removal_1504686794869
- repeat-intrauterine-palpation_1504687023070
- manual-compression-ended-aorta-compression_1504687099431
- invasive_1504687163659
- after-removal-of-placenta_1504687463704
- action-right-after-removal-of-placenta_1504687629838

Module: Maternal Sepsis

Icon: /icon/module/maternal_sepsis

Module Key: maternal-sepsis_1487679043550

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Action Cards:

- definitions_1486838289048
- emergency-management_1486839145753
- following-management_1486840964786
- identify-cause_1486841405518
- emergency-referral_1490716065855

Procedures:

- urine-catheter_1487245590353
- intravenous-access-equipment_1487245827833

Videos:

- /India english/Maternal sepsis/intro
- /India english/Maternal sepsis/how_to_diagnose_definitions
- /India english/Maternal sepsis/how_to_manage
- /India english/Maternal sepsis/following_management
- /India english/Maternal sepsis/origin_of_infection
- /India english/Maternal sepsis/hiv

Drugs:

- ampicillin_1486975748548
- artesunate_1487078838130
- betamethasone_1487080183154
- chloramphenicol_1487097224306
- dexamethasone_1487101787771
- dicloxacillin_1487103485964
- gentamicin_1487679704293
- metronidazole_1487112262252
- penicillin-g-benzylpenicillin_1487157542019
- quinine_1487158788622

Key Learning Points:

- meaning-of-sepsis_1504944629301
- diagnose-sepsis_1504944721115
- septic-shock_1504944788616
- vital-signs_1504944907638
- importance-of-fluid-management_1504944997773
- assess-of-fluid-balance_1504945090408
- fluid-balance-be-calculated_1504945280195
- fluid-balance-be_1504945346611
- in-case-of-a-septic-shock-it-should-be-calculated-each-hour_1504945415855
- complications-to-sepsis_1504945501258
- initial-management_1504945569353

- circulation-_1504945650697
- drugs_1504945729088
- ampicillin-sepsis_1504945793742
- gentamicin_1504945982881
- metronidazole_1504946039271
- ceftriaxone_1504946097782
- tetracycline_1504946176147
- chloramphenicol_1504946438452
- malaria_1504946617389
- malaria-drugs_1504946711304
- artesunate_1504946886646
- artesunate_1504946909791
- quinine_1504947056059
- dosage-artesunate_1504947115207
- dosage-quinine_1504947179913
- the-6-ws_1504947235003
- the-6-ws-continue_1504947447368
- chorioamnionitis_1504947765349
- endometritis_1504947835819
- treatment-for-chorioamnionitis_1504947946364
- treatment-for-endometritis_1504948024311
- endometritis-and-no-effect-of-ab_1504948113282
- mastitis_1504948177504
- breast-abscess_1504948229532
- wound-infections_1504948296628
- other-causes-of-infection_1504948359563

Module: Neonatal Resuscitation

Icon: /icon/module/neonatal_resuscitation

Module Key: neonatal-resuscitation_1487679104279

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Action Cards:

- neonatal-resuccitation_1486842879218

- emergency-referral_1490716065855

Videos:

- /India english/Neonatal resuscitation/intro
- /India english/Neonatal resuscitation/assess_the_baby
- /India english/Neonatal resuscitation/airway
- /India english/Neonatal resuscitation/breathing
- /India english/Neonatal resuscitation/
when_to_stop_resuscitation
- /India english/Neonatal resuscitation/monitoring

Drugs:

- ampicillin_1486975748548
- ceftriaxone_1487080443380
- gentamicin_1487679704293
- phenobarbital_1487158287290

Key Learning Points:

- adequate-breathing_1504254228875
- golden-minute_1504254468378
- assess-the-newborn-after-birth_1504256308739
- skin-to-the-mother_1504256398872
- first-step_1504256461191
- meconium-suction-_1504256548380
- correct-order-suction_1504256615224
- stimulation-the-newborn_1504256686246
- newborn-is-not-breathing-well_1504256807203
- when-to-resuscitate_1504256982883
- airway-call-for-help_1504257037350
- airway-position-of-head_1504257274893
- airway-position-of-mask_1504258127166
- airway-effective-ventilation_1504258182479

- number-of-inflation-breaths_1504258233083
- jaw-thrust_1504258315698
- successful-inflation-breaths_1504258364953
- if-no-chest-movements_1504258411252
- inflation-breaths_1504258492447
- ventilation-breath_1504258548225
- not-breathing-well_1504258620844
- continue-ventilation-breaths_1504258677078
- most-important-ventilation-breaths_1504258732578
- rhythm-ventilation-breaths_1504258780270
- good-ventilation_1504258849221
- breathing---assessment_1504258930435
- breathing---assessment_1504258984771
- breathing-assessment---continued_1504259182601
- action-if-breathing-well_1504259239916
- give-oxygen_1504259318506
- when-to-stop_1504259374882
- resuscitation-stop-next_1504259443850
- skin-to-skin-and-breastfeeding_1504259573821
- monitoring_1504259668644
- low-birth-weight-and-resuscitation_1504259731434
- challenge-low-birth-weight_1504259788567
- successful-inflation-breaths_1505392891735

Module: Newborn Management

Icon: /icon/module/newborn_management

Module Key: newborn-management_1487679126990

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Action Cards:

- prevention_1486813119229
- routine-care_1486842306870
- danger-signs_1486807603214
- management-of-danger-signs_1486808272156

- pediatric-drugs-and-fluids_1486735917639
- emergency-referral_1490716065855

Procedures:

- intravenous-access-equipment_1487245827833
- nasogastric-tube_1487245852188

Videos:

- /India english/Newborn management/intro
- /India english/Newborn management/prevention
- /India english/Newborn management/
initial_care_of_the_newborn
- /India english/Newborn management/danger_signs
- /India english/Newborn management/
management_of_danger_signs

Drugs:

- ampicillin_1486975748548
- antibiotic-eye-drops_1490261954893
- betamethasone_1487080183154
- ceftriaxone_1487080443380
- dexamethasone_1487101787771
- gentamicin_1487679704293
- oxytocin_1487156988152
- phenobarbital_1487158287290
- vitamin-k_1487160613336

Key Learning Points:

- common-causes-of-newborn-death_1500553429571
- death-in-first-week_1500553499970
- avoidable-death_1500553555177

- prevent-respiratory-distress-preterm_1500553601195
- risk-factors-infection-newborn_1500553681339
- prevent-newborn-infections_1500553770880
- risk-from-resuscitation_1500554103700
- action-after-resuscitation_1500554254443
- early-feeding_1500554299315
- initial-care-of-the-newborn_1500554337833
- not-crying-or-breathing-well_1500554397207
- initial-care-of-the-newborn-_1500554450890
- initial-care-of-lbw_1500554526669
- lbw-unable-to-suck-adequately_1500554570547
- signs-of-illness_1500554688877
- management-of-danger-signs_1500554757277
- danger-signs---airway_1500554845568
- danger-signs---breathing_1500554890466
- grunting_1500554955221
- apnoea_1500554996619
- shock_1500555055530
- disability_1500555142461
- danger-signs---exposure_1500555231782
- jaundice_1500555273140
- management---case_1500555311125
- obstructed-airways_1500555401801
- management---breathing_1500555450027
- management---circulation_1500555489191
- management-drowsy-and-unconscious_1500555531251
- convulsions_1500555605930
- management-of-danger-signs_1500555700421
- management---jaundice_1500555760414
- management---kangaroo-motherhood-care-_1500555801259
- management---infections_1500555888659
- no-condition-improved_1500555927749
- risk-of-low-blood-sugar_1500555965754
- newborn-cannot-suck_1500556011761
- rural-facility-newborn-danger-signs_1500556085207

- kangaroo-care_1504774517466

Module: Low Birth Weight

Icon: /key learning points/Neonatal resuscitation/
low_birth_weight_preterm

Module Key: low-birth-weight_1539679206162

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Module: Female Genital Mutilation

Icon: /icon/module/female_genital_mutilation

Module Key: female-genital-mutilation_1494414132178

lanId: ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Key Learning Points:

- fgm-is-an-unhealthy-practice_1499073619747
- fgm-is-a-dangerous-practice_1499073729938
- fgm-is-forbidden_1499073844486
- four-types-of-fgm_1499073935484
- mix-of-the-types_1499073988266
- type-1_1499074047139
- type-2_1499074470165
- type-3_1499074607765
- type-4_1499074689724
- type-2-becomes-type-3_1499075316336
- immediate-health-risk_1499075598511
- fgm-and-infertility_1499078717543
- long-term-consequences_1499078794803
- fgm-antenatal_1499078908785
- birth-complications_1499079019921
- why-birth-complications-_1499079100900
- urethra_1499079409674
- deinfibulation-and-urethra_1499082946166
- steps-for-deinfibulation_1499083113362
- interrupted-sutures_1499083209086

- analgesia-after-deinfibulation_1499083297174
- hygiene-for-deinfibulation_1499083352056
- antenatal-deinfibulation_1499083466972
- when-antenatal-deinfibulation_1499083597995
- whom-to-deinfibulate-at-birth_1499083681432
- when-deinfibulation-at-birth_1499083768252
- deinfibulation-and-second-stage_1499083842981
- episiotomy_1499083920090
- episiotomy-procedure_1499083991489
- forbidden-re-infibulation_1499084226935
- wash-after-deinfibulation_1499084307332
- separate-labia_1499084395928
- re-infibulation-no-health-benefits_1499084511660
- re-infibulation-is-a-health-risk_1499084576800
- healthcare-worker-fgm-knowledge_1499085160949
- no-fgm-for-newborn-girl-_1499085427989