

## Modulewise Data of LagId : ee722f96-fcf6-4bcf-9f4e-c5fd285eaac3

Module: Post Abortion Care

Icon: /icon/module/post\_abortion\_care

Module Key: post-abortion-care\_1487676484604

Action Cards:

- definitions\_1486843717068
- emergency-management\_1486843944185
- following-management\_1486845338095
- emergency-referral\_1490716065855

Procedures:

- cleaning-manual-vacuum-aspirator\_1487245553833
- urine-catheter\_1487245590353
- intravenous-access-equipment\_1487245827833

Videos:

- /India english/Post abortion care/intro
- /India english/Post abortion care/how\_to\_diagnose
- /India english/Post abortion care/how\_to\_manage
- /India english/Post abortion care/evacuation\_of\_the\_uterus
- /India english/Post abortion care/manual\_vacuum\_aspiration

Drugs:

- ampicillin\_1486975748548
- diazepam\_1487102080825
- gentamicin\_1487679704293
- ibuprofen\_1490364323757
- metronidazole\_1487112262252

- morphine\_1487151518002

### Key Learning Points:

- test-version\_1498639439941
- septic-abortion-\_1504773307717
- complication-to-unsafe-abortion\_1504773499553
- urine-test\_1504773825842
- complication-to-abortion\_1504774087137
- fever-and-septic-abortion\_1504774293226
- what-to-measure-\_1504774413976
- signs-of-septic-abortion\_1504887365737
- suspect-lower-abdominal-tenderness\_1504887830839
- septic-abortion-can-be-fatal\_1504888032303
- most-important-action-at-septic-abortion\_1504889099044
- action-at-septic-abortion\_1504889244498
- ab-and-evacuation-of-the-uterus\_1504889778909
- which-ab-to-give\_1504889919542
- urgent-management-\_1504890132226
- correct-order-of-actions\_1504890487022
- peritonitis\_1504890766676
- evacuation-of-the-uterus-\_1504891033708
- manual-vacuum-aspiration\_1504891220598
- uterus-size-in-week-12\_1504891361270
- preparation-for-mva\_1504891579618
- analgesic-for-mva\_1504891713356
- valves-closed\_1504891839206
- vacuum-to-mva-syringe\_1504891916767
- tenaculum\_1504892022670
- dilate-the-cervix\_1504892212948
- no-touch-technique\_1504892332161
- size-of-cannula-for-mva\_1504893593707
- insert-of-the-cannula\_1504893750649
- attachment-of-the-syringe\_1504893864011
- rotate-and-move-cannula-back-and-forth\_1504893996349

- end-of-suctioning\_1504894096498
- use-the-cannula-again\_1504894223366
- ab-after-mva\_1504894299772
- counselling\_1504894423681
- advise-before-dismissal\_1504894568322

## Module: Infection Prevention

Icon: /icon/module/infection\_prevention

Module Key: infection-prevention\_1487676001934

## Procedures:

- mixing-alcohol-based-handrub\_1487245431108
- mixing-chlorine-solution\_1487245480781

## Videos:

- /India english/Infection prevention/intro
- /India english/Infection prevention/hand\_hygiene
- /India english/Infection prevention/procedure\_for\_handwash
- /India english/Infection prevention/procedure\_for\_handrub
- /India english/Infection prevention/  
personal\_protective\_equipment
- /India english/Infection prevention/cleaning\_decontamination
- /India english/Infection prevention/storage
- /India english/Infection prevention/housekeeping
- /India english/Infection prevention/hiv\_hepatitis\_b\_prevention

## Drugs:

- post-exposure-prophylaxis---pep\_1490708337530

## Key Learning Points:

- prevent-infections\_1497356424386

- hand-hygiene-wash-and-alcohol\_1497356912190
- use-alcohol-based-handrub\_1497357082394
- soap-for-handwashing\_1497357310306
- personal-protective-equipment\_1497357545527
- gloves-for-different-patients\_1497357682494
- soap-solution\_1497357821040
- mix-chlorine-solution\_1497357928324
- package-of-sterile-items\_1497358040371
- storage-of-sterile-items\_1497358161852
- cleaning-and-delivery-bed\_1497358695244
- recommended-vaccination\_1497358858644
- test\_1497612420029
- test-1\_1497612450477
- test-2\_1497612458572
- test-3\_1497612477070
- testing-order\_1497875988625
- infections-and-maternal-death\_1498119404097
- infection-and-neonatal-death\_1498119889065
- infection-and-germs\_1498122575522
- healthcare-workers-hands\_1498122780668
- alcohol-most-effective\_1498123474560
- no-jewellery-on-hands\_1498129645819
- towel\_1498130033393
- when-hand-hygiene\_1498130470830
- soak-items-in-soap-solution\_1498130755471
- order-of-cleaning-and-storage\_1498131043643
- disinfection-solution\_1498131845883
- spread-of-hiv-and-hep-b\_1498132068449
- when-hand-hygiene\_1498206602602
- wash-hands\_1498466635307
- procedure-for-handwashing\_1498466869003
- amount-of-handrub\_1498468033321
- product-in-alcohol-based-handrub\_1498468332125
- label-of-alcohol-based-handrub\_1498468847400
- removing-gloves\_1498469162217

- gloves-for-birth\_1498469312963
- cleaning-and-decontamination\_1498469480918
- high-risk-area\_1498469624522
- transmission-most-likely\_1498469768452
- action-if-exposed\_1498469888180
- soap-wet-container\_1504268064997
- air-dry-hands\_1504268482403
- product-in-alcohol-based-handrub-2\_1511791111025

Module: Hypertension

Icon: /icon/module/hypertension

Module Key: hypertension\_1487677992452

Action Cards:

- definitions\_1486814136250
- management-hypertension\_1486815930026
- management-eclampsia\_1486815158568
- emergency-referral\_1490716065855

Procedures:

- urine-catheter\_1487245590353
- intravenous-access-equipment\_1487245827833

Videos:

- /India english/Hypertension/intro
- /India english/Hypertension/definitions
- /India english/Hypertension/causes
- /India english/Hypertension/how\_to\_diagnose
- /India english/Hypertension/management\_of\_hypertension
- /India english/Hypertension/severe\_preeclampsia\_eclampsia

## Drugs:

- calcium-gluconate\_1490273124941
- hydralazine\_1487109388353
- labetalol\_1490625535572
- magnesium-sulphate\_1487111473832
- methyldopa\_1487111904464
- nifedipine\_1487151847555

## Key Learning Points:

- severe-preeclampsia\_1498640805283
- gestational-hypertension\_1498725687229
- bp-gestational-hypertension\_1498725828989
- preeclampsia\_1498726056310
- severely-elevated-blood-pressure\_1498726396275
- severe-headache\_1498726747222
- visual-disturbances\_1498727185096
- upper-abdominal-pain\_1498727830882
- respiratory-distress\_1498728011726
- oliguria\_1498728193444
- eclampsia\_1498728592426
- eclampsia-normal-blood-pressure\_1498728833764
- differential-diagnosis\_1498728983703
- causes-of-pre-eclampsia\_1498729147494
- anc-blood-pressure\_1498729463504
- confirm-bp\_1498732684396
- confirmed-elevated-bp\_1498733012381
- eclampsia-maternal-and-perinatal-death\_1498733194577
- keep-systolic-bp-below-160\_1498733307152
- keep-diastolic-bp-above-80\_1498733486432
- labetalol\_1498733591134
- methyldopa\_1498733669777
- iv-with-bp-above-160-\_1498733752579
- hydralazine\_1498733943552
- eclampsia-seizures\_1498734086062

- magnesium-sulphate-24-hours\_1498734258534
- severe-pre-eclampsia-or-eclampsia\_1498734362115
- hypertension-and-urine-catheter\_1498735254202
- eclampsia---monitor-12-hour\_1498735362891
- give-magnesium-sulphate\_1498735437004
- mix-magnesium-sulphate\_1498735538053
- toxicity\_1498735622169
- antidote-to-magnesium-sulphate\_1498735735523
- maintain-dose-magnesium-sulphate\_1498735869673
- refer-and-eclampsia\_1498735994297
- lung-maturity\_1498736081492
- proteinuria\_1505054086730
- cure-of-pre-eclampsia\_1505055063577
- maternal-death-and-pre-eclampsia\_1505055560037
- action-eclampsia\_1505059069497
- fluid-overload\_1505059297071
- calcium-gluconate\_1505059639547

Module: Normal Labour and Birth

Icon: /icon/module/infection\_prevention

Module Key: normal-labour-and-birth\_1592462784361

Module: Active Management of Third Stage Labour

Icon: /icon/module/active\_management\_of\_third\_stage\_labour

Module Key: active-management-of-third-stage-labour\_1487678773861

Action Cards:

- emergency-referral\_1490716065855

Videos:

- /India english/Active management of third stage labour/intro
- /India english/Active management of third stage labour/

give\_a\_uterotonic\_drug

- /India english/Active management of third stage labour/  
assess\_the\_baby

- /India english/Active management of third stage labour/  
clamp\_and\_cut\_the\_umbilical\_cord

- /India english/Active management of third stage labour/  
controlled\_cord\_traction

- /India english/Active management of third stage labour/  
examine\_the\_placenta

- /India english/Active management of third stage labour/  
monitoring

- /India english/Active management of third stage labour/  
early\_breastfeeding

Drugs:

- antibiotic-eye-drops\_1490261954893

- misoprostol\_1487112562173

- oxytocin\_1487156988152

- vitamin-k\_1487160613336

Key Learning Points:

- test-1\_1497203846089

- active-management\_1499780658036

- uterotonic-drug\_1499780861769

- oxytocin\_1499780939312

- exclude-second-baby\_1499781002066

- dose-of-oxytocin\_1499781138851

- when-to-give-oxytocin\_1499781203804

- storage-of-oxytocin\_1499781280757

- misoprostol\_1499781426287

- administration-of-misoprostol\_1499781506837

- when-to-assess-the-newborn\_1499781574406

- thick-meconium\_1499782019890



- suction-thick-meconium\_1499782064819
- dry-of-the-baby\_1499782139883
- wipe-of-baby\_1499782205916
- warm-baby\_1499782287277
- newborn-not-breathing-well\_1499782350880
- stable-newborn\_1499782411878
- clamp-umbilical-cord\_1499782532294
- cord-traction\_1499782592747
- counter-pressure-pubic-bone\_1499782667213
- rounded-uterus\_1499782790928
- push-for-the-placenta\_1499782842943
- check-uterus\_1499782915737
- measure-blood-loss\_1499783000460
- examine-the-placenta\_1499783053169
- maternal-surface-of-placenta\_1499783139366
- no-unattended-mother-and-newborn\_1499783196382
- skin-to-skin-contact\_1499783293686
- check-mother-after-birth\_1499783355207
- how-often-check-mother\_1499783434630
- how-often-check-newborn\_1499783488277
- check-the-newborn-after-birth\_1499783546639
- cord-monitor\_1499783616183
- eye-drop-and-vit-k\_1499783708999
- early-breastfeeding-newborn\_1499783782397
- early-breastfeeding-mother\_1499783921892
- active-management\_1504529794145
- delivery-of-membranes\_1504531639638
- estimation-of-blood-loss\_1504531689289

Module: Prolonged Labour

Icon: /icon/module/prolonged\_labour

Module Key: prolonged-labour\_1487678855904

Action Cards:

- indications\_1486847771915
- prerequisites\_1486847909849
- procedures-by-also\_1486848037447
- emergency-referral\_1490716065855

#### Procedures:

- urine-catheter\_1487245590353
- foetal-heart-rate\_1487245616159
- vacuum-extractor\_1487245649303
- intravenous-access-equipment\_1487245827833

#### Videos:

- /India english/Prolonged labour/intro
- /India english/Prolonged labour/causes
- /India english/Prolonged labour/1st\_stage\_latent\_phase
- /India english/Prolonged labour/1st\_stage\_active\_phase
- /India english/Prolonged labour/2nd\_stage\_prolonged
- /India english/Prolonged labour/2nd\_stage\_vacuum\_extraction

#### Drugs:

- ampicillin\_1486975748548
- diazepam\_1487102080825
- ergometrine\_1487103846814
- iron\_1487110835869
- misoprostol\_1487112562173
- morphine\_1487151518002
- oxytocin\_1487156988152

#### Key Learning Points:

- complication-prolonged-mother\_1500303341759
- complication-prolonged-infant\_1500303472241

- meaning-of-prolonged\_1500303544246
- four-ps\_1500303686918
- power\_1500303769050
- power-seconds\_1500303835797
- pelvis-passage\_1500303897179
- small-pelvis\_1500303985877
- small-pelvis-refer\_1500304101224
- passenger\_1500304149296
- psyche\_1500304254893
- normal-progression-first-stage\_1500304315481
- false-labour\_1500304449630
- false-labour-and-urinary-tract-infections\_1500304515323
- start-of-active-phase\_1500304605377
- start-partograph\_1500304739350
- vaginal-examination-interval\_1500304824357
- when-prolonged-first-labour\_1500304876769
- action-for-prolonged-first-labour\_1500304936911
- reassess-in-two-hours\_1500305041698
- cervix-fully-dilated\_1500305095706
- push\_1500305543117
- push-for-60-min\_1500305592951
- vacuum-after-60-min\_1500305658715
- indication-for-vacuum-extraction-\_1500305701962
- assess-of-fhr\_1500305759343
- critical-fhr\_1500305923791
- condition-for-vacuum\_1500305971761
- ga-less-37-at-a-hospital\_1500306017747
- risk-of-vacuum-extraction\_1500306057945
- preparation-of-vacuum\_1500306101811
- address-the-patient\_1500306177213
- flexion-point\_1500306221884
- steady-traction\_1500306294118
- give-up-vacuum-delivery\_1500306346908
- incision-rarely\_1500306426261
- performing-vacuum\_1500306468033

- vacuum-extraction-fails\_1500306519814
- cephalic-occiput-posterior\_1504777759514
- more-comfortable\_1504777788729
- prolonged-first-stage-latent-phase\_1504777971422
- contractions-causes-if-not-prolonged-latent-phase\_1504777990862
- prolonged-first-stage-active-phase\_1504778019268
- progress-of-labour\_1504778029127
- partograph-with-dilatation-between-alert-and-action-line-\_1504778050950
- cervical-dilatation\_1504778060957
- immediate-delivery\_1504778233462
- head-above-the-pubic-bone\_1504778275952
- posterior-fontanelle-where\_1504778354078
- posterior-fontanelle\_1504778363034
- placing-the-cup\_1504778371941
- check-for-maternal-tissue\_1504778388031
- flexion-point\_1504778425046
- intact-perineum\_1504778470373
- prolonged-second-stage\_1504778541913

Module: Post Partum Hemorrhage

Icon: /icon/module/post\_partum\_hemorrhage

Module Key: post-partum-hemorrhage\_1487678905876

Action Cards:

- 1-uterus\_1486846813345
- 2-legs\_1486846790935
- 3-arms\_1486846769297
- emergency-referral\_1490716065855

Procedures:

- urine-catheter\_1487245590353

- intravenous-access-equipment\_1487245827833
- condom-balloon-tamponade\_1490714664807

#### Videos:

- /India english/Post partum hemorrhage/intro
- /India english/Post partum hemorrhage/uterus\_massage
- /India english/Post partum hemorrhage/empty\_the\_bladder
- /India english/Post partum hemorrhage/bimanual\_compression
- /India english/Post partum hemorrhage/resuscitation
- /India english/Post partum hemorrhage/give\_uterotonic\_drugs
- /India english/Post partum hemorrhage/trauma
- /India english/Post partum hemorrhage/tissue
- /India english/Post partum hemorrhage/monitoring\_after\_bleeding

#### Drugs:

- oxytocin\_1487156988152
- ergometrine\_1487103846814
- misoprostol\_1487112562173
- morphine\_1487151518002
- diazepam\_1487102080825
- ampicillin\_1486975748548
- iron\_1487110835869

#### Key Learning Points:

- maternal-death-\_1504532310198
- loss-of-tone\_1504532426966
- call-for-help-then-stop-the-bleeding-\_1504532471069
- symptoms-for-pph\_1504532547937
- occult-bleeding\_1504532588274
- moderate-bleeding---characteristics-\_1504532614848
- moderate-bleeding-actions-act-differently-depending-on-the-character-of-the-bleeding\_1504532653096

- uterine-massage-\_1504532689921
- empty-the-bladder-\_1504532728261
- vital-signs\_1504532772329
- strong-bleeding--characteristics\_1504532942008
- bimanual-compression\_1504532979962
- bimanual-compression-procedure\_1504533043410
- bimanual-compression-type-of-gloves-\_1504533076462
- pph-drugs\_1504533111438
- misoprostol-\_1504533151987
- ergometrine-\_1504533241147
- prioritize-manual-maneuvers\_1504533285341
- bleeding-from-a-vaginal-tear-\_1504533312898
- always-do-a-rectal-exploration-\_1504533348068
- placental-tissue-missing-\_1504533398776
- treatment-if-there-is-placental-tissue-missing-\_1504533431739
- removal-tissue-1\_1504533465688
- removal-tissue-2\_1504533503753
- when-to-measure-hemoglobin-\_1504533536483
- observation-after-treatment-pph-\_1504533565253
- cbt-is-indicated-\_1504533594043
- the-causes-of-vaginal-bleeding-\_1504533620719
- positioning-pph\_1504533661330
- positioning---vital-organs-\_1504533702197
- actions-helper-at-the-head\_1504533732416
- actions-helper-at-the-arm\_1504533786043
- referral-pph\_1504533836750
- iv-fluids-pph\_1504533868169
- oxytocin-storage\_1504533913864
- cardiovascular-symptoms-pph\_1504533944971
- blood-inside-uterus\_1504534828505

Module: Manual Removal of Placenta

Icon: /icon/module/manual\_removal\_of\_placenta

Module Key: manual-removal-of-placenta\_1487678949720

### Action Cards:

- emergency-referral\_1490716065855

### Procedures:

- urine-catheter\_1487245590353
- intravenous-access-equipment\_1487245827833
- condom-balloon-tamponade\_1490714664807

### Videos:

- /India english/Manual removal of placenta/intro
- /India english/Manual removal of placenta/aorta\_compression
- /India english/Manual removal of placenta/resuscitation
- /India english/Manual removal of placenta/drugs\_before\_removal
- /India english/Manual removal of placenta/gloves
- /India english/Manual removal of placenta/remove\_the\_placenta
- /India english/Manual removal of placenta/drugs\_after\_removal
- /India english/Manual removal of placenta/monitoring

### Drugs:

- ampicillin\_1486975748548
- diazepam\_1487102080825
- ergometrine\_1487103846814
- iron\_1487110835869
- morphine\_1487151518002
- misoprostol\_1487112562173
- oxytocin\_1487156988152

### Key Learning Points:

- time-before-placenta-be-born\_1500550057787
- remove-manually\_1500550146760
- act-fast\_1500550214224

- life-threatening\_1500550548020
- overall-action\_1500550609958
- when-aorta-compression\_1500550713329
- mother-in-shock\_1500550773682
- how-aorta-compression\_1500550835318
- adequate-aorta-compression\_1500550901996
- maintain-compression\_1500550952103
- woman-lie-down\_1500551007297
- vital-signs\_1500551043524
- iv-fluid\_1500551093153
- fluid-gives-fast-\_1500551155005
- blood-transfusion\_1500551204516
- pain-relief\_1500551251327
- pain-relief-and-aorta-compression\_1500551299297
- gloves-and-manual-removal-of-placenta\_1500551362316
- womans-consent\_1500551424020
- first-hand\_1500551465462
- second-hand\_1500551522653
- right-order-for-removal-of-placenta\_1500551561159
- withdrawing-and-second-hand\_1500551610790
- fragile-uterine-wall\_1500551652340
- placenta-detached\_1500551743553
- palpate-the-uterine-cavity\_1500551802100
- the-overall-action-to-control-pph\_1500551855322
- condom-tamponade\_1500551920106
- bolus-drug-after-removal-of-placenta\_1500551982282
- substituted-drug-for-oxytocin\_1500552127586
- oxytocin-drip-after-removal-of-placenta\_1500552209102
- how-to-mix-oxytocin-drip\_1500552277806
- massage-the-fundus\_1500552326288
- day-after-recommendations\_1500552382430
- refer-manual-removal\_1500552464602
- ab-to-newborn\_1500553942631
- manual-removal-bemonc\_1504534513078
- when-manual-removal\_1504534643086



- heavy-bleeding\_1504534768047
- excessive-bleeding\_1504535064090
- stop-bleeding-retained-placenta\_1504535188380
- resuscitation\_1504535401875
- stop-compression-at-resuscitation\_1504682726957
- how-to-manually-remove-the-placenta\_1504686597840
- aorta-compression-and-manual-removal\_1504686794869
- repeat-intrauterine-palpation\_1504687023070
- manual-compression-ended-aorta-compression\_1504687099431
- invasive\_1504687163659
- after-removal-of-placenta\_1504687463704
- action-right-after-removal-of-placenta\_1504687629838

Module: Maternal Sepsis

Icon: /icon/module/maternal\_sepsis

Module Key: maternal-sepsis\_1487679043550

Action Cards:

- definitions\_1486838289048
- emergency-management\_1486839145753
- following-management\_1486840964786
- identify-cause\_1486841405518
- emergency-referral\_1490716065855

Procedures:

- urine-catheter\_1487245590353
- intravenous-access-equipment\_1487245827833

Videos:

- /India english/Maternal sepsis/intro
- /India english/Maternal sepsis/how\_to\_diagnose\_definitions

- /India english/Maternal sepsis/how\_to\_manage
- /India english/Maternal sepsis/following\_management
- /India english/Maternal sepsis/origin\_of\_infection
- /India english/Maternal sepsis/hiv

#### Drugs:

- ampicillin\_1486975748548
- artesunate\_1487078838130
- betamethasone\_1487080183154
- chloramphenicol\_1487097224306
- dexamethasone\_1487101787771
- dicloxacillin\_1487103485964
- gentamicin\_1487679704293
- metronidazole\_1487112262252
- penicillin-g-benzylpenicillin\_1487157542019
- quinine\_1487158788622

#### Key Learning Points:

- meaning-of-sepsis\_1504944629301
- diagnose-sepsis\_1504944721115
- septic-shock\_1504944788616
- vital-signs\_1504944907638
- importance-of-fluid-management\_1504944997773
- assess-of-fluid-balance\_1504945090408
- fluid-balance-be-calculated\_1504945280195
- fluid-balance-be\_1504945346611
- in-case-of-a-septic-shock-it-should-be-calculated-each-hour\_1504945415855
- complications-to-sepsis\_1504945501258
- initial-management\_1504945569353
- circulation-\_1504945650697
- drugs\_1504945729088
- ampicillin-sepsis\_1504945793742

- gentamicin\_1504945982881
- metronidazole\_1504946039271
- ceftriaxone\_1504946097782
- tetracycline\_1504946176147
- chloramphenicol\_1504946438452
- malaria\_1504946617389
- malaria-drugs\_1504946711304
- artesunate\_1504946886646
- artesunate\_1504946909791
- quinine\_1504947056059
- dosage-artesunate\_1504947115207
- dosage-quinine\_1504947179913
- the-6-ws\_1504947235003
- the-6-ws-continue\_1504947447368
- chorioamnionitis\_1504947765349
- endometritis\_1504947835819
- treatment-for-chorioamnionitis\_1504947946364
- treatment-for-endometritis\_1504948024311
- endometritis-and-no-effect-of-ab\_1504948113282
- mastitis\_1504948177504
- breast-abscess\_1504948229532
- wound-infections\_1504948296628
- other-causes-of-infection\_1504948359563

Module: Neonatal Resuscitation

Icon: /icon/module/neonatal\_resuscitation

Module Key: neonatal-resuscitation\_1487679104279

Action Cards:

- neonatal-resuccitation\_1486842879218
- emergency-referral\_1490716065855

Videos:

- /India english/Neonatal resuscitation/intro
- /India english/Neonatal resuscitation/assess\_the\_baby
- /India english/Neonatal resuscitation/airway
- /India english/Neonatal resuscitation/breathing
- /India english/Neonatal resuscitation/  
when\_to\_stop\_resuscitation
- /India english/Neonatal resuscitation/monitoring

#### Drugs:

- ampicillin\_1486975748548
- ceftriaxone\_1487080443380
- gentamicin\_1487679704293
- phenobarbital\_1487158287290

#### Key Learning Points:

- adequate-breathing\_1504254228875
- golden-minute\_1504254468378
- assess-the-newborn-after-birth\_1504256308739
- skin-to-the-mother\_1504256398872
- first-step\_1504256461191
- meconium-suction-\_1504256548380
- correct-order-suction\_1504256615224
- stimulation-the-newborn\_1504256686246
- newborn-is-not-breathing-well\_1504256807203
- when-to-resuscitate\_1504256982883
- airway-call-for-help\_1504257037350
- airway-position-of-head\_1504257274893
- airway-position-of-mask\_1504258127166
- airway-effective-ventilation\_1504258182479
- number-of-inflation-breaths\_1504258233083
- jaw-thrust\_1504258315698
- successful-inflation-breaths\_1504258364953
- if-no-chest-movements\_1504258411252

- inflation-breaths\_1504258492447
- ventilation-breath\_1504258548225
- not-breathing-well\_1504258620844
- continue-ventilation-breaths\_1504258677078
- most-important-ventilation-breaths\_1504258732578
- rhythm-ventilation-breaths\_1504258780270
- good-ventilation\_1504258849221
- breathing---assessment\_1504258930435
- breathing---assessment\_1504258984771
- breathing-assessment---continued\_1504259182601
- action-if-breathing-well\_1504259239916
- give-oxygen\_1504259318506
- when-to-stop\_1504259374882
- resuscitation-stop-next\_1504259443850
- skin-to-skin-and-breastfeeding\_1504259573821
- monitoring\_1504259668644
- low-birth-weight-and-resuscitation\_1504259731434
- challenge-low-birth-weight\_1504259788567
- successful-inflation-breaths\_1505392891735

Module: Newborn Management

Icon: /icon/module/newborn\_management

Module Key: newborn-management\_1487679126990

Action Cards:

- prevention\_1486813119229
- routine-care\_1486842306870
- danger-signs\_1486807603214
- management-of-danger-signs\_1486808272156
- pediatric-drugs-and-fluids\_1486735917639
- emergency-referral\_1490716065855

Procedures:

- intravenous-access-equipment\_1487245827833
- nasogastric-tube\_1487245852188

#### Videos:

- /India english/Newborn management/intro
- /India english/Newborn management/prevention
- /India english/Newborn management/  
initial\_care\_of\_the\_newborn
- /India english/Newborn management/danger\_signs
- /India english/Newborn management/  
management\_of\_danger\_signs

#### Drugs:

- ampicillin\_1486975748548
- antibiotic-eye-drops\_1490261954893
- betamethasone\_1487080183154
- ceftriaxone\_1487080443380
- dexamethasone\_1487101787771
- gentamicin\_1487679704293
- oxytocin\_1487156988152
- phenobarbital\_1487158287290
- vitamin-k\_1487160613336

#### Key Learning Points:

- common-causes-of-newborn-death\_1500553429571
- death-in-first-week\_1500553499970
- avoidable-death\_1500553555177
- prevent-respiratory-distress-preterm\_1500553601195
- risk-factors-infection-newborn\_1500553681339
- prevent-newborn-infections\_1500553770880
- risk-from-resuscitation\_1500554103700
- action-after-resuscitation\_1500554254443

- early-feeding\_1500554299315
- initial-care-of-the-newborn\_1500554337833
- not-crying-or-breathing-well\_1500554397207
- initial-care-of-the-newborn-\_1500554450890
- initial-care-of-lbw\_1500554526669
- lbw-unable-to-suck-adequately\_1500554570547
- signs-of-illness\_1500554688877
- management-of-danger-signs\_1500554757277
- danger-signs---airway\_1500554845568
- danger-signs---breathing\_1500554890466
- grunting\_1500554955221
- apnoea\_1500554996619
- shock\_1500555055530
- disability\_1500555142461
- danger-signs---exposure\_1500555231782
- jaundice\_1500555273140
- management---case\_1500555311125
- obstructed-airways\_1500555401801
- management---breathing\_1500555450027
- management---circulation\_1500555489191
- management-drowsy-and-unconscious\_1500555531251
- convulsions\_1500555605930
- management-of-danger-signs\_1500555700421
- management---jaundice\_1500555760414
- management---kangaroo-motherhood-care-\_1500555801259
- management---infections\_1500555888659
- no-condition-improved\_1500555927749
- risk-of-low-blood-sugar\_1500555965754
- newborn-cannot-suck\_1500556011761
- rural-facility-newborn-danger-signs\_1500556085207
- kangaroo-care\_1504774517466

Module: Low Birth Weight

Icon: /key learning points/Neonatal resuscitation/

low\_birth\_weight\_preterm

Module Key: low-birth-weight\_1539679206162

Module: Female Genital Mutilation

Icon: /icon/module/female\_genital\_mutilation

Module Key: female-genital-mutilation\_1494414132178

Key Learning Points:

- fgm-is-an-unhealthy-practice\_1499073619747
- fgm-is-a-dangerous-practice\_1499073729938
- fgm-is-forbidden\_1499073844486
- four-types-of-fgm\_1499073935484
- mix-of-the-types\_1499073988266
- type-1\_1499074047139
- type-2\_1499074470165
- type-3\_1499074607765
- type-4\_1499074689724
- type-2-becomes-type-3\_1499075316336
- immediate-health-risk\_1499075598511
- fgm-and-infertility\_1499078717543
- long-term-consequences\_1499078794803
- fgm-antenatal\_1499078908785
- birth-complications\_1499079019921
- why-birth-complications-\_1499079100900
- urethra\_1499079409674
- deinfibulation-and-urethra\_1499082946166
- steps-for-deinfibulation\_1499083113362
- interrupted-sutures\_1499083209086
- analgesia-after-deinfibulation\_1499083297174
- hygiene-for-deinfibulation\_1499083352056
- antenatal-deinfibulation\_1499083466972
- when-antenatal-deinfibulation\_1499083597995
- whom-to-deinfibulate-at-birth\_1499083681432
- when-deinfibulation-at-birth\_1499083768252
- deinfibulation-and-second-stage\_1499083842981



- episiotomy\_1499083920090
- episiotomy-procedure\_1499083991489
- forbidden-re-infibulation\_1499084226935
- wash-after-deinfibulation\_1499084307332
- separate-labia\_1499084395928
- re-infibulation-no-health-benefits\_1499084511660
- re-infibulation-is-a-health-risk\_1499084576800
- healthcare-worker-fgm-knowledge\_1499085160949
- no-fgm-for-newborn-girl-\_1499085427989