

Numerical Mathematics II for Engineers - Homework 6

Group 11

Hetvi Chaniyara, Nelleke Kortleven, Hande Pamuksuz

November 30, 2025

Exercise 6.1

(a)

Deriving the weak form of the Boundary Value Problem (BVP)

Our original equation with the boundary condition is:

$$\begin{aligned} -u''(x) &= f(x) \text{ in } \Omega = (0, 1) \\ u(0) &= u(1) = 0 \end{aligned}$$

where the inhomogeneity is

$$f(x) = \begin{cases} 3, & \text{if } 0 < x < \frac{1}{2}, \\ -1, & \text{if } \frac{1}{2} \leq x < 1. \end{cases}$$

We now choose an arbitrary function v and multiply both sides of our equation by $v(x)$ and integrate over Ω .

$$\int_{\Omega} -u''(x)v(x)dx = \int_{\Omega} f(x)v(x)dx$$

We can employ the integration by parts method to eliminate the second derivative of u .

$$\begin{aligned} \int_{\Omega} -u''(x)v(x)dx &= \int_{\Omega} f(x)v(x)dx \\ [-u'(x)v(x)]_0^1 + \int_{\Omega} u'(x)v'(x)dx &= \int_{\Omega} f(x)v(x)dx \\ (-u'(1)v(1) + u'(0)v(0)) + \int_{\Omega} u'(x)v'(x)dx &= \int_{\Omega} f(x)v(x)dx \end{aligned}$$

The boundary term $(-u'(1)v(1) + u'(0)v(0))$ must be zero to obtain a simplified weak form. This is achieved by carefully defining the space V for the test function v .

We select the function space $V = H_0^1(\Omega)$. The H^1 space (Sobolev space of order 1) is chosen because it ensures that u and v have square-integrable first (weak) derivatives, making the integral $\int_{\Omega} u'(x)v'(x)dx$ well-defined and finite. The crucial subscript '0' restricts v to satisfy the homogeneous essential (Dirichlet) boundary conditions, meaning $v(0) = 0$ and $v(1) = 0$. This ensures the boundary term vanishes: $(-u'(1) \cdot 0 + u'(0) \cdot 0) = 0$, leading directly to the simplified weak form (these results are from the lecture).

The weak/variational problem can be restated as:

$$\text{Find } u \in H_0^1(\Omega) \text{ such that } \int_{\Omega} u'(x)v'(x)dx = \int_{\Omega} f(x)v(x)dx \text{ for all } v \in H_0^1(\Omega)$$

(b)

Since $f(x)$ is piecewise constant, the solution $u(x)$ is a piecewise quadratic polynomial (because if we wanted to find $u(x)$ we would have to integrate $f(x)$ twice). The solution must satisfy the Dirichlet boundary conditions and the C^1 continuity conditions at the interface $x = 1/2$.

We split the domain as defined in $f(x)$. We solve $u''(x) = -f(x)$ in the two regions $\Omega_1 = (0, 1/2)$ and $\Omega_2 = (1/2, 1)$.

$$\begin{aligned} \text{If } 0 < x < 1/2 \text{ (} f(x) = 3 \text{): } u_1''(x) &= -3 \implies u_1(x) = \iint -3dx \\ &= u_1(x) = -\frac{3}{2}x^2 + C_1x + C_2 \\ \text{If } 1/2 < x < 1 \text{ (} f(x) = -1 \text{): } u_2''(x) &= 1 \implies u_2(x) = \iint 2dx \\ &= u_2(x) = \frac{1}{2}x^2 + C_3x + C_4 \end{aligned}$$

We now need to solve for the constants

1. $u(0) = 0 \implies C_2 = 0$ not C_4 because of the domain
2. $u(1) = 0 \implies \frac{1}{2} + C_3 + C_4 = 0 \implies C_3 + C_4 = -1/2$ (I)
3. While H^1 only strictly requires u to be continuous, the successful elimination of the $[-u'(x)v(x)]_0^1$ in the weak problem relies on the continuity of $u' \implies u_1'(1/2) = u_2'(1/2)$:

$$-3/2 + C_1 = 1/2 + C_3$$

$$C_1 - C_3 = 2 \quad \text{(II)}$$

4. $u_1(1/2) = u_2(1/2)$ (because we need continuity of u since $u \in H_0^1(\Omega)$):

$$-\frac{3}{8} + \frac{C_1}{2} = \frac{1}{8} + \frac{C_3}{2} + C_4$$

$$4C_1 - 4C_3 - 8C_4 = 4 \quad \text{(III)}$$

Substituting $C_1 = C_3 + 2$ from (II) into (III) gives:

$$\begin{aligned} 4(C_3 + 2) - 4C_3 - 8C_4 &= 4 \\ 8 - 8C_4 &= 4 \\ C_4 &= 1/2 \end{aligned}$$

Substituting C_4 into (I):

$$\begin{aligned} C_3 + 1/2 &= -1/2 \\ C_3 &= -1 \end{aligned}$$

Finding C_1 from (II):

$$\begin{aligned} C_1 &= -1 + 2 \\ C_1 &= 1 \end{aligned}$$

The solution to the Weak Problem:

$$u(x) = \begin{cases} u_1(x) = -\frac{3}{2}x^2 + x & \text{if } 0 \leq x \leq 1/2 \\ u_2(x) = \frac{1}{2}x^2 - x + \frac{1}{2} & \text{if } 1/2 \leq x \leq 1 \end{cases}$$

(c)

Is the solution of the weak problem also a classical solution of the original BVP?

A function $u(x)$ is a classical solution if it is twice continuously differentiable on the domain, $u \in C^2(\Omega)$, and satisfies the differential equation $-u''(x) = f(x)$ everywhere.

The weak solution $u(x)$ found in (b) is:

$$u(x) = \begin{cases} u_1(x) = -\frac{3}{2}x^2 + x & \text{if } 0 \leq x \leq 1/2 \\ u_2(x) = \frac{1}{2}x^2 - x + \frac{1}{2} & \text{if } 1/2 \leq x \leq 1 \end{cases}$$

The second derivatives are:

$$\begin{aligned} u_1''(x) &= -3 \\ u_2''(x) &= 1 \end{aligned}$$

Since $u_1''(1/2) = -3$ and $u_2''(1/2) = 1$, the second derivative $u''(x)$ is discontinuous at $x = 1/2$. This means $u \notin C^2(\Omega)$.

Because the solution $u(x)$ fails the C^2 smoothness requirement, it is not a classical solution.

However, the weak formulation is necessary precisely because it allows for solutions with lower regularity than C^2 .

- The trial space $u \in H_0^1(\Omega)$ only requires the first derivative u' to be square-integrable.
- Our solution is C^1 -continuous ($u_1'(1/2) = u_2'(1/2) = -1/2$), which is the minimal requirement for the weak formulation to hold, and ensures the flux is continuous across the jump in $f(x)$.

Thus, the weak solution is the mathematical generalisation that holds even when the source term $f(x)$ is not smooth enough to produce a strict classical solution.

Exercise 6.2

(a)

The assumption is

$$|u - u_h|_\infty \leq c h^p$$

Applying this to mesh size at $h/2$, we get the following:

$$|u - u_{h/2}|_\infty \leq c h^p / 2^p$$

The maximum difference between u_h and $u_{h/2}$ can also be written as:

$$|(u - u_{h/2}) - (u - u_h)|_\infty$$

Based on the assumption, the right-hand side can also be completed:

$$|u_h - u_{h/2}| \leq c h^p + \frac{c h^p}{2^p} = c \left(1 + \frac{1}{2^p}\right) h^p$$

We can define $\tilde{c} = c \left(1 + \frac{1}{2^p}\right)$. Then we reach equation (1);

$$\|u_h - u_{h/2}\|_\infty \leq \tilde{c} h^p \quad \square$$

(b)

The assumption is

$$|u - u_h|_\infty \leq c h^p$$

We can copy the logic from part (a) and write this for mesh sizes h , $h/2$ and $h/4$ for experimentation.

Looking at the maximum difference between h and $h/2$, and $h/2$ and $h/4$, we can calculate a ratio of change for each time we halve h . First:

$$u_h - u_{h/2} = (u - u_{h/2}) - (u - u_h) \approx c (h/2)^p - c h^p = c h^p ((1/2)^p - 1)$$

and:

$$u_{h/2} - u_{h/4} = (u - u_{h/4}) - (u - u_{h/2}) \approx c (h/4)^p - c (h/2)^p = c h^p ((1/4)^p - (1/2)^p)$$

The ratio R is:

$$R = \frac{D_1}{D_2} \approx \frac{|c| h^p \left(1 - \frac{1}{2^p}\right)}{|c| \frac{h^p}{2^p} \left(1 - \frac{1}{2^p}\right)} = 2^p.$$

The ratio of the differences between successive solutions is 2^p , where p is the unknown convergence order.

Here we got the experimental order of convergence as

$$p = \log_2(R)$$

Exercise 6.3

(a)

To ensure stability in the maximum norm, we apply the first theorem from Lecture 11. In this theorem, the matrix $M = \tau(1 - \theta)L_h$ has to satisfy the condition $M_{ii} \leq 1$. In 2D, the diagonal entry of L_h is $\frac{4}{h^2}$, so the condition becomes

$$\tau(1 - \theta) \frac{4}{h^2} \leq 1.$$

Substituting $\tau = \frac{T}{M}$ and $h = \frac{1}{N+1}$, we determine the following condition for M :

$$\begin{aligned} \frac{T}{M}(1 - \theta) \frac{4}{\left(\frac{1}{N+1}\right)^2} &\leq 1 \\ \Rightarrow M &\geq T(1 - \theta) \frac{4}{\left(\frac{1}{N+1}\right)^2}. \end{aligned}$$

If $\theta = 1$, the term $1 - \theta$ vanishes and the scheme is unconditionally stable.

(e)




Figure 1: Theta = 0




Figure 2: Theta = 0.5




Figure 3: Theta = 1.0

Figure 4: Comparison of the 2D heat equation solution at $T = 0.05$ using different θ -schemes ($N = 80$).