

STAT 330 - Mathematical Statistics

Cameron Roopnarine

Last updated: September 8, 2020

Contents

Contents	1
2 Random Variable	2
2.1 Probability Model	2
2.2 Random Variable	5

Chapter 2

Random Variable

Review of:

- Probability
- Random variables (discrete and continuous)
- Expectation and variance
- Moment generating function

2.1 Probability Model

DEFINITION 2.1.1: Probability model

A **probability model** is used for a random experiment, which consists of three components:

- (I) Sample space
- (II) Event
- (III) Probability function

DEFINITION 2.1.2: Sample space

A **sample space** S is a set of all the distinct outcomes for a random experiment, with the property that in a single trial, one and only one of these outcomes occurs.

EXAMPLE 2.1.3

Toss a coin twice. This is a random experiment because we do not know the outcome before we toss the coin twice.

- $S = \{(H, H), (H, T), (T, H), (T, T)\}$

Define A : First toss is an H .

Clearly, $A = \{(H, H), (H, T)\} \subseteq S$, so A is an event.

DEFINITION 2.1.4: † Sigma algebra

A collection of subsets of a set S is called **sigma algebra**, denoted by β , if it satisfies the following properties:

- (I) $\emptyset \in \beta$
- (II) If $A \in \beta$, then $\bar{A} \in \beta$
- (III) If $A_1, A_2, \dots \in \beta$, then $\bigcup_{i=1}^{\infty} A_i \in \beta$

DEFINITION 2.1.5: Probability set function

Let β be a sigma algebra associated with the sample space S . A **probability set function** is a function P with domain β that satisfies the following axioms:

- (I) $P(A) \geq 0$ for all $A \in \beta$
- (II) $P(S) = 1$
- (III) *Additivity property:* If $A_1, A_2, A_3, \dots \in \beta$ are pairwise mutually exclusive events; that is, $A_i \cap A_j = \emptyset$ for all $i \neq j$, then

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i)$$

EXAMPLE 2.1.6

Toss a coin twice, given one event A ,

$$P(A) = \frac{\# \text{ of outcomes in } A}{4}$$

since $|S| = 4$. P satisfies the three properties, therefore P is a probability function.

PROPOSITION 2.1.7: Additional Properties of the Probability Set Function

Let β be a sigma algebra associated with the sample space S and let P be a probability set function with domain β . If $A, B \in \beta$, then:

- (1) $P(\emptyset) = 0$
- (2) If A and B are mutually exclusive events, then $P(A \cup B) = P(A) + P(B)$
- (3) $P(\bar{A}) = 1 - P(A)$
- (4) If $A \subset B$, then $P(A) \leq P(B)$

Note for (4), $A \subset B$ means $a \in A$ implies $a \in B$.

Proof of: 2.1.7

Proof of (1): Let $A_1 = S$ and $A_i = \emptyset$ for $i = 2, 3, \dots$. Since $\bigcup_{i=1}^{\infty} A_i = S$, then by (III) it follows that

$$P(S) = P(S) + \sum_{i=2}^{\infty} P(\emptyset)$$

and by (II) we have

$$1 = 1 + \sum_{i=2}^{\infty}$$

By (I) the right side is a series of non-negative numbers which must converge to the left side which is 1 which is finite which results in a contradiction unless $P(\emptyset) = 0$ as required.

Proof of (2): Let $A_1 = A$, $A_2 = B$, and $A_i = \emptyset$ for $i = 3, 4, \dots$. Since $\bigcup_{i=1}^{\infty} A_i = A \cup B$, then by (III)

$$P(A \cup B) = P(A) + P(B) + \sum_{i=3}^{\infty} P(\emptyset)$$

and since $P(\emptyset) = 0$ by the result of (1) it follows that

$$P(A \cup B) = P(A) + P(B)$$

Proof of (3): Since $S = A \cup \bar{A}$ and $A \cap \bar{A} = \emptyset$ then by (II) and by (2) it follows that

$$1 = P(S) = P(A \cup \bar{A}) = P(A) + P(\bar{A})$$

as required.

Proof of (4): Since

$$B = (A \cap B) \cup (\bar{A} \cap B) = A \cup (\bar{A} \cap B)$$

and $A \cap (\bar{A} \cap B) = \emptyset$ then by (2)

$$P(B) = P(A) + P(\bar{A} \cap B)$$

But by (1), $P(\bar{A} \cap B) \geq 0$, so the result now follows.

EXERCISE 2.1.8

Let β be a sigma algebra associated with the sample space S and let P be a probability set function with domain β . If $A, B \in \beta$ then prove the following:

1. $0 \leq P(A) \leq 1$
2. $P(A \cap \bar{B}) = P(A) - P(A \cap B)$
3. $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

1. $P(A) \geq 0$ follows from (I). From (3) we have $P(\bar{A}) = 1 - P(A)$. But from (I) $P(\bar{A}) \geq 0$ and therefore $P(A) \leq 1$.
2. Since $A = (A \cap B) \cup (A \cap \bar{B})$ and $(A \cap B) \cap (A \cap \bar{B}) = \emptyset$, then by (2)

$$P(A) = P(A \cap B) + P(A \cap \bar{B})$$

as required.

3. $P(A \cup B) = (A \cap \bar{B}) \cup (A \cap B) \cup (\bar{A} \cap B)$. By the previous result,

$$P(A \cap \bar{B}) = P(A) - P(A \cap B) \text{ and } P(\bar{A} \cap B) = P(B) - P(A \cap B)$$

Therefore,

$$\begin{aligned} P(A \cup B) &= (P(A) - P(A \cap B)) + P(A \cap B) + (P(B) - P(A \cap B)) \\ &= P(A) + P(B) - P(A \cap B) \end{aligned}$$

as required.

DEFINITION 2.1.9: Conditional probability

Let β be a sigma algebra associated with the sample space S and suppose $A, B \in \beta$ with $P(B) > 0$. Then the **conditional probability** of A given that B has occurred is

$$P(A | B) = \frac{P(A \cap B)}{P(B)}$$

DEFINITION 2.1.10: Independent events

Let β be a sigma algebra associated with the sample space S and suppose $A, B \in \beta$. A and B are **independent events** if

$$P(A \cap B) = P(A)P(B)$$

Clearly, $P(A | B) = P(A)$ if A and B are independent since

$$P(A | B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A)P(B)}{P(B)} = P(A)$$

EXAMPLE 2.1.11

Toss a coin twice.

- A : First toss is H
- B : Second toss is T

$$P(A) = \frac{\# \text{ of outcomes in } A}{4} = \frac{2}{4}$$

also

$$P(B) = \frac{2}{4}$$

$$P(A \cap B) = \frac{1}{4} = P(A)P(B)$$

therefore A and B are independent.

2.2 Random Variable

DEFINITION 2.2.1: Random variable

A **random variable** X is a function from a sample space S to the real numbers \mathbb{R} ; that is,

$$X : S \rightarrow \mathbb{R}$$

satisfies for any given $x \in \mathbb{R}$ $\{X \leq x\}$ is an event.

$$\{X \leq x\} = \{\omega \in S : X(\omega) \leq x\} \subseteq S$$

EXAMPLE 2.2.2

Toss a coin twice. X : # of H in two tosses

Possible values of X : 0, 1, 2. Given $x \in \mathbb{R}$.

$$\{X \leq x\}$$

- $x < 0$ then $\{X \leq x\} = \emptyset$
- $0 \leq x < 1$ then

then

$$\{X \leq x\} = \{X = 0\} = \{(T, T)\} \subseteq S$$

therefore X is a random variable.

DEFINITION 2.2.3: Cumulative distribution function

The **cumulative distribution function** (c.d.f.) of a random variable X is defined by

$$F(x) = P(X \leq x)$$

for all $x \in \mathbb{R}$. Note that the c.d.f. is defined for all \mathbb{R}

DEFINITION 2.2.4: Properties of the cumulative distribution function

- (1) F is a non-decreasing function; that is, if $x_1 \leq x_2$, then $F(x_1) \leq F(x_2)$.

By looking at: $\{X \leq x_1\} \subseteq \{X \leq x_2\}$ if $x_1 \leq x_2$.

- (2) $\lim_{x \rightarrow \infty} F(x) = 1$ and $\lim_{x \rightarrow -\infty} F(x) = 0$.

By looking at: $x \rightarrow \infty$: $\{X \leq x\} \rightarrow S$ $x \rightarrow -\infty$: $\{X \leq x\} \rightarrow \emptyset$

- (3) $F(x)$ is a right continuous function; that is, for any $a \in \mathbb{R}$,

$$\lim_{x \rightarrow a^+} F(x) = F(a)$$

- (4) For all $a < b$

$$P(a < X \leq b) = P(X \leq b) - P(X \leq a) = F(b) - F(a)$$

- (5) For all b

$$P(X = b) = P(\text{jump at } b) = \lim_{t \rightarrow b^+} F(t) - \lim_{t \rightarrow b^-} F(t) = F(b) - \lim_{t \rightarrow b^-} F(t)$$