

LECTURE 14

Estimators, Bias, and Variance

Exploring the different sources of error in the predictions that our models make.

Today's Roadmap

Sample Statistics

Prediction vs. Inference

- Modeling: Assumptions of Randomness

The Bias-Variance Tradeoff

Interpreting Slopes

[Extra] Review of the Bootstrap

[Terminology] Random Variable

Suppose we draw a random sample of size n from a population.

A **random variable** is a numerical function of a sample.


sample was drawn at random

value depends on how the sample came out

- Often denoted with uppercase “variable-like” letters (e.g. X, Y).
- Also known as a sample statistic, or **statistic**.
- Domain (input): all random samples of size n
- Range (output): number line

Descriptive Properties of Random Variables

There are several ways to describe a random variable:



The expectation and variance of a random variable are **numerical summaries** of X .

They are numbers and are not random!

Definition of Expectation

The **expectation** of a random variable X is the **weighted average** of the values of X , where the weights are the probabilities of the values.

Two equivalent ways to apply the weights:

1. One sample at a time:

$$\mathbb{E}[X] = \sum_{\text{all samples } s} X(s)P(s)$$

1. One possible value at a time:

$$\mathbb{E}[X] = \sum_{\text{all possible } x} xP(X = x)$$

} More common (we are usually given the distribution, not all possible samples)

Expectation is a **number, not a **random variable**!**

- It is analogous to the **average** (same units as the random variable).
- It is the center of gravity of the probability histogram.
- It is the long run average of the random variable, if you simulate the variable many times.

Properties of Expectation [1/3]

Instead of simulating full distributions, we often just compute expectation and variance directly.

Recall definitions of expectation:

$$\mathbb{E}[X] = \sum_x x P(X = x)$$

$$\mathbb{E}[X] = \sum_{\text{all samples}} X(s) P(s)$$

Properties:

1. **Expectation is linear.**

Intuition: summations are linear.

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$$

Properties of Expectation [2/3]

Instead of simulating full distributions, we often just compute expectation and variance directly.

Recall definitions of expectation:

$$\mathbb{E}[X] = \sum_x x P(X = x)$$

$$\mathbb{E}[X] = \sum_{\text{all samples}} X(s) P(s)$$

Properties:

1. **Expectation is linear.**

Intuition: summations are linear.

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$$

2. Expectation is linear in sums of RVs,
for any relationship between X and Y.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

Properties of Expectation [3/3]

Instead of simulating full distributions, we often just compute expectation and variance directly.

Recall definitions of expectation:

$$\mathbb{E}[X] = \sum_x xP(X = x)$$

$$\mathbb{E}[X] = \sum_{\text{all samples}} X(s)P(s)$$

Properties:

1. Expectation is linear.

Intuition: summations are linear.

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$$

2. Expectation is linear in sums of RVs,
for any relationship between X and Y.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

3. If g is a non-linear function, then in general $\mathbb{E}[g(X)] \neq g(\mathbb{E}[X])$.

- Example: if X is -1 or 1 with equal probability, then $E^2[X] = 0$ but $E[X^2] = 1 \neq 0$.

Definition of Variance

Variance is the **expected squared deviation from the expectation** of X.

$$\text{Var}(X) = \mathbb{E} [(X - \mathbb{E}[X])^2]$$

- The units of the variance are the square of the units of X.
- To get back to the right scale, use the **standard deviation** of X: $\text{SD}(X) = \sqrt{\text{Var}(X)}$

Variance is a number, not a random variable!

- The main use of variance is to **quantify chance error**. How far away from the expectation could X be, just by chance?

By [Chebyshev's inequality](#) :

- No matter what the shape of the distribution of X is, the vast majority of the probability lies in the interval "expectation plus or minus a few SDs."

Properties of Variance [1/2]

Recall definition of variance:

$$\text{Var}(X) = \mathbb{E} [(X - \mathbb{E}[X])^2]$$




Properties:

1. **Variance is non-linear:**

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Intuition: Consider the
Standard Deviation for $Y = -3X + 2$:

$$\text{SD}(aX + b) = |a| \text{SD}(X)$$



Properties of Variance [2/2]

Recall definition of variance:

$$\text{Var}(X) = \mathbb{E} [(X - \mathbb{E}[X])^2]$$

Properties:

1. Variance is non-linear:

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Intuition (full proof): Consider the Standard Deviation for $Y = -3X + 2$:

$$\text{SD}(aX + b) = |a| \text{SD}(X)$$

2. Variance of sums of RVs is affected by the (in)dependence of the RVs

$$\text{Var}(X_1 + X_2) = \text{Var}(X_1) + \text{Var}(X_2) + 2\text{Cov}(X, Y)$$



Covariance of X and Y (next slide).
If X, Y independent,
then $\text{Cov}(X, Y) = 0$.


From Populations to Samples

If we know the **distribution of a random variable**, we can reliably compute expectation, variance, functions of the random variable, etc.

However, in Data Science, we often collect **samples**.

- We don't know the distribution of our population.
- We'd like to use the distribution of your sample to estimate/infer properties of the population.


The **big assumption** we make in modeling/inference: Our random sample datapoints are **IID**.



The Central Limit Theorem

No matter what population you are drawing from:

If an IID sample of size n is large,
the probability distribution of the **sample mean**
is **roughly normal** with mean μ and SD σ/\sqrt{n} .



Any theorem that provides the rough distribution of a statistic
and **doesn't need the distribution of the population** is valuable to data scientists.

- Because we rarely know a lot about the population!

Properties of the Sample Mean

Consider an IID sample X_1, X_2, \dots, X_n drawn from a numerical population with **mean μ and SD σ** .

Define the **sample mean**:

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

Expectation:

$$\begin{aligned}\mathbb{E}[\bar{X}_n] &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}[X_i] \\ &= \frac{1}{n}(n\mu) = \mu\end{aligned}$$

Variance/Standard Deviation:

$$\begin{aligned}\text{Var}(\bar{X}_n) &= \frac{1}{n^2} \text{Var}\left(\sum_{i=1}^n X_i\right) = \frac{1}{n^2} \underbrace{\left(\sum_{i=1}^n \text{Var}(X_i)\right)}_{\text{IID} \rightarrow \text{Cov}(X_i, X_j) = 0} \\ &= \frac{1}{n^2} (n\sigma^2) = \frac{\sigma^2}{n}\end{aligned}$$

$$\text{SD}(\bar{X}_n) = \frac{\sigma}{\sqrt{n}}$$

Using the Sample Mean to Estimate the Population Mean


Our goal is often to **estimate** some characteristic of a population.

- Example: average height of Cal undergraduates.
- We typically can collect a **single sample**. It has just one average.
- Since that sample was random, it *could have* come out differently.

We should consider the **average value and spread** of all possible sample means, and how it scales with the sample size n .

$$\mathbb{E}[\bar{X}_n] = \mu$$

$$\text{SD}(\bar{X}_n) = \frac{\sigma}{\sqrt{n}}$$



Using the Sample Mean to Estimate the Population Mean


Our goal is often to **estimate** some characteristic of a population.

- Example: average height of Cal undergraduates.
- We typically can collect a **single sample**. It has just one average.
- Since that sample was random, it *could have come out differently*.

We should consider the **average value and spread** of all possible sample means, and what this means for how big n should be.

$$\mathbb{E}[\bar{X}_n] = \mu$$

$$\text{SD}(\bar{X}_n) = \frac{\sigma}{\sqrt{n}}$$



For every sample size, the expected value of the sample mean is the population mean.

We call the sample mean an **unbiased estimator** of the population mean.
(more on this in a bit)

Using the Sample Mean to Estimate the Population Mean

Our goal is often to **estimate** some characteristic of a population.


- Example: average height of Cal undergraduates.
- We typically can collect a **single sample**. It has just one average.
- Since that sample was random, it *could have come out differently*.

We should consider the **average value and spread** of all possible sample means, and what this means for how big n should be.

$$\mathbb{E}[\bar{X}_n] = \mu$$

For every sample size, the expected value of the sample mean is the population mean.

We call the sample mean an **unbiased estimator** of the population mean.
(more on this in a bit)



$$\text{SD}(\bar{X}_n) = \frac{\sigma}{\sqrt{n}}$$

Square root law: If you increase the sample size by a factor, the SD decreases by the square root of the factor.

The sample mean is more likely to be close to the population mean if we have a larger sample size.

Prediction vs. Inference

Sample Statistics

Prediction vs. Inference

- Modeling: Assumptions of Randomness

The Bias-Variance Tradeoff

Interpreting Slopes

[Extra] Review of the Bootstrap

Prediction vs. Inference

Why do we build models?

To make **accurate predictions** about unseen data.

Prediction is the task of using our model to make predictions for the response (output) of unseen data.

To understand **complex phenomena** occurring in the world we live in.

Inference is the task of using our model to draw conclusions about the underlying true relationship(s) between our features and response.

Prediction vs. Inference

Why do we build models?

To make **accurate predictions** about unseen data.

Prediction is the task of using our model to make predictions for the response (output) of unseen data.

To understand **complex phenomena** occurring in the world we live in.

Inference is the task of using our model to draw conclusions about the underlying true relationship(s) between our features and response.

Example: Suppose we are interested in studying the relationship between the value of a home and a view of a river, school districts, property size, income level of community, etc.

Prediction: Given the attributes of some house, how much is it worth?

We care more about making accurate predictions, don't care so much about how.

Inference: How much extra will a house be worth if it has a view of the river?

We care more about having model parameters that are interpretable and meaningful.


[Terminology] Parameters, Statistics, and Estimators

Inference is all about **drawing conclusions** about
population parameters, given only a **random sample**.



[Terminology] Parameters, Statistics, and Estimators

Inference is all about **drawing conclusions** about **population parameters**, given only a **random sample**.



A **parameter** is a numerical function of the **population**.


Population mean
 μ

Sample mean
 $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$

A **statistic** is a numerical function of the random **sample**.

[Terminology] Parameters, Statistics, and Estimators

Inference is all about **drawing conclusions** about **population parameters**, given only a **random sample**.



Random Sampling With Replacement →

X_1, X_2, \dots, X_n
Sample

x
0 6
1 8
2 6
3 6
4 3
...
95 8



Population mean

$$\mu$$

A **parameter** is a numerical function of the **population**.

Estimate

$$\text{np.mean}(\dots) = 5.71$$

Sample statistic

Sample mean

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

A **statistic** is a numerical function of the random **sample**.


We can then use the sample statistic as an **estimator** of the true population parameter.

Since our **sample is random**, our statistic (which we use as our estimator) could have been different.


Example: When we use the sample mean to estimate the population mean, our estimator is almost always going to be somewhat off.

Data Generation Process: Estimating a Value

One View: Randomly draw a random sample, then compute the statistic for that sample.




Another View: Randomly draw from the distribution of the statistic (generated from all possible samples).




We often use the 2nd view because we want to interpret the estimator directly, and not the random sample.

If We Drew a Different Sample, We'd Get A Different Estimator

population




X_1, X_2, \dots, X_n
Sample



Sample mean

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

`np.mean(...)` = 5.74



5.74

The value of our estimator is a function of the random sample. The estimator is therefore also random.

Modeling: Estimating a Relationship


What if we wanted to estimate the relationship between input x and random response Y ?

$$Y = g(x) + \epsilon$$

We would like to find the true relationship g .

Each individual in the population has:

- **Fixed features** x , and hence fixed $g(x)$.



Modeling: Estimating a Relationship

What if we wanted to estimate the relationship between input x and random response Y ?


$$Y = g(x) + \epsilon$$

We would like to find the true relationship g .

Each individual in the population has:

- **Fixed features** x , and hence fixed $g(x)$.
- Random **error/noise** ϵ
- Random **observation/response** $Y = g(x) + \epsilon$

Errors ϵ are assumed expectation 0 ("zero mean")
and i.i.d. across individuals



Modeling: Estimating a Relationship

What if we wanted to estimate the relationship between input x and random response Y ?


$$Y = g(x) + \epsilon$$

We would like to find the true relationship g .

Each individual in the population has:

- **Fixed features** x , and hence fixed $g(x)$.
- Random **error/noise** ϵ
- Random **observation/response** $Y = g(x) + \epsilon$


Errors ϵ are assumed expectation 0 ("zero mean") and i.i.d. across individuals



We only can only observe our random sample. From this we'd like to estimate the true relationship g .

Modeling: Estimating a Relationship

What if we wanted to estimate the relationship between input x and random response Y ?



We would like to find the true relationship g .

Each individual in the population has:

- **Fixed features** x , and hence fixed $g(x)$.
- Random **error/noise** ϵ
- Random **observation/response** $Y = g(x) + \epsilon$

Errors ϵ are assumed expectation 0 ("zero mean") and i.i.d. across individuals

We build a **model** for predictions based on our observed sample of (x, y) pairs. Our model **estimates** the true relationship g .

At every x , our **prediction** for Y is $\hat{Y}(x)$.

Modeling: Estimating a Relationship




If we assume our true relationship g is **linear**, then we express the response as $Y = f_\theta(x)$.



- The true relationship has true, unobservable parameters θ .
- There is still random noise ϵ , so we still can never observe the true relationship.

Estimating a linear relationship

If we assume our true relationship g is **linear**, then we express the response as $Y = f_\theta(x)$.



- The true relationship has true, unobservable parameters θ .
- There is still random noise ϵ , so we still can never observe the true relationship.

- Obtain a sample \mathbb{X}, \mathbb{Y} of n observed relationships (x, Y) .
- Train a model and obtain estimates $\hat{\theta}$.

Which expressions are random?

Suppose we have an individual with fixed input x . Assume the true relationship g is linear.



$$Y = g(x) + \epsilon$$

$$f_{\theta}(x) = \theta_0 + \sum_{j=1}^p \theta_j x_j + \epsilon$$

$$\begin{aligned}\hat{Y}(x) &= f_{\hat{\theta}}(x) \\ &= \hat{\theta}_0 + \sum_{j=1}^p \hat{\theta}_j x_j\end{aligned}$$

- The true relationship has true, unobservable parameters θ .
- There is still random noise ϵ , so we still can never observe the true relationship.

- Obtain a sample \mathbb{X}, \mathbb{Y} of n observed relationships (x, Y) .
- Train a model and obtain estimates $\hat{\theta}$.

Which expressions are random?

Random in blue

Suppose we have an individual with fixed input x . Assume the true relationship g is linear.

$$Y = g(x) + \epsilon$$
$$f_{\theta}(x) = \theta_0 + \sum_{j=1}^p \theta_j x_j + \epsilon$$

- The true relationship has true, unobservable parameters θ .
- There is still random noise ϵ , so we still can never observe the true relationship.

$$\hat{Y}(x) = f_{\hat{\theta}}(x)$$
$$= \hat{\theta}_0 + \sum_{j=1}^p \hat{\theta}_j x_j$$

- Obtain a sample \mathbb{X}, \mathbb{Y} of n observed relationships (x, Y) .
- Train a model and obtain estimates $\hat{\theta}$.

The Bias-Variance Tradeoff

Sample Statistics

Prediction vs. Inference

- Modeling: Assumptions of Randomness


The Bias-Variance Tradeoff

Interpreting Slopes

[Extra] Review of the Bootstrap

The Bias-Variance Tradeoff

This is the abstract version of the plot on the previous slide:



Terms we will define:

- Model Risk
- Observation Variance
- Model Bias
- Model Variance

For a new individual at (x, Y) :

Model Risk is the mean squared prediction error.

$$\text{model risk} = \mathbb{E}[(Y - \hat{Y}(x))^2]$$

Expectation over **multiple** random variables $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n, Y$:

- All possible samples we could have gotten when fitting our model
- All possible new observations at this fixed x

This Entire Section is Theoretical

For a new individual at (x, Y) :

Model Risk is the mean squared prediction error.

$$\text{model risk} = \mathbb{E}[(Y - \hat{Y}(x))^2]$$

Expectation over **multiple** random variables X_1, X_2, \dots, X_n, Y :

- All possible samples we could have gotten when fitting our model
- All possible new observations at this fixed x

Contrast with numerical functions of the actual collected sample:

$$(\text{L2 loss})_i = (y_i - \hat{y}_i)^2$$

- The i -th collected response $Y = y_i$
- The prediction \hat{y}_i using the model you fit to the collected sample

$$\text{empirical risk} = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

Mean Squared error over all individuals in the collected sample $\{(x_1, y_1), \dots, (y_n, y_n)\}$

The Three Components of Model Risk

There are three types of error that contribute to model risk:

1. **Observation variance**,


because Y has random noise ϵ ;

2. **Model variance**,

because sample X_1, X_2, \dots, X_n is random; and

3. **Model bias**,

because our model is different from the true underlying function g .



How do you think each of the types of error are encoded into the diagram?



The Three Components of Model Risk

There are three types of error that contribute to model risk:

1. **Observation variance**,


because Y has random noise ϵ ;

2. **Model variance**,

because sample X_1, X_2, \dots, X_n is random; and

3. **Model bias**,

because our model is different from
the true underlying function g .



We'll spend this section **defining** each component of
the below equation.

$$\text{model risk} = \text{observation variance} + (\text{model bias})^2 + \text{model variance}$$


1. Observation Variance

The observation Y is random because by definition, our observation is noisy.

We assume **random error** ϵ to have zero mean and variance σ^2 .

$$Y = g(x) + \epsilon$$

 random error



1. Observation Variance

The observation Y is random because by definition, our observation is noisy.

We assume **random error** ϵ to have zero mean and variance σ^2 .

Define **observation variance** as the variance of random error:

$$\text{observation variance} = \sigma^2$$

Reasons:


- Measurement error
- Missing information acting like noise

Remedies:

- Could try to get more precise measurements
- But often this is **beyond the control** of the data scientist.

$$Y = g(x) + \epsilon$$


random error





2. Model Variance

Our fitted model is based on a **random sample**.

If the sample came out differently, then the fitted model would have been different.

$\hat{Y}(x)$
Prediction for the individual at x
A random variable



2. Model Variance

Our fitted model is based on a **random sample**.

If the sample came out differently, then the fitted model would have been different.

$\hat{Y}(x)$ Prediction for the individual at x .
A random variable

Define the **model variance** as the variance of our prediction at x :


$$\text{model variance} = \text{Var}(\hat{Y}(x))$$

Main Reason:

- **Overfitting**. Small differences in random samples lead to large differences in the fitted model

Remedy:

- Reduce model complexity
- Don't fit the noise




3. Model Bias

Define the **model bias** as the difference between our predicted value and the true $g(x)$.

- The fit of our model (for a linear model, the estimate $\hat{\theta}$) is based on a random sample.
- So model bias is averaged over all possible samples.

$\hat{Y}(x)$ Prediction for the individual at x .
A random variable



3. Model Bias


Define the **model bias** as the difference between our predicted value and the true $g(x)$.

- The fit of our model (for a linear model, the estimate $\hat{\theta}$) is based on a random sample.
- So model bias is averaged over all possible samples.

$$\text{model bias} = \mathbb{E}[\hat{Y}(x)] - g(x)$$

- Bias is an average measure for a specific individual x :
- If positive, the model tends to overestimate at this x .
 - If negative, the model tends to underestimate at this x .
 - If zero, the model is **unbiased**.

$\hat{Y}(x)$ Prediction for the individual at x .
A random variable



3. Model Bias

Define the **model bias** as the difference between our predicted value and the true $g(x)$.

- The fit of our model (for a linear model, the estimate $\hat{\theta}$) is based on a random sample.
- So model bias is averaged over all possible samples.

$\hat{Y}(x)$ Prediction for the individual at x .
A random variable


$$\text{model bias} = \mathbb{E}[\hat{Y}(x)] - g(x)$$

Reasons:

- **Underfitting**.
- Lack of domain knowledge.

Remedies:

- Increase model complexity (but don't overfit!)
- Consult domain experts to see which models make sense.



3. [Definition] Unbiased Estimators

$$\text{model bias} = \mathbb{E}[\hat{Y}(x)] - g(x)$$

An **unbiased model** is one where model bias = 0.

In other words, on average, the model predicts $g(x)$.

We can define bias for estimators, too.

For example, the sample mean is an **unbiased estimator** of the population mean.

- By the CLT, $\mathbb{E}[\bar{X}_n] = \mu$.
- Therefore estimator bias = $\mathbb{E}[\bar{X}_n] - \mu = 0$.

The Bias-Variance Decomposition

We've spent this section **defining** each component of the below equation.

model risk = observation variance + (model bias)² + model variance



$$\mathbb{E}[(Y - \hat{Y}(x))^2] = \sigma^2 + \left(\mathbb{E}[\hat{Y}(x)] - g(x) \right)^2 + \text{Var}(\hat{Y}(x))$$


Interpret:

- Model risk is an expectation and is therefore a fixed number (for a given x and model $\hat{Y}(x)$).
- Observation variance is irreducible.
- As models **increase in complexity**, they **overfit** the sample data and will have **higher model variance**. This often corresponds to a decrease in bias.
- As models **decrease in complexity**, they **underfit** the sample data and have lower model variance. This corresponds to an **increase in bias**.

This is the **Bias-Variance Tradeoff**.

The Bias-Variance Tradeoff

model risk = observation variance + (model bias)² + model variance



Interpreting Slopes

Sample Statistics

Prediction vs. Inference



- Modeling: Assumptions of Randomness

The Bias-Variance Tradeoff

Interpreting Slopes


[Extra] Review of the Bootstrap

Inference: The right model structure matters!



Ptolemaic Astronomy, a geocentric model based on circular orbits (epicycles and deferents).

High accuracy but very high model complexity.



Copernicus and Kepler: a heliocentric model with elliptical orbits.

Small model complexity yet high accuracy.

Inference for Linear Regression

Assume the true relationship is linear:

$$f_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_p x_p + \epsilon$$

Unknown true parameters θ



Our estimation from our sample (design matrix \mathbb{X} , response vector \mathbb{Y}):

$$f_{\hat{\theta}}(x) = \hat{\theta}_0 + \hat{\theta}_1 x_1 + \hat{\theta}_2 x_2 + \dots + \hat{\theta}_p x_p$$

Estimated parameters $\hat{\theta}$



The meaning of “slope”:

1. What if the true parameter θ_1 is 0?
2. What does the parameter θ_1 even mean?

What can we **infer** about our true parameter given our estimate $\hat{\theta}_1$?

Inferring Zero slope

Our **estimation** from our sample (design matrix \mathbb{X} , response vector \mathbb{Y}):

$$f_{\hat{\theta}}(x) = \hat{\theta}_0 + \hat{\theta}_1 x_1 + \hat{\theta}_2 x_2 \cdots + \hat{\theta}_p x_p$$



This estimate $\hat{\theta}_1$ for the true parameter θ_1 depends on our sample.

What if the true parameter θ_1 was 0?

Then the feature x_1 has **no effect** on the response!

Approximate confidence interval for true slope

How do we test if the true parameter θ_1 was 0?

- We get one estimate $\hat{\theta}_1$ from our sample of size n .
- But we must imagine all the other random samples that could have happened, and draw our conclusion based on this distribution of estimates.

Enter **hypothesis testing!**

Null hypothesis: The true parameter θ_1 is 0.

Alternative hypothesis: The true parameter θ_1 is not 0.

If your p-value is small, reject the null hypothesis at the cutoff level (say, 5%).

Equivalently ([duality argument](#)):

- Compute an approximate 95% confidence interval with **bootstrapping**.
- If the interval does not contain 0, reject the null hypothesis at the 5% level.
- Otherwise, data are consistent with null hypothesis (the true parameter could be 0).

The Snowy Plover

Data on the tiny [Snowy Plover](#) bird was collected at the Point Reyes National Seashore.

The bigger a newly hatched chick, the more likely it is to survive.



Demo

Assumed true relationship for newborn weight $Y = f_{\theta}(x)$:

$$f_{\theta}(x) = \theta_0 + \theta_1 \text{egg_weight} + \theta_2 \text{egg_length} + \theta_3 \text{egg_breadth} + \epsilon$$

Estimating the Snowy Plover

Assumed true relationship for newborn weight $Y = f_\theta(x)$:

$$f_\theta(x) = \theta_0 + \theta_1 \text{egg_weight} + \theta_2 \text{egg_length} + \theta_3 \text{egg_breadth} + \epsilon$$

Estimated model for newborn weight $\hat{Y} = f_{\hat{\theta}}(x)$:

	theta_hat
$\hat{\theta}_0$	intercept -4.605670
$\hat{\theta}_1$	egg_weight 0.431229
$\hat{\theta}_2$	egg_length 0.066570
$\hat{\theta}_3$	egg_breadth 0.215914

Demo

Is this the right linear model for newborn weight?

Let's test the **null hypothesis**: The true parameter θ_1 is 0.

Bootstrapped Confidence Interval for θ_1

We can estimate the distribution of $\hat{\theta}_1$ by bootstrapping.

Bootstrap the sample to build an **approximate 95% confidence interval** for the parameter θ_1 :

```
sample_df = ... # call this the bootstrap population
n = len(sample_df)
estimates = []
repeat 10000 times:
    # resample n times with replacement
    resample = sample_df.sample(n, replace=True)
    ... # fit the new model to the new resampled X, y
    estimate = get_theta_hat1(model)
    estimates.append(estimate)
lower = np.percentile(estimates, 2.5)
upper = np.percentile(estimates, 97.5)
conf_interval = (lower, upper)
```

Demo

Bootstrapped Confidence Interval for θ_1

We can estimate the distribution of $\hat{\theta}_1$ by bootstrapping.

Bootstrap the sample to build an **approximate 95% confidence interval** for the parameter θ_1 :

Our bootstrapped 95% confidence interval for the true θ_1 :

$$(-0.262, 1.115)$$

Demo

We cannot reject the null hypothesis at cutoff 5%
(our true parameter θ_1 could be 0).



Are all of our true parameters 0?

Let's bootstrap 95% confidence intervals for all our parameters:

True param		lower	upper
θ_0	intercept	-15.457398	5.518540
θ_1	theta_egg_weight	-0.271299	1.136913
θ_2	theta_egg_length	-0.102671	0.212089
θ_3	theta_egg_breadth	-0.271769	0.765737

Demo



Wait....something's off here!



Collinearity

Our estimation from our sample (design matrix \mathbb{X} , response vector \mathbb{Y}):

$$f_{\hat{\theta}}(x) = \hat{\theta}_0 + \hat{\theta}_1 x_1 + \hat{\theta}_2 x_2 \cdots + \hat{\theta}_p x_p$$




The **slope** $\hat{\theta}_p$ measures the change in y per unit change in x_p ,
provided all the other variables are held constant.

$$f_{\theta}(x) = \theta_0 + \theta_1 \text{egg_weight} + \theta_2 \text{egg_length} + \theta_3 \text{egg_breadth} + \epsilon$$

If two chihuahuas have a 1 unit difference in egg weight **and others keep the same**, their estimated weight difference is θ_1 .

If variables are **related** to each other, then **interpretation fails!**
E.g., if a change in egg weight always came with a change in change in length and breadth.



Collinearity


If features are related to each other, it might not be possible to have a change in one of them **while holding the others constant**.

- Then the individual slopes will have no meaning!

Collinearity: When a feature can be predicted pretty accurately by a **linear** function of the others, i.e., the feature is highly correlated with the others.

- Slopes can't be interpreted
- $\mathbb{X}^T \mathbb{X}$ might not be invertible, i.e., solution might not be uniquely determined
- Small changes in the data sample can lead to big changes in the estimated slopes
- Also known as **multicollinearity**

Cross-wise comparison of egg features



```
px.scatter_matrix(eggs)
```

	egg_weight	egg_length	egg_breadth	bird_weight
egg_weight	1.000000	0.792449	0.839077	0.847228
egg_length	0.792449	1.000000	0.402764	0.676142
egg_breadth	0.839077	0.402764	1.000000	0.733687
bird_weight	0.847228	0.676142	0.733687	1.000000

```
eggs.corr()
```

Demo

A more interpretable model

If we instead assume a true relationship using only egg weight:

$$f_{\theta}(x) = \hat{\theta}_0 + \hat{\theta}_1 \text{egg_weight} + \epsilon$$

theta_hat
$\hat{\theta}_0$
intercept -0.058272
$\hat{\theta}_1$
egg_weight 0.718515

This model performs almost as well as our other model (RMSE 0.0464, old RMSE 0.0454), and the confidence interval for the true parameter $\hat{\theta}_1$ doesn't contain zero:

$$(0.604, 0.819)$$

Demo

In retrospect, it's no surprise that the weight of an egg best predicts the weight of a newly-hatched chick.

A model with **highly correlated variables** prevents us from interpreting how the variables are related to the prediction.

Reminder: Assumptions matter

Keep the following in mind:

- All inference assumes that the regression model holds.
 - If the model doesn't hold, the inference might not be valid.
 - If the [assumptions of the bootstrap](#) don't hold...
 - Sample size n is large
 - Sample is representative of population distribution (drawn IID, unbiased)
- ...then the results of the bootstrap might not be valid.

[Extra] Review of the Bootstrap

Sample Statistics

Prediction vs. Inference

- Modeling: Assumptions of Randomness

The Bias-Variance Tradeoff

Interpreting Slopes

[Extra] Review of the Bootstrap



- To determine the properties (e.g. variance) of the sampling distribution of an estimator, we'd need to have access to the population.
 - We would have to consider all possible samples, and compute an estimate for each sample.
- But we don't, we only have one random sample from the population.

Idea: Treat our random sample as a “population”, and resample from it.


- Intuition: a random sample resembles the population, so a random resample resembles a random sample.

The Distribution of an Estimator

Resampling the population to estimate the sample distribution.




Variability in my estimation procedure.




Bootstrap the Distribution of an Estimator

Simulation method to estimate the sample distribution.




Bootstrap the Distribution of an Estimator

Simulation method to estimate the sample distribution.




Bootstrap resampling is a technique for estimating the sampling distribution of an estimator.

Impractical:



Bootstrap:



Bootstrapping pseudocode

```
collect random sample of size n (called the bootstrap population)
initiate list of estimates
repeat 10,000 times:
    resample with replacement n times from bootstrap population
    apply estimator f to resample
    store in list
list of estimates is the bootstrapped sampling distribution of f
```

Why **must** we resample
with replacement?

The **bootstrapped sampling distribution of an estimator** does not exactly match the **sampling distribution of that estimator**.

- The center and spread are both wrong (but often close).

The center of the bootstrapped distribution is the estimator applied to our original sample.

- We have no way of recovering the estimator's true expected value.

The variance of the bootstrapped distribution is often close to the true variance of the estimator.

The quality of our bootstrapped distribution depends on the quality of our original sample.

- If our original sample was not representative of the population, bootstrap is next to useless.