

? This sculpture of a woolly mammoth, just 3.7 cm (1.5 in.) in length, was carved from a mammoth's ivory tusk by an artist who lived in southwestern Germany 35,000 years ago. It is possible to date biological specimens such as this because (i) ancient materials were more radioactive than modern ones; (ii) ancient materials were less radioactive than modern ones; (iii) biological specimens continue to take in radioactive substances after they die; (iv) biological specimens no longer take in radioactive substances after they die; (v) more than one of these.



43 Nuclear Physics

LEARNING OUTCOMES

In this chapter, you'll learn...

- 43.1 Some key properties of atomic nuclei, including radii, densities, spins, and magnetic moments.
- 43.2 How the binding energy of a nucleus depends on the numbers of protons and neutrons that it contains.
- 43.3 The most important ways in which unstable nuclei undergo radioactive decay.
- 43.4 How the decay rate of a radioactive substance depends on time.
- 43.5 Some of the biological hazards and medical uses of radiation.
- 43.6 How to analyze some important types of nuclear reactions.
- 43.7 What happens in a nuclear fission chain reaction, and how it can be controlled.
- 43.8 The nuclear reactions that allow the sun to shine.

You'll need to review...

- 5.5 The strong interaction.
- 18.3 Kinetic energy of gas molecules.
- 21.1 Proton and neutron.
- 26.4 Discharging capacitor.
- 37.8 Rest mass and rest energy.
- 39.2 Discovery of the nucleus.
- 40.3, 40.4 Square-well potential; tunneling.
- 41.4–41.6 Magnetic moments; spin- $\frac{1}{2}$ particles; central-field approximation.

Every atom contains at its center an extremely dense, positively charged *nucleus*, which is much smaller than the overall size of the atom but contains most of its total mass. In this chapter we'll look at several important general properties of nuclei and of the nuclear force that holds protons and neutrons together within a nucleus. The stability or instability of a particular nucleus is determined by the competition between the attractive nuclear force among the protons and neutrons and the repulsive electrical interactions among the protons. Unstable nuclei *decay*, transforming themselves spontaneously into other nuclei by a variety of processes. Nuclear reactions can also be induced by impact on a nucleus of a particle or another nucleus. Two classes of reactions of special interest are *fission* and *fusion*. Fission is the process that takes place within a nuclear reactor used for generating power. We could not survive without the energy released by one nearby fusion reactor, our sun.

43.1 PROPERTIES OF NUCLEI

As we described in Section 39.2, Rutherford found that the nucleus is tens of thousands of times smaller in radius than the atom itself. Since Rutherford's initial experiments, many additional scattering experiments have been performed with high-energy protons, electrons, neutrons, and alpha particles. These experiments show that we can model a nucleus as a sphere with a radius R that depends on the total number of *nucleons* (neutrons and protons) in the nucleus. This number is called the **nucleon number** A . The radii of most nuclei are represented quite well by the equation

$$R = R_0 A^{1/3} \quad (43.1)$$

Experimentally determined constant = $1.2 \times 10^{-15} \text{ m} = 1.2 \text{ fm}$
 Radius of an atomic nucleus $\rightarrow R$ $\leftarrow$ Nucleon number = total number of protons and neutrons A

The nucleon number A in Eq. (43.1) is also called the **mass number** because it is the nearest whole number to the mass of the nucleus measured in unified atomic mass units (u).

(The proton mass and the neutron mass are both approximately 1 u.) The best current value is

$$1 \text{ u} = 1.660539040(20) \times 10^{-27} \text{ kg}$$

In Section 43.2 we'll discuss the masses of nuclei in more detail.

Nuclear Density

The volume V of a sphere is equal to $4\pi R^3/3$, so Eq. (43.1) shows that the *volume* of a nucleus is approximately proportional to A . Since the proton and the neutron both have about the same mass, the *mass* of a nucleus is also approximately proportional to the total number A of protons and neutrons. When we divide the mass by the volume to calculate the density, the factors of A cancel. Thus *all nuclei have approximately the same density*. We'll see in Section 43.2 that this observation is important in understanding nuclear structure.

CAUTION Nuclear masses are rest masses When we speak of the masses of the proton, the neutron, and nuclei, we always mean their *rest* masses (see Section 37.7). We'll use this same convention when we discuss other subatomic particles in Chapter 44. 

EXAMPLE 43.1 Calculating nuclear properties

The most common kind of iron nucleus has mass number $A = 56$. Find the approximate radius, mass, and density of the nucleus.

IDENTIFY and SET UP Equation (43.1) tells us how the nuclear radius R depends on the mass number A . The mass of the nucleus in atomic mass units is approximately equal to the value of A , and the density ρ is mass divided by volume.

EXECUTE The radius and mass are

$$\begin{aligned} R &= R_0 A^{1/3} = (1.2 \times 10^{-15} \text{ m})(56)^{1/3} \\ &= 4.6 \times 10^{-15} \text{ m} = 4.6 \text{ fm} \\ m &\approx (56 \text{ u})(1.66 \times 10^{-27} \text{ kg/u}) = 9.3 \times 10^{-26} \text{ kg} \end{aligned}$$

The volume V of the nucleus (which we treat as a sphere of radius R) and its density ρ are

$$\begin{aligned} V &= \frac{4}{3}\pi R^3 = \frac{4}{3}\pi R_0^3 A = \frac{4}{3}\pi (4.6 \times 10^{-15} \text{ m})^3 \\ &= 4.1 \times 10^{-43} \text{ m}^3 \\ \rho &= \frac{m}{V} \approx \frac{9.3 \times 10^{-26} \text{ kg}}{4.1 \times 10^{-43} \text{ m}^3} = 2.3 \times 10^{17} \text{ kg/m}^3 \end{aligned}$$

EVALUATE As we mentioned above, *all* nuclei have approximately this same density. The density of solid iron is about 7000 kg/m^3 ; the iron nucleus is more than 10^{13} times as dense as iron in bulk. Such densities are also found in *neutron stars*, which are similar to gigantic nuclei made almost entirely of neutrons. A 1 cm cube of material with this density would have a mass of $2.3 \times 10^{11} \text{ kg}$, or 230 million metric tons!

KEYCONCEPT The radius of a nucleus is approximately proportional to the $\frac{1}{3}$ power of the number of nucleons A that the nucleus contains. The volume is approximately proportional to A .

Nuclides and Isotopes

The building blocks of the nucleus are the proton and the neutron. In a neutral atom, there is one electron for every proton in the nucleus. We introduced these particles in Section 21.1; we'll recount the discovery of the neutron and proton in Chapter 44. The masses of these particles are

$$\begin{aligned} \text{Proton:} \quad m_p &= 1.007276 \text{ u} = 1.672622 \times 10^{-27} \text{ kg} \\ \text{Neutron:} \quad m_n &= 1.008665 \text{ u} = 1.674927 \times 10^{-27} \text{ kg} \\ \text{Electron:} \quad m_e &= 0.000548580 \text{ u} = 9.10938 \times 10^{-31} \text{ kg} \end{aligned}$$


The number of protons in a nucleus is the **atomic number** Z . The number of neutrons is the **neutron number** N . The nucleon number or mass number A is the sum of the number of protons Z and the number of neutrons N :

$$A = Z + N \quad (43.2)$$

A single nuclear species having specific values of both Z and N is called a **nuclide**. **Table 43.1** (next page) lists values of A , Z , and N for some nuclides. The electron structure of an atom, which is responsible for its chemical properties, is determined by the charge Ze of the nucleus. The table shows some nuclides that have the same number of protons Z

APPLICATION Using Isotopes to Measure Ancient Climate

This sample of ice from Antarctica was deposited tens of thousands of years ago. The deeper the sample, the further in the past the ice was deposited. Most of the water molecules (H_2O) in the ice contain the oxygen isotope ^{16}O , but a small percentage contain the heavier isotope ^{18}O . Water molecules that contain the lighter isotope evaporate more readily, but condense less readily, than water molecules that include the heavier isotope, and these processes vary with temperature. Measuring the ratio of ^{18}O to ^{16}O in an ancient ice sample thus allows scientists to determine the average ocean temperature at the time the sample was deposited. Scientists also measure the amount of atmospheric carbon dioxide (CO_2) that was trapped in the ice when it was deposited. These observations have helped confirm the idea that high atmospheric CO_2 concentrations go hand in hand with high temperatures, a key principle for understanding 21st-century climate change (see Section 17.7).

**TABLE 43.1** Compositions of Some Common Nuclides Z = atomic number (number of protons) N = neutron number $A = Z + N$ = mass number (total number of nucleons)

Nucleus	Z	N	$A = Z + N$
^1_1H	1	0	1
^2_1H	1	1	2
^4_2He	2	2	4
^6_3Li	3	3	6
^7_3Li	3	4	7
^9_4Be	4	5	9
$^{10}_5\text{B}$	5	5	10
$^{11}_5\text{B}$	5	6	11
$^{12}_6\text{C}$	6	6	12
$^{13}_6\text{C}$	6	7	13
$^{14}_7\text{N}$	7	7	14
$^{16}_8\text{O}$	8	8	16
$^{23}_{11}\text{Na}$	11	12	23
$^{65}_{29}\text{Cu}$	29	36	65
$^{200}_{80}\text{Hg}$	80	120	200
$^{235}_{92}\text{U}$	92	143	235
$^{238}_{92}\text{U}$	92	146	238

but a different number of neutrons N . These nuclides are called **isotopes** of that element. A familiar example is chlorine (Cl , $Z = 17$). About 76% of chlorine nuclei have $N = 18$; the other 24% have $N = 20$. Different isotopes of an element usually have slightly different physical properties such as melting and boiling temperatures and diffusion rates. The two common isotopes of uranium with $A = 235$ and 238 are usually separated industrially by taking advantage of the different diffusion rates of gaseous uranium hexafluoride (UF_6) containing the two isotopes.

Table 43.1 also shows the usual notation for individual nuclides: the symbol of the element, with a pre-subscript equal to Z and a pre-superscript equal to the mass number A . The general format for an element El is ^A_ZEl . The isotopes of chlorine mentioned above, with $A = 35$ and 37 , are written $^{35}_{17}\text{Cl}$ and $^{37}_{17}\text{Cl}$ and pronounced “chlorine-35” and “chlorine-37,” respectively. This name of the element determines the atomic number Z , so the pre-subscript Z is sometimes omitted, as in ^{35}Cl .

Table 43.2 gives the masses of some common atoms, including their electrons. Note that this table gives masses of *neutral* atoms (with Z electrons) rather than masses of *bare* nuclei, because it is much more difficult to measure masses of bare nuclei with high precision. The mass of a neutral carbon-12 atom is exactly 12 u; that’s how the unified atomic mass unit is defined. The masses of other atoms are *approximately* equal to A atomic mass units, as we stated earlier. In fact, the atomic masses are *less* than the sum of the masses of their parts (the Z protons, the Z electrons, and the N neutrons). We’ll explain this very important mass difference in the next section.

Nuclear Spins and Magnetic Moments

Like electrons, nucleons (protons and neutrons) are spin- $\frac{1}{2}$ particles with spin angular momenta given by the same equations as in Section 41.5. The magnitude of the spin angular momentum $\vec{S}$ of a nucleon is

$$S = \sqrt{\frac{1}{2}(\frac{1}{2} + 1)}\hbar = \sqrt{\frac{3}{4}}\hbar \quad (43.3)$$

TABLE 43.2 Neutral Atomic Masses for Some Light Nuclides

Element and Isotope	Atomic Number, Z	Neutron Number, N	Atomic Mass (u)	Mass Number, A
Hydrogen (^1_1H)	1	0	1.007825	1
Deuterium (^2_1H)	1	1	2.014102	2
Tritium (^3_1H)	1	2	3.016049	3
Helium (^3_2He)	2	1	3.016029	3
Helium (^4_2He)	2	2	4.002603	4
Lithium (^6_3Li)	3	3	6.015123	6
Lithium (^7_3Li)	3	4	7.016003	7
Beryllium (^9_4Be)	4	5	9.012183	9
Boron ($^{10}_5\text{B}$)	5	5	10.012937	10
Boron ($^{11}_5\text{B}$)	5	6	11.009305	11
Carbon ($^{12}_6\text{C}$)	6	6	12.000000	12
Carbon ($^{13}_6\text{C}$)	6	7	13.003355	13
Nitrogen ($^{14}_7\text{N}$)	7	7	14.003074	14
Nitrogen ($^{15}_7\text{N}$)	7	8	15.000109	15
Oxygen ($^{16}_8\text{O}$)	8	8	15.994915	16
Oxygen ($^{17}_8\text{O}$)	8	9	16.999132	17
Oxygen ($^{18}_8\text{O}$)	8	10	17.999160	18

Source: International Atomic Energy Agency.

and the z -component is

$$S_z = \pm \frac{1}{2} \hbar \quad (43.4)$$

In addition to its spin angular momentum, a nucleon may have *orbital* angular momentum $\vec{L}$ associated with its motion within the nucleus. The values of $\vec{L}$ and of its z -component L_z for a nucleon are quantized in the same way as for an electron in an atom.

The *total* angular momentum $\vec{J}$ of the nucleus is the vector sum of the individual spin and orbital angular momenta of all the nucleons. It has magnitude

$$J = \sqrt{j(j+1)} \hbar \quad (43.5)$$

and z -component

$$J_z = m_j \hbar \quad (m_j = -j, -j+1, \dots, j-1, j) \quad (43.6)$$

The total nuclear angular momentum quantum number j is usually called the *nuclear spin*, even though in general it refers to a combination of the orbital and spin angular momenta of the nucleons that make up the nucleus. When the total number of nucleons A is *even*, j is an integer; when it is *odd*, j is a half-integer. All nuclides for which both Z and N are even have $J = 0$. As we'll see, this happens because nucleons tend to form pairs with opposite spin components.

Associated with nuclear angular momentum is a *magnetic moment*. When we discussed *electron* magnetic moments in Section 41.4, we introduced the Bohr magneton $\mu_B = e\hbar/2m_e$ as a natural unit of magnetic moment. We found that the magnitude of the z -component of the electron spin magnetic moment is almost exactly equal to μ_B ; that is, $|\mu_{sz}|_{\text{electron}} \approx \mu_B$. In discussing *nuclear* magnetic moments, we can define an analogous quantity, the **nuclear magneton** μ_n :

$$\mu_n = \frac{e\hbar}{2m_p} = 5.05078 \times 10^{-27} \text{ J/T} = 3.15245 \times 10^{-8} \text{ eV/T} \quad (43.7)$$

(nuclear magneton)

The proton mass m_p is 1836 times larger than the electron mass m_e , so the nuclear magneton μ_n is 1836 times smaller than the Bohr magneton μ_B .

We might expect the magnitude of the z -component of the spin magnetic moment of the proton to be approximately μ_n . Instead, it turns out to be

$$|\mu_{sz}|_{\text{proton}} = 2.7928\mu_n \quad (43.8)$$

Even more surprising, the neutron, which has zero charge, has a spin magnetic moment; its z -component has magnitude

$$|\mu_{sz}|_{\text{neutron}} = 1.9130\mu_n \quad (43.9)$$

The proton has a positive charge; as expected, its spin magnetic moment $\vec{\mu}$ is parallel to its spin angular momentum $\vec{S}$. However, $\vec{\mu}$ and $\vec{S}$ are opposite for a neutron, as would be expected for a *negative* charge distribution. These *anomalous* magnetic moments arise because the proton and neutron aren't really fundamental particles but are made of simpler particles called *quarks*. We'll discuss quarks in some detail in Chapter 44.

The magnetic moment of an entire nucleus is typically a few nuclear magnetons. When a nucleus is placed in an external magnetic field $\vec{B}$, there is an interaction energy $U = -\vec{\mu} \cdot \vec{B} = -\mu_z B$ just as with atomic magnetic moments. The components of the magnetic moment in the direction of the field μ_z are quantized, so a series of energy levels results from this interaction.

EXAMPLE 43.2 Proton spin flips

Protons are placed in a 2.30 T magnetic field that points in the positive z -direction. (a) What is the energy difference between states with the z -component of proton spin angular momentum parallel and antiparallel to the field? (b) A proton can make a transition from one of these states to the other by emitting or absorbing a photon with the appropriate energy. Find the frequency and wavelength of such a photon.

IDENTIFY and SET UP The proton is a spin- $\frac{1}{2}$ particle with a magnetic moment $\vec{\mu}$ in the same direction as its spin $\vec{S}$, so its energy depends on the orientation of its spin relative to an applied magnetic field $\vec{B}$. If the z -component of $\vec{S}$ is aligned with $\vec{B}$, then μ_z is equal to the positive value given in Eq. (43.8). If the z -component of $\vec{S}$ is opposite $\vec{B}$, then μ_z is the negative of this value. The interaction energy in either case is $U = -\mu_z B$; the difference between these energies is our target variable in part (a). We find the photon frequency and wavelength by using $E = hf = hc/\lambda$.

EXECUTE (a) When the z -components of $\vec{S}$ and $\vec{\mu}$ are parallel to $\vec{B}$, the interaction energy is

$$\begin{aligned} U &= -|\mu_z|B = -(2.7928)(3.152 \times 10^{-8} \text{ eV/T})(2.30 \text{ T}) \\ &= -2.025 \times 10^{-7} \text{ eV} \end{aligned}$$

When the z -components of $\vec{S}$ and $\vec{\mu}$ are antiparallel to the field, the energy is $+2.025 \times 10^{-7} \text{ eV}$. Hence the energy *difference* between the states is

$$\Delta E = 2(2.025 \times 10^{-7} \text{ eV}) = 4.05 \times 10^{-7} \text{ eV}$$

(b) The corresponding photon frequency and wavelength are

$$\begin{aligned} f &= \frac{\Delta E}{h} = \frac{4.05 \times 10^{-7} \text{ eV}}{4.136 \times 10^{-15} \text{ eV} \cdot \text{s}} = 9.79 \times 10^7 \text{ Hz} = 97.9 \text{ MHz} \\ \lambda &= \frac{c}{f} = \frac{3.00 \times 10^8 \text{ m/s}}{9.79 \times 10^7 \text{ s}^{-1}} = 3.06 \text{ m} \end{aligned}$$

EVALUATE This frequency is in the middle of the FM radio band. When a hydrogen specimen is placed in a 2.30 T magnetic field and irradiated with radio waves of this frequency, proton *spin flips* can be detected by the absorption of energy from the radiation.


KEYCONCEPT Both the proton and the neutron have a magnetic moment $\vec{\mu}$ associated with its spin. When placed in a magnetic field $\vec{B}$, a proton or neutron has two states of different energy: one where the component of $\vec{\mu}$ in the direction of $\vec{B}$ is positive and one where it is negative.

Nuclear Magnetic Resonance and MRI

Spin-flip experiments of the sort referred to in Example 43.2 are called *nuclear magnetic resonance* (NMR). They have been carried out with many different nuclides. Frequencies and magnetic fields can be measured very precisely, so this technique permits precise measurements of nuclear magnetic moments. An elaboration of this basic idea leads to *magnetic resonance imaging* (MRI), a noninvasive medical imaging technique that discriminates among various types of body tissues on the basis of the differing environments of protons in the tissues (**Fig. 43.1**).

The magnetic moment of a nucleus is also the *source* of a magnetic field. In an atom the interaction of an electron's magnetic moment with the field of the nucleus's magnetic

Figure 43.1 Magnetic resonance imaging (MRI).



moment causes additional splittings in atomic energy levels and spectra. We called this effect *hyperfine structure* in Section 41.5. Measurements of the hyperfine structure may be used to directly determine the nuclear spin.

TEST YOUR UNDERSTANDING OF SECTION 43.1 (a) By what factor must the mass number of a nucleus increase to double its volume? (i) $\sqrt[3]{2}$; (ii) $\sqrt{2}$; (iii) 2; (iv) 4; (v) 8. (b) By what factor must the mass number increase to double the radius of the nucleus? (i) $\sqrt[3]{2}$; (ii) $\sqrt{2}$; (iii) 2; (iv) 4; (v) 8.

ANSWER


(a) (iii), (b) (v) The radius R is proportional to the cube root of the mass number A , while the volume is proportional to R^3 and hence to $(A^{1/3})^3 = A$. Therefore, doubling the volume requires increasing the mass number by a factor of 2; doubling the radius implies increasing both the volume and the mass number by a factor of $2^3 = 8$.

43.2 NUCLEAR BINDING AND NUCLEAR STRUCTURE

Because energy must be added to a nucleus to separate it into its individual protons and neutrons, the total rest energy E_0 of the separated nucleons is greater than the rest energy of the nucleus. The minimum energy that must be added to separate the nucleons is called the **binding energy** E_B ; it is the magnitude of the energy by which the nucleons are bound together. Thus the rest energy of the nucleus is $E_0 - E_B$. The binding energy is defined as

BIO APPLICATION Deuterium and Heavy Water Toxicity

A crucial step in plant and animal cell division is the formation of a spindle, which separates the two sets of daughter chromosomes. If a plant is given only heavy water—in which one or both of the hydrogen atoms in an H_2O molecule are replaced with a deuterium atom—cell division stops and the plant stops growing. The reason is that deuterium is more massive than ordinary hydrogen, so the O–H bond in heavy water has a slightly different binding energy and heavy water has slightly different properties as a solvent. The biochemical reactions that occur during cell division are very sensitive to these solvent properties, so a spindle never forms and the cell cannot reproduce.



$$E_B = (ZM_H + Nm_n - \frac{A}{Z}M)c^2 \quad (43.10)$$

Labels for Equation (43.10):

- Binding energy of a nucleus with Z protons, N neutrons:** E_B
- Atomic number:** Z
- Mass of hydrogen atom:** M_H
- Neutron number:** N
- Neutron mass:** m_n
- (Speed of light in vacuum)² = 931.5 MeV/u:** c^2
- Mass of neutral atom containing nucleus:** $\frac{A}{Z}M$

Note that Eq. (43.10) does not include Zm_p , the mass of Z protons. Rather, it contains ZM_H , the mass of Z protons and Z electrons combined as Z neutral ${}^1_1\text{H}$ atoms, to balance the Z electrons included in $\frac{A}{Z}M$, the mass of the neutral atom.

The simplest nucleus is that of hydrogen, a single proton. Next comes ${}^2_1\text{H}$, the isotope of hydrogen with mass number 2, usually called *deuterium*. Its nucleus consists of a proton and a neutron bound together to form a particle called the *deuteron*. By using values from Table 43.2 in Eq. (43.10), we find that the binding energy of the deuteron is

$$\begin{aligned} E_B &= (1.007825 \text{ u} + 1.008665 \text{ u} - 2.014102 \text{ u})(931.5 \text{ MeV/u}) \\ &= 2.224 \text{ MeV} \end{aligned}$$

This much energy would be required to pull the deuteron apart into a proton and a neutron. An important measure of how tightly a nucleus is bound is the *binding energy per nucleon*, E_B/A . At $(2.224 \text{ MeV})/(2 \text{ nucleons}) = 1.112 \text{ MeV per nucleon}$, ${}^2_1\text{H}$ has the lowest binding energy per nucleon of all nuclides. The nuclide with the highest value of E_B/A is ${}^{62}_{28}\text{Ni}$ (see Example 43.3).

Using the equivalence of rest mass and energy (Section 37.8), we see that the mass of a nucleus is always *less* than the total mass of its nucleons by an amount $\Delta M = E_B/c^2$, called the *mass defect*. For example, the mass defect of ${}^2_1\text{H}$ is $\Delta M = E_B/c^2 = (2.224 \text{ MeV})/(931.5 \text{ MeV/u}) = 0.002388 \text{ u}$.

PROBLEM-SOLVING STRATEGY 43.1 Nuclear Properties

IDENTIFY *the relevant concepts:* The key properties of a nucleus are its mass, radius, binding energy, mass defect, binding energy per nucleon, and angular momentum.

SET UP *the problem:* Once you have identified the target variables, assemble the equations needed to solve the problem from this section and Section 43.1.

EXECUTE *the solution:* Solve for the target variables. Binding-energy calculations that use Eq. (43.10) often involve subtracting two nearly equal quantities. To get enough precision in the difference, you may need to carry as many as nine significant figures.

EVALUATE *your answer:* It's useful to be familiar with the following benchmark magnitudes. Protons and neutrons are about 1840 times as massive as electrons. Nuclear radii are of the order of 10^{-15} m . The electric potential energy of two protons in a nucleus is roughly 10^{-13} J or 1 MeV, so nuclear interaction energies are typically a few MeV rather than a few eV as with atoms. The binding energy per nucleon is about 1% of the nucleon rest energy. (The ionization energy of the hydrogen atom is only 0.003% of the electron's rest energy.) Angular momenta are determined only by the value of $\hbar$, so they are of the same order of magnitude in both nuclei and atoms. Nuclear magnetic moments, however, are about a factor of 1000 *smaller* than those of electrons in atoms because nuclei are so much more massive than electrons.

EXAMPLE 43.3 The most strongly bound nuclide**WITH VARIATION PROBLEMS**

Find the mass defect, total binding energy, and binding energy per nucleon of ${}^{62}_{28}\text{Ni}$, which has neutral atomic mass 61.928345 u.

IDENTIFY and SET UP The mass defect ΔM is the difference between the mass of the nucleus and the combined mass of its constituent nucleons. The binding energy E_B is this quantity multiplied by c^2 , and the binding energy per nucleon is E_B divided by the mass number A . We use Eq. (43.10), $\Delta M = ZM_H + Nm_n - \frac{A}{Z}M$, to determine both the mass defect and the binding energy.

EXECUTE With $Z = 28$, $M_H = 1.007825 \text{ u}$, $N = A - Z = 62 - 28 = 34$, $m_n = 1.008665 \text{ u}$, and $\frac{A}{Z}M = 61.928345 \text{ u}$, Eq. (43.10) gives $\Delta M = 0.585365 \text{ u}$. The binding energy is then

$$E_B = (0.585365 \text{ u})(931.5 \text{ MeV/u}) = 545.3 \text{ MeV}$$

The binding energy *per nucleon* is $E_B/A = (545.3 \text{ MeV})/62$, or 8.795 MeV per nucleon. This value is the highest of any nuclide (**Fig. 43.2**).

EVALUATE Our result means that it would take a minimum of 545.3 MeV to pull a ${}^{62}_{28}\text{Ni}$ completely apart into 28 protons and 34 neutrons. The mass defect of ${}^{62}_{28}\text{Ni}$ is about 1% of the atomic (or the nuclear) mass. The binding energy is therefore about 1% of the rest energy of the nucleus, and the binding energy per nucleon is about 1% of the rest energy of a nucleon. Note that the mass defect is more than half the mass of a nucleon, which suggests how tightly bound nuclei are.

KEYCONCEPT The mass defect of a nucleus is the difference between the sum of the masses of its nucleons and the mass of the nucleus itself. We calculate this using the mass of a hydrogen atom rather than the mass of a proton, and using the mass of a neutral atom that contains the nucleus rather than the nuclear mass. To find the binding energy, multiply the mass defect by c^2 ; to then find the binding energy per nucleon, divide by the nucleon number A .

Figure 43.2 Approximate binding energy per nucleon as a function of mass number A (the total number of nucleons) for stable nuclides.




Figure 43.2 is a graph of binding energy per nucleon as a function of the mass number A . This graph shows that nearly all stable nuclides have binding energies per nucleon in the range 7–9 MeV. Note the spike at $A = 4$, showing the unusually large binding energy per nucleon of the ${}^4_2\text{He}$ nucleus (alpha particle) relative to its neighbors. To explain this curve, we must consider the interactions among the nucleons.

The Nuclear Force

The force that binds protons and neutrons together in the nucleus, despite the electrical repulsion of the protons, is an example of the *strong interaction* that we mentioned in Section 5.5. In the context of nuclear structure, this interaction is called the *nuclear force*. Here are some of its characteristics:

1. The nuclear force does not depend on charge; the binding is the same for both neutrons and protons. It also has a short range, of the order of nuclear dimensions—that is, 10^{-15} m. (If it had a longer range, a nucleus would grow by pulling in additional protons and neutrons.) Within its range, the attractive nuclear force is much stronger than the repulsive electric forces between protons; otherwise, the nucleus could never be stable. Unfortunately, there is no simple equation like Coulomb's law for how the nuclear force depends on the distance between nucleons.
2. The observation that nuclei all have about the same density and about the same binding energy per nucleon, E_B/A , shows that a particular nucleon cannot interact simultaneously through the nuclear force with *all* the other nucleons in a nucleus. (If it could, then in larger nuclei the nucleons would crowd closer together, making the nucleus denser and increasing the value of E_B/A .) Instead, a nucleon interacts with only the few other nucleons in its immediate vicinity. (This is different from electric forces; *every* proton in the nucleus exerts an electric repulsion on *every* other proton.) This limit on the number of interactions via the nuclear force is called *saturation*.
3. The nuclear force favors the binding of *pairs* of protons or neutrons with opposite spins and of *pairs of pairs*—that is, a pair of protons and a pair of neutrons, with each pair having opposite spins. This explains the spike in binding energy per nucleon shown in Fig. 43.2 for ${}^4_2\text{He}$ (two protons and two neutrons). We'll see other evidence for pairing effects in nuclei in the next subsection. (In Section 42.8 we described an analogous pairing that binds opposite-spin electrons in a superconductor.)

Analyzing nuclear structure is a challenge because the nature of the nuclear force is so complex and because electric forces are also involved. Even so, we can gain some insight into nuclear structure by the use of simple models. We'll discuss briefly two rather different but successful models, the *liquid-drop model* and the *shell model*.

The Liquid-Drop Model

The **liquid-drop model**, first proposed in 1928 by the Russian physicist George Gamow and later expanded by Niels Bohr, is suggested by the observation that all nuclei have nearly the same density. The individual nucleons are analogous to molecules of a liquid, held together by short-range interactions and surface-tension effects. We can use this simple picture to derive a formula for the estimated total binding energy of a nucleus. We'll include five contributions:

1. An individual nucleon interacts with only a few of its nearest neighbors (saturation). If every nucleon in a nucleus were surrounded by nearest neighbors, the binding energy per nucleon, E_B/A , would have the same constant value for all nucleons. If we call this constant C_1 , then the binding energy of a nucleus with A nucleons would be C_1A .
2. The nucleons on the surface of the nucleus have no neighbors outside the surface and so are less tightly bound than those in the interior. This decrease in the binding energy gives a *negative* energy term proportional to the surface area $4\pi R^2$. Because R is proportional to $A^{1/3}$, this term is proportional to $A^{2/3}$; we write it as $-C_2A^{2/3}$, where C_2 is another constant.
3. Every one of the Z protons repels every one of the $(Z - 1)$ other protons. The total repulsive electric potential energy is proportional to $Z(Z - 1)$ and inversely proportional to the radius R and thus to $A^{1/3}$. This energy term is negative because the nucleons are less tightly bound than they would be without the electrical repulsion. We write this correction as $-C_3Z(Z - 1)/A^{1/3}$.
4. Observations show that nuclei are most tightly bound if N is close to Z for small A and N is greater than Z (but not too much greater) for larger A . We need a negative energy term corresponding to the difference $|N - Z|$. The best agreement with observed binding energies is obtained if this term is proportional to $(N - Z)^2/A$. If we use $N = A - Z$ to express this energy in terms of A and Z , this correction is $-C_4(A - 2Z)^2/A$.
5. Finally, the nuclear force favors *pairing* of protons and of neutrons. This energy term is positive (more binding) if both Z and N are even, negative (less binding) if both Z and N are odd, and zero otherwise. The best fit to the data occurs with the form $\pm C_5A^{-4/3}$ for this term.

The total estimated binding energy E_B is the sum of these five terms:

$$E_B = C_1A - C_2A^{2/3} - C_3\frac{Z(Z - 1)}{A^{1/3}} - C_4\frac{(A - 2Z)^2}{A} \pm C_5A^{-4/3} \quad (43.11)$$

(nuclear binding energy)

The constants C_1 , C_2 , C_3 , C_4 , and C_5 , chosen to make this formula best fit the observed binding energies of nuclides, are

$$C_1 = 15.75 \text{ MeV}$$

$$C_2 = 17.80 \text{ MeV}$$

$$C_3 = 0.7100 \text{ MeV}$$

$$C_4 = 23.69 \text{ MeV}$$

$$C_5 = 39 \text{ MeV}$$

The constant C_1 is the binding energy per nucleon due to the saturated nuclear force. This energy is almost 16 MeV per nucleon, about double the *total* binding energy per nucleon in most nuclides. The other terms in Eq. (43.11) are responsible for the difference. The negative terms with C_3 and C_4 become more important for larger nuclides that have more protons and have A greater than $2Z$; that explains why the binding energy per nucleon gradually decreases beyond $A = 62$, as Fig. 43.2 shows.

If we use Eq. (43.11) to estimate the binding energy E_B , we can solve Eq. (43.10) to use it to estimate the mass of any neutral atom:

$${}^A_ZM = ZM_H + Nm_n - \frac{E_B}{c^2} \quad (\text{semiempirical mass formula}) \quad (43.12)$$

Equation (43.12) is called the *semiempirical mass formula*. The name is apt; the equation is *empirical* in the sense that the C 's have to be determined empirically (experimentally), yet it does have a sound theoretical basis.

EXAMPLE 43.4 Estimating binding energy and mass

WITH VARIATION PROBLEMS

For the nuclide ${}^{62}_{28}\text{Ni}$ of Example 43.3, (a) calculate the five terms in the binding energy and the total estimated binding energy, and (b) find the neutral atomic mass using the semiempirical mass formula.

IDENTIFY and SET UP We use the liquid-drop model of the nucleus and its five contributions to the binding energy, as given by Eq. (43.11), to calculate the total binding energy E_B . We then use Eq. (43.12) to find the neutral atomic mass ${}^{62}_{28}M$.

EXECUTE (a) With $Z = 28$, $A = 62$, and $N = 34$, the five terms in Eq. (43.11) are

1. $C_1A = (15.75 \text{ MeV})(62) = 976.5 \text{ MeV}$
2. $-C_2A^{2/3} = -(17.80 \text{ MeV})(62)^{2/3} = -278.8 \text{ MeV}$
3. $-C_3\frac{Z(Z-1)}{A^{1/3}} = -(0.7100 \text{ MeV})\frac{(28)(27)}{(62)^{1/3}} = -135.6 \text{ MeV}$
4. $-C_4\frac{(A-2Z)^2}{A} = -(23.69 \text{ MeV})\frac{(62-56)^2}{62} = -13.8 \text{ MeV}$
5. $+C_5A^{-4/3} = (39 \text{ MeV})(62)^{-4/3} = 0.2 \text{ MeV}$

The pairing correction (term 5) is by far the smallest of all the terms; it is positive because both Z and N are even. The sum of all five terms is the total estimated binding energy, $E_B = 548.5 \text{ MeV}$.

(b) We use $E_B = 548.5 \text{ MeV}$ in Eq. (43.12):

$$\begin{aligned} {}^{62}_{28}M &= 28(1.007825 \text{ u}) + 34(1.008665 \text{ u}) - \frac{548.5 \text{ MeV}}{931.5 \text{ MeV/u}} \\ &= 61.925 \text{ u} \end{aligned}$$

EVALUATE The binding energy of ${}^{62}_{28}\text{Ni}$ calculated in part (a) is only about 0.6% larger than the true value of 545.3 MeV found in Example 43.3, and the mass calculated in part (b) is only about 0.005% smaller than the measured value of 61.928345 u. The semiempirical mass formula can be quite accurate!

KEYCONCEPT You can use the liquid-drop model and its associated semiempirical mass formula to calculate the binding energy and mass of a nuclide.

The liquid-drop model and the mass formula derived from it are quite successful in correlating nuclear masses, and we'll see later that they help in understanding decay processes of unstable nuclides. Other aspects of nuclei, such as angular momentum and excited states, are better approached with different models.

The Shell Model


The **shell model** of nuclear structure is analogous to the central-field approximation in atomic physics (see Section 41.6). We picture each nucleon as moving in a potential that represents the averaged-out effect of all the other nucleons. Although this is a very simplified model, in several respects it works out quite well.

The potential-energy function for the nuclear force is the same for protons as for neutrons. **Figure 43.3a** (next page) shows a reasonable assumption for the shape of this function: a spherical version of the square-well potential we discussed in Section 40.3. The function is somewhat rounded because the nucleus doesn't have a sharply defined surface at radius R . For a proton there is an additional electric potential energy due to interactions with the other $Z - 1$ protons, which we collectively treat as a uniformly charged sphere of radius R and total charge $(Z - 1)e$. **Figure 43.3b** shows the nuclear, electric, and total potential energies for a proton as functions of the distance r from the center of the nucleus.


In principle, we could solve the Schrödinger equation for a proton or neutron moving in such a potential. For any spherically symmetric potential energy, the angular-momentum states are the same as for the electrons in the central-field approximation in atomic physics.

Figure 43.3 Approximate potential-energy functions for a nucleon in a nucleus. The approximate nuclear radius is R .

(a) The potential energy U_{nuc} due to the nuclear force is the same for protons and neutrons. For neutrons, it is the *total* potential energy.



(b) For protons, the total potential energy U_{tot} is the sum of the nuclear (U_{nuc}) and electric (U_{el}) potential energies.



In particular, we can use the concept of *filled shells and subshells* and their relationship to stability. As we saw in Section 41.6, the exclusion principle forbids more than one electron from occupying any given quantum-mechanical state in a multielectron atom. This explains why the values $Z = 2, 10, 18, 36, 54,$ and 86 (the atomic numbers of the noble gases) correspond to atoms with particularly stable electron arrangements.

A comparable effect occurs in nuclear structure. Like electrons, protons and neutrons are spin- $\frac{1}{2}$ particles. So no more than one proton can be in any given quantum-mechanical state in a nucleus, and likewise for neutrons. Just as for electrons in atoms, there are certain numbers of protons *or* of neutrons, called *magic numbers*, that correspond to particularly stable nuclei—that is, nuclei with particularly high binding energies. The magic numbers are 2, 8, 20, 28, 50, 82, and 126. These numbers are different from those for electrons in atoms because the potential-energy function is different and the nuclear spin-orbit interaction is much stronger and of opposite sign than in atoms. So nuclear subshells fill up in a different order from those for electrons in an atom. Nuclides in which Z is a magic number tend to have an above-average number of stable isotopes. (Nuclides with $Z = 126$ have not been observed in nature.) There are several *doubly magic* nuclides for which both Z and N are magic, including



All these nuclides have substantially higher binding energy per nucleon than do nuclides with neighboring values of N or Z . They also all have zero nuclear spin. The magic numbers correspond to filled-shell or filled-subshell configurations of nucleon energy levels with a relatively large jump in energy to the next allowed level.

TEST YOUR UNDERSTANDING OF SECTION 43.2 Rank the following nuclei in order from largest to smallest value of the binding energy per nucleon. (i) ${}^4_2\text{He}$; (ii) ${}^{52}_{24}\text{Cr}$; (iii) ${}^{152}_{62}\text{Sm}$; (iv) ${}^{200}_{80}\text{Hg}$; (v) ${}^{252}_{92}\text{Cf}$.

ANSWER

(i), (ii), (iii), (iv), (v). You can find the answers by inspecting Fig. 43.2. The binding energy per nucleon is lowest for very light nuclei such as ${}^4_2\text{He}$, is greatest around $A = 60$, and then decreases with increasing A .

43.3 NUCLEAR STABILITY AND RADIOACTIVITY

Among about 2500 known nuclides, fewer than 300 are stable. The others are unstable structures that decay to form other nuclides by emitting particles and electromagnetic radiation, a process called **radioactivity**. The time scale of these decay processes ranges from a small fraction of a microsecond to billions of years. The *stable* nuclides are shown by dots on the graph in **Fig. 43.4**, where the neutron number N and proton number (or atomic number) Z for each nuclide are plotted. Such a chart is called a *Segrè chart*, after its inventor, the Italian-American physicist Emilio Segrè (1905–1989).

Each blue line in Fig. 43.4 represents a specific value of the mass number $A = Z + N$. Most lines of constant A pass through only one or two stable nuclides; that is, there is usually a very narrow range of stability for a given mass number. The lines at $A = 20, 40, 60,$ and 80 are examples. In four exceptional cases ($A = 94, 124, 130,$ and 136), these lines pass through *three* stable nuclides.

Only four stable nuclides have both odd Z and odd N :



These are called *odd-odd nuclides*. The absence of other odd-odd nuclides shows the importance of pairing in adding to nuclear stability. Also, there is *no* stable nuclide with $A = 5$ or $A = 8$. The doubly magic ${}^4_2\text{He}$ nucleus, with a pair of protons and a pair of neutrons, has no interest in accepting a fifth particle into its structure. Collections of eight nucleons decay to smaller nuclides, with a ${}^8_4\text{Be}$ nucleus immediately splitting into two ${}^4_2\text{He}$ nuclei.





Figure 43.4 Segrè chart showing neutron number and proton number for stable nuclides.

The stable nuclides define a rather narrow region on the Segrè chart. For low mass numbers, the numbers of protons and neutrons are approximately equal, $N \approx Z$. The ratio N/Z increases gradually with A , up to about 1.6 at large mass numbers, because of the increasing influence of the electrical repulsion of the protons. Points to the right of the stability region represent nuclides that have too many protons relative to neutrons. In these cases, repulsion wins, and the nucleus comes apart. To the left are nuclides with too many neutrons relative to protons. In these cases the energy associated with the neutrons is out of balance with that associated with the protons, and the nuclides decay in a process that converts neutrons to protons. The graph also shows that no nuclide with $A > 209$ or $Z > 83$ is stable. A nucleus is unstable if it is too big. Note that there is no stable nuclide with $Z = 43$ (technetium) or 61 (promethium).

Nearly 90% of the 2500 known nuclides are *radioactive*; they are not stable but decay into other nuclides. Many of these radioactive nuclides occur in nature. For example, you are very slightly radioactive because of unstable nuclides such as carbon-14 ($^{14}_6\text{C}$) and potassium-40 ($^{40}_{19}\text{K}$) that are present throughout your body. The study of radioactivity began in 1896, one year after Wilhelm Röntgen discovered x rays (Section 36.6). Henri Becquerel discovered a radiation from uranium salts that seemed similar to x rays. Investigation in the following two decades by Marie and Pierre Curie, Ernest Rutherford, and many others revealed that the emissions consist of positively and negatively charged particles and neutral rays. These particles were given the names *alpha*, *beta*, and *gamma* because of their differing penetration characteristics.

Figure 43.5 Alpha decay of the unstable radium nuclide $^{226}_{88}\text{Ra}$. The alpha particles used in the Rutherford scattering experiment (Section 39.2) were emitted by this nuclide.



Alpha Decay

When unstable nuclides decay into different nuclides, they usually emit alpha (α) or beta (β) particles. An **alpha particle** is a ^4_2He nucleus, two protons and two neutrons bound together, with total spin zero. Alpha decay occurs principally with nuclei that are too large to be stable. The nucleus gets rid of the excess nucleons in the form of a ^4_2He nucleus because this is a particularly stable grouping (see Fig. 43.2). When a nucleus emits an alpha particle, its N and Z values each decrease by 2 and A decreases by 4, moving it closer to stable territory on the Segrè chart.

Figure 43.5a shows the alpha decay of radium-226 ($^{226}_{88}\text{Ra}$). Spontaneous alpha decay of this kind can occur only if energy is released in the process; this released energy goes into the kinetic energy of the emitted α particle and of the nucleus that remains, called the *daughter nucleus*. (For the decay shown in Fig. 43.5a, the daughter nucleus is radon-222, $^{222}_{86}\text{Rn}$.) The original nucleus (in this case, $^{226}_{88}\text{Ra}$) is called the *parent nucleus*. You can use mass-energy conservation to show that

alpha decay is possible whenever the mass of the original neutral atom is greater than the sum of the masses of the final neutral atom and a neutral ^4_2He atom.

In alpha decay, the α particle tunnels through a potential-energy barrier, as Fig. 43.5b shows. You may want to review the discussion of tunneling in Section 40.4.

Alpha particles are always emitted with definite kinetic energies, determined by conservation of momentum and energy in the alpha-decay process. As Fig. 43.5c shows, an α particle emitted in the decay of $^{226}_{88}\text{Ra}$ can have either of *two* possible energies, depending on the energy level of the $^{222}_{86}\text{Rn}$ daughter nucleus just after the decay. (Later in this section we'll discuss the photon-emission process shown in Fig. 43.5c.)

Alpha particles are emitted at high speeds, typically a few percent of the speed of light (see the following example). Nonetheless, because of their charge and mass, alpha particles can travel only several centimeters in air, or a few tenths or hundredths of a millimeter through solids, before they are brought to rest by collisions.

EXAMPLE 43.5 Alpha decay of radium

WITH VARIATION PROBLEMS

Show that the α -emission process $^{226}_{88}\text{Ra} \rightarrow ^{222}_{86}\text{Rn} + ^4_2\text{He}$ (Fig. 43.5a) is energetically possible, and calculate the kinetic energy of the emitted α particle. The neutral atomic masses are 226.025410 u for $^{226}_{88}\text{Ra}$, 222.017578 u for $^{222}_{86}\text{Rn}$, and 4.002603 u for ^4_2He .

IDENTIFY and SET UP Alpha emission is possible if the mass of the $^{226}_{88}\text{Ra}$ atom is greater than the sum of the atomic masses of $^{222}_{86}\text{Rn}$ and ^4_2He . The mass difference between the initial radium atom and the final

radon and helium atoms corresponds (through $E = mc^2$) to the energy E released in the decay. Because momentum is conserved as well as energy, *both* the alpha particle and the $^{222}_{86}\text{Rn}$ atom are in motion after the decay; we'll have to account for this in determining the kinetic energy of the alpha particle.

EXECUTE The difference in mass between the original nucleus and the decay products is

$$226.025410 \text{ u} - (222.017578 \text{ u} + 4.002603 \text{ u}) = +0.005229 \text{ u}$$

Since this is positive, α decay is energetically possible. The energy equivalent of this mass difference is

$$E = (0.005229 \text{ u})(931.5 \text{ MeV/u}) = 4.871 \text{ MeV}$$

In this process the $^{222}_{86}\text{Rn}$ nucleus is produced in its ground level (Fig. 43.5c). Thus we expect the decay products to emerge with total kinetic energy 4.871 MeV. Momentum is also conserved; if the parent $^{226}_{88}\text{Ra}$ nucleus is at rest, the daughter $^{222}_{86}\text{Rn}$ nucleus and the α particle have momenta of equal magnitude p but opposite direction. Kinetic energy is $K = \frac{1}{2}mv^2 = p^2/2m$: Since p is the same for the two particles, the kinetic energy divides inversely as their masses. Hence the α particle gets $222/(222 + 4)$ of the total, or 4.78 MeV.

EVALUATE Experiment shows that $^{226}_{88}\text{Ra}$ does emit α particles with a kinetic energy of 4.78 MeV. Check your results by verifying that the alpha particle and the $^{222}_{86}\text{Rn}$ nucleus produced in the decay have the same magnitude of momentum $p = mv$. You can calculate the speed v of each of the decay products from its respective kinetic energy [note that the $^{222}_{86}\text{Rn}$ nucleus gets $4/(222 + 4)$ of the 4.871 MeV released]. You'll find that the alpha particle moves at $0.0506c = 1.52 \times 10^7 \text{ m/s}$; if momentum is conserved, you should find that the $^{222}_{86}\text{Rn}$ nucleus moves $\frac{4}{222}$ as fast. Does it?

KEYCONCEPT In alpha decay, a large, unstable nucleus with Z protons and N neutrons emits an alpha particle (a ^4_2He nucleus). The daughter nucleus has $Z - 2$ protons and $N - 2$ neutrons. The combined mass of the decay products is less than the mass of the parent; the lost rest energy goes into kinetic energy of the decay products.

Beta Decay

There are three different simple types of *beta decay*: *beta-minus*, *beta-plus*, and *electron capture*. A **beta-minus particle** (β^-) is an electron. There are no electrons in the nucleus waiting to be emitted; instead, emission of a β^- involves *transformation* of a neutron into a proton, an electron, and a third particle called an *antineutrino*. In fact, if you freed a neutron from a nucleus, it would decay into a proton, an electron, and an antineutrino in an average time of about 15 minutes.

Beta particles can be identified and their speeds can be measured with techniques that are similar to the Thomson e/m experiment we described in Section 27.5. The speeds of beta particles range up to 0.9995 of the speed of light, so their motion is highly relativistic. They are emitted with a continuous spectrum of energies. This would not be possible if the only two particles were the β^- and the recoiling nucleus, since energy and momentum conservation would then require a definite speed for the β^- . Thus there must be a *third* particle involved. From conservation of charge, it must be neutral, and from conservation of angular momentum, it must be a spin- $\frac{1}{2}$ particle.

This third particle is an antineutrino, the *antiparticle* of a **neutrino**. The symbol for a neutrino is ν_e (the Greek letter nu). Both the neutrino and the antineutrino have zero charge and very small mass and therefore produce very little observable effect when passing through matter. Both evaded detection until 1953, when Frederick Reines and Clyde Cowan succeeded in observing the antineutrino directly. We now know that there are at least three varieties of neutrinos, each with its corresponding antineutrino; one is associated with beta decay and the other two are associated with the decay of two unstable particles, the muon and the tau particle. We'll discuss these particles in more detail in Chapter 44. The antineutrino that is emitted in β^- decay is denoted as $\bar{\nu}_e$. The basic process of β^- decay is

$$n \rightarrow p + \beta^- + \bar{\nu}_e \quad (43.13)$$

Beta-minus decay usually occurs with nuclides for which the neutron-to-proton ratio N/Z is too large for stability. In β^- decay, N decreases by 1, Z increases by 1, and A doesn't change. You can use mass-energy conservation to show that

beta-minus decay can occur whenever the mass of the original neutral atom is larger than that of the final atom.

EXAMPLE 43.6 Why cobalt-60 is a beta-minus emitter**WITH VARIATION PROBLEMS**

The nuclide ${}^{60}_{27}\text{Co}$, an odd-odd unstable nucleus, is used in medical and industrial applications of radiation. Show that it is unstable relative to β^- decay. The atomic masses you need are 59.933817 u for ${}^{60}_{27}\text{Co}$ and 59.930786 u for ${}^{60}_{28}\text{Ni}$.

IDENTIFY and SET UP Beta-minus decay is possible if the mass of the original neutral atom is greater than that of the final atom. We must first identify the nuclide that will result if ${}^{60}_{27}\text{Co}$ undergoes β^- decay and then compare its neutral atomic mass to that of ${}^{60}_{27}\text{Co}$.

EXECUTE In the presumed β^- decay of ${}^{60}_{27}\text{Co}$, Z increases by 1 from 27 to 28 and A remains at 60, so the final nuclide is ${}^{60}_{28}\text{Ni}$. The neutral atomic mass of ${}^{60}_{27}\text{Co}$ is greater than that of ${}^{60}_{28}\text{Ni}$ by 0.003031 u, so β^- decay *can* occur.

EVALUATE With three decay products in β^- decay—the ${}^{60}_{28}\text{Ni}$ nucleus, the electron, and the antineutrino—the energy can be shared in many different ways that are consistent with conservation of energy and momentum. It's impossible to predict precisely how the energy will be shared for the decay of a particular ${}^{60}_{27}\text{Co}$ nucleus. By contrast, in alpha decay there are just two decay products, and their energies and momenta are determined uniquely (see Example 43.5).

KEYCONCEPT In beta-minus decay, an unstable nucleus with Z protons and N neutrons decays by converting a neutron into a proton and emitting an electron and an antineutrino. The daughter nucleus has $Z + 1$ protons and $N - 1$ neutrons. This is possible only if the neutral parent atom has a greater mass than the neutral daughter atom.

We have noted that β^- decay occurs with nuclides that have too large a neutron-to-proton ratio N/Z . Nuclides for which N/Z is too *small* for stability can emit a *positron*, the electron's antiparticle, which is identical to the electron but with positive charge. (We'll discuss the positron in more detail in Chapter 44.) The basic process, called *beta-plus decay* (β^+), is

$$p \rightarrow n + \beta^+ + \nu_e \quad (43.14)$$

where β^+ is a positron and ν_e is the electron neutrino.

Beta-plus decay can occur whenever the mass of the original neutral atom is at least two electron masses larger than that of the final atom.

You can show this by using mass-energy conservation.

The third type of beta decay is *electron capture*. There are a few nuclides for which β^+ emission is not energetically possible but in which an orbital electron (usually in the innermost K shell) can combine with a proton in the nucleus to form a neutron and a neutrino. The neutron remains in the nucleus and the neutrino is emitted. The basic process is

$$p + \beta^- \rightarrow n + \nu_e \quad (43.15)$$

You can use mass-energy conservation to show that

electron capture can occur whenever the mass of the original neutral atom is larger than that of the final atom.

In all types of beta decay, A remains constant. However, in beta-plus decay and electron capture, N increases by 1 and Z decreases by 1 as the neutron-to-proton ratio increases toward a more stable value. The reaction of Eq. (43.15) also helps explain the formation of a neutron star, mentioned in Example 43.1.

CAUTION Beta decay inside and outside nuclei The beta-decay reactions given by Eqs. (43.13), (43.14), and (43.15) occur *within* a nucleus. Although the decay of a neutron outside the nucleus proceeds through the reaction of Eq. (43.13), the reaction of Eq. (43.14) is forbidden by mass-energy conservation for a proton outside the nucleus. The reaction of Eq. (43.15) can occur outside the nucleus only with the addition of some extra energy, as in a collision. **|**

EXAMPLE 43.7 Why cobalt-57 is not a beta-plus emitter**WITH VARIATION PROBLEMS**

The nuclide $^{57}_{27}\text{Co}$ is an odd-even unstable nucleus. Show that it cannot undergo β^+ decay, but that it *can* decay by electron capture. The atomic masses you need are 56.936291 u for $^{57}_{27}\text{Co}$ and 56.935394 u for $^{57}_{26}\text{Fe}$.

IDENTIFY and SET UP Beta-plus decay is possible if the mass of the original neutral atom is greater than that of the final atom plus two electron masses (0.001097 u). Electron capture is possible if the mass of the original atom is greater than that of the final atom. We must first identify the nuclide that will result if $^{57}_{27}\text{Co}$ undergoes β^+ decay or electron capture and then find the corresponding mass difference.

EXECUTE The original nuclide is $^{57}_{27}\text{Co}$. In both the presumed β^+ decay and electron capture, Z decreases by 1 from 27 to 26, and A remains at 57, so the final nuclide is $^{57}_{26}\text{Fe}$. Its mass is less than that of $^{57}_{27}\text{Co}$ by 0.000897 u, a value smaller than 0.001097 u (two electron masses),

so β^+ decay *cannot* occur. However, the mass of the original atom is greater than the mass of the final atom, so electron capture *can* occur.

EVALUATE In electron capture there are just two decay products, the final nucleus and the emitted neutrino. As in alpha decay (Example 43.5) but unlike in β^- decay (Example 43.6), the decay products of electron capture have unique energies and momenta. In Section 43.4 we'll see how to relate the probability that electron capture will occur to the *half-life* of this nuclide.

KEYCONCEPT In beta-plus decay, an unstable nucleus with Z protons and N neutrons decays by converting a proton into a neutron and emitting a positron and a neutrino. The daughter nucleus has $Z - 1$ protons and $N + 1$ neutrons. This is possible only if the neutral parent atom has a greater mass than the neutral daughter atom plus two electrons. If this is not the case, electron capture may be possible.

Gamma Decay

The energy of internal motion of a nucleus is quantized. A typical nucleus has a set of allowed energy levels, including a *ground state* (state of lowest energy) and several *excited states*. Because of the great strength of nuclear interactions, excitation energies of nuclei are typically of the order of 1 MeV, compared with a few eV for atomic energy levels. In ordinary physical and chemical transformations the nucleus always remains in its ground state. When a nucleus is placed in an excited state, either by bombardment with high-energy particles or by a radioactive transformation, it can decay to the ground state by emission of one or more photons called **gamma rays** or *gamma-ray photons*, with typical energies of 10 keV to 5 MeV. This process is called *gamma (γ) decay*. For example, alpha particles emitted from ^{226}Ra have two possible kinetic energies, either 4.784 MeV or 4.602 MeV. Including the recoil energy of the resulting ^{222}Rn nucleus, these correspond to a total released energy of 4.871 MeV or 4.685 MeV, respectively (see Fig. 43.5c). When an alpha particle with the smaller energy is emitted, the ^{222}Rn nucleus is left in an excited state. It then decays to its ground state by emitting a gamma-ray photon with energy $(4.871 - 4.685) \text{ MeV} = 0.186 \text{ MeV}$.

CAUTION Comparing α , β , and γ decays In both α and β decay, the Z value of a nucleus changes and the nucleus of one element becomes the nucleus of a different element. In γ decay, the element does *not* change; the nucleus merely goes from an excited state to a less excited state. Remember that in each of these decays, the initial unstable nucleus does not simply disappear; it's just replaced by a different one (or, in the case of γ decay, a less-excited version of the initial nucleus).

Radioactive Decay Series

When a radioactive nucleus decays, the daughter nucleus may also be unstable. In this case a *series* of successive decays occurs until a stable configuration is reached. One of the most abundant radioactive nuclides found on earth is the uranium isotope ^{238}U , which undergoes a series of 14 decays, including eight α emissions and six β^- emissions, terminating at a stable isotope of lead, ^{206}Pb (Fig. 43.6).


Radioactive decay series can be represented on a Segrè chart, as in Fig. 43.7 (next page). The neutron number N is plotted vertically, and the atomic number Z is plotted horizontally. In alpha emission, both N and Z decrease by 2. In β^- emission, N decreases by 1 and Z increases by 1. The decays can also be represented in equation form; the first two decays in the series are written as



Figure 43.6 This earthquake damage to a road was caused in part by the radioactive decay of ^{238}U in the earth's interior. These decays release energy that helps produce convection currents in the earth's interior. Such currents drive the motions of the earth's crust, including the sudden sharp motions that we call earthquakes. Other radioactive nuclides whose decays heat our planet's interior include ^{235}U , ^{232}Th , and ^{40}K .



Figure 43.7 Segrè chart showing the uranium ^{238}U decay series, terminating with the stable nuclide ^{206}Pb . The times are half-lives (discussed in the next section), given in years (y), days (d), hours (h), minutes (m), or seconds (s).



or more briefly as



In the second process, the beta decay leaves the daughter nucleus ^{234}Pa in an excited state, from which it decays to the ground state by emitting a gamma-ray photon. An excited state is denoted by an asterisk, so we can represent the γ emission as



Note the branching of the ^{238}U decay series that occurs at ^{214}Bi . This nuclide decays to ^{210}Pb by emission of an α and a β^- , which can occur in either order. Note also that the series includes unstable isotopes of several elements that also have stable isotopes, including thallium (Tl), lead (Pb), and bismuth (Bi). These unstable isotopes of these elements all have too many neutrons to be stable.

Many other decay series are known. Two of these occur in nature, one starting with the uncommon isotope ^{235}U and ending with ^{207}Pb , the other starting with thorium (^{232}Th) and ending with ^{208}Pb .

TEST YOUR UNDERSTANDING OF SECTION 43.3 A nucleus with atomic number Z and neutron number N undergoes two decay processes. The result is a nucleus with atomic number $Z - 3$ and neutron number $N - 1$. Which decay processes may have taken place? (i) Two β^- decays; (ii) two β^+ decays; (iii) two α decays; (iv) an α decay and a β^- decay; (v) an α decay and a β^+ decay.

ANSWER

(v) Two protons and two neutrons are lost in an α decay, so Z and N each decrease by 2. A β^+ decay changes a proton to a neutron, so Z decreases by 1 and N increases by 1. The net result is that Z decreases by 3 and N decreases by 1.

43.4 ACTIVITIES AND HALF-LIVES

Suppose you have a certain number of nuclei of a particular radioactive nuclide. If no more are produced, that number decreases in a simple manner as the nuclei decay. This decrease is a statistical process; there is no way to predict when any individual nucleus will decay. No change in physical or chemical environment, such as chemical reactions or heating or cooling, greatly affects most decay rates. The rate varies over an extremely wide range for different nuclides.

Radioactive Decay Rates

Let $N(t)$ be the (very large) number of radioactive nuclei in a sample at time t , and let $dN(t)$ be the (negative) change in that number during a short time interval dt . (We'll use $N(t)$ to minimize confusion with the neutron number N .) The number of decays during the interval dt is $-dN(t)$. The rate of change of $N(t)$ is the negative quantity $dN(t)/dt$; thus $-dN(t)/dt$ is called the *decay rate* or the **activity** of the specimen. The larger the number of nuclei in the specimen, the more nuclei decay during any time interval. That is, the activity is directly proportional to $N(t)$; it equals a constant λ multiplied by $N(t)$:

$$-\frac{dN(t)}{dt} = \lambda N(t) \quad (43.16)$$

The constant λ is called the **decay constant**, and it has different values for different nuclides. A large value of λ corresponds to rapid decay; a small value corresponds to slower decay. Solving Eq. (43.16) for λ shows us that λ is the ratio of the number of decays per time to the number of remaining radioactive nuclei; λ can then be interpreted as the *probability per unit time* that any individual nucleus will decay.

The situation is reminiscent of a discharging capacitor, which we studied in Section 26.4. Equation (43.16) has the same form as the negative of Eq. (26.15), with q and $1/RC$ replaced by $N(t)$ and λ . Then we can make the same substitutions in Eq. (26.16), with the initial number of nuclei $N(0) = N_0$, to find

$$N(t) = N_0 e^{-\lambda t} \quad (43.17)$$

Number of remaining nuclei at time t in sample of radioactive element
Number of nuclei at $t = 0$
Time
Decay constant


Figure 43.8 is a graph of this function.

The **half-life** $T_{1/2}$ is the time required for the number of radioactive nuclei to decrease to one-half the original number N_0 . Then half of the remaining radioactive nuclei decay during a second interval $T_{1/2}$, and so on. The numbers remaining after successive half-lives are $N_0/2$, $N_0/4$, $N_0/8$, $\dots$

To get the relationship between the half-life $T_{1/2}$ and the decay constant λ , we set $N(t)/N_0 = \frac{1}{2}$ and $t = T_{1/2}$ in Eq. (43.17), obtaining

$$\frac{1}{2} = e^{-\lambda T_{1/2}}$$

Figure 43.8 The number of nuclei in a sample of a radioactive element as a function of time. The sample's activity has an exponential decay curve with the same shape.



We take logarithms of both sides and solve for $T_{1/2}$:

$$T_{1/2} = \frac{\ln 2}{\lambda} = \frac{0.693}{\lambda} \quad (43.18)$$

The mean lifetime T_{mean} , generally called the *lifetime*, of an unstable nucleus or particle is the time for $N(t)$ to decrease by $1/e$. It is proportional to the half-life $T_{1/2}$:

$$T_{\text{mean}} = \frac{1}{\lambda} = \frac{T_{1/2}}{\ln 2} = \frac{T_{1/2}}{0.693} \quad (43.19)$$

Lifetime of unstable nucleus or particle $\rightarrow$ T_{mean} $\leftarrow$ Half-life of nucleus or particle
Decay constant of nucleus or particle $\leftarrow$ λ

In particle physics the life of an unstable particle is usually described by the lifetime, not the half-life.

Because the activity $-dN(t)/dt$ at any time equals $\lambda N(t)$, Eq. (43.17) tells us that the activity also depends on time as $e^{-\lambda t}$. Thus the graph of activity versus time has the same shape as Fig. 43.8. Also, after successive half-lives, the activity is one-half, one-fourth, one-eighth, and so on of the original activity.

CAUTION A half-life may not be enough It is sometimes implied that any radioactive sample will be safe after a half-life has passed. That's wrong. If your radioactive waste initially has ten times too much activity for safety, it is not safe after one half-life, when it still has five times too much. Even after three half-lives it still has 25% more activity than is safe. The number of radioactive nuclei and the activity approach zero only as t approaches infinity. ■

A common unit of activity is the **curie**, abbreviated Ci, which is defined to be 3.70×10^{10} decays per second. This is approximately equal to the activity of one gram of radium-226. The SI unit of activity is the *becquerel*, abbreviated Bq. One becquerel is one decay per second, so

$$1 \text{ Ci} = 3.70 \times 10^{10} \text{ Bq} = 3.70 \times 10^{10} \text{ decays/s}$$

EXAMPLE 43.8 Activity of ^{57}Co

WITH VARIATION PROBLEMS

The isotope ^{57}Co decays by electron capture to ^{57}Fe with a half-life of 272 d. The ^{57}Fe nucleus is produced in an excited state, and it almost instantaneously emits gamma rays that we can detect. (a) Find the mean lifetime and decay constant for ^{57}Co . (b) If the activity of a ^{57}Co radiation source is now $2.00 \mu\text{Ci}$, how many ^{57}Co nuclei does the source contain? (c) What will be the activity after one year?

IDENTIFY and SET UP This problem uses the relationships among decay constant λ , lifetime T_{mean} , and activity $-dN(t)/dt$. In part (a) we use Eq. (43.19) to find λ and T_{mean} from $T_{1/2}$. In part (b), we use Eq. (43.16) to calculate the number of nuclei $N(t)$ from the activity. Finally, in part (c) we use Eqs. (43.16) and (43.17) to find the activity after one year.

EXECUTE (a) It's convenient to convert the half-life from days to seconds:

$$\begin{aligned} T_{1/2} &= (272 \text{ d})(86,400 \text{ s/d}) \\ &= 2.35 \times 10^7 \text{ s} \end{aligned}$$

From Eq. (43.19), the mean lifetime and the decay constant are

$$\begin{aligned} T_{\text{mean}} &= \frac{T_{1/2}}{\ln 2} = \frac{2.35 \times 10^7 \text{ s}}{0.693} \\ &= 3.39 \times 10^7 \text{ s} = 392 \text{ days} \\ \lambda &= \frac{1}{T_{\text{mean}}} = 2.95 \times 10^{-8} \text{ s}^{-1} \end{aligned}$$

(b) The activity $-dN(t)/dt$ is given as $2.00 \mu\text{Ci}$, so

$$\begin{aligned} -\frac{dN(t)}{dt} &= 2.00 \mu\text{Ci} = (2.00 \times 10^{-6})(3.70 \times 10^{10} \text{ s}^{-1}) \\ &= 7.40 \times 10^4 \text{ decays/s} \end{aligned}$$

From Eq. (43.16) this is equal to $\lambda N(t)$, so we find

$$\begin{aligned} N(t) &= -\frac{dN(t)/dt}{\lambda} = \frac{7.40 \times 10^4 \text{ s}^{-1}}{2.95 \times 10^{-8} \text{ s}^{-1}} \\ &= 2.51 \times 10^{12} \text{ nuclei} \end{aligned}$$

If you feel we're being too cavalier about the "units" decays and nuclei, you can use decays/(nucleus · s) as the unit for λ .

(c) From Eq. (43.17) the number $N(t)$ of nuclei remaining after one year (3.156×10^7 s) is

$$\begin{aligned} N(t) &= N_0 e^{-\lambda t} = N_0 e^{-(2.95 \times 10^{-8} \text{ s}^{-1})(3.156 \times 10^7 \text{ s})} \\ &= 0.394 N_0 \end{aligned}$$

The number of nuclei has decreased to 0.394 of the original number. Equation (43.16) says that the activity is proportional to the number of nuclei, so the activity has decreased by this same factor to $(0.394)(2.00 \text{ } \mu\text{Ci}) = 0.788 \text{ } \mu\text{Ci}$.

EVALUATE The number of nuclei found in part (b) is equivalent to $4.17 \times 10^{-12} \text{ mol}$, with a mass of $2.38 \times 10^{-10} \text{ g}$. This is a far smaller mass than even the most sensitive balance can measure.

After one 272 day half-life, the number of ^{57}Co nuclei has decreased to $N_0/2$; after $2(272 \text{ d}) = 544 \text{ d}$, it has decreased to $N_0/2^2 = N_0/4$. This result agrees with our answer to part (c), which says that after 365 d the number of nuclei is between $N_0/2$ and $N_0/4$.

KEYCONCEPT Because radioactive decay is a statistical process, the number of radioactive nuclei $N(t)$ present in a radioactive sample and the activity $-dN(t)/dt$ of the sample both decay exponentially with time. Both $N(t)$ and $-dN(t)/dt$ decrease by $\frac{1}{2}$ in one half-life and by $1/e$ in one mean lifetime.

Radioactive Dating

An important application of radioactivity is the dating of archaeological and geological specimens by measuring the concentration of radioactive isotopes. The most familiar example is *carbon dating*. The unstable isotope ^{14}C , produced during nuclear reactions in the atmosphere that result from cosmic-ray bombardment, gives a small proportion of ^{14}C in the CO_2 in the atmosphere. Plants that obtain their carbon from this source contain the same proportion of ^{14}C as the atmosphere. When a plant dies, it stops taking in carbon, and its ^{14}C β^- decays to ^{14}N with a half-life of 5730 years. By measuring the proportion of ^{14}C in the remains, we can determine how long ago the organism died.

One difficulty with radiocarbon dating is that the ^{14}C concentration in the atmosphere changes over long time intervals. Corrections can be made on the basis of other data such as measurements of tree rings that show annual growth cycles. Similar radioactive techniques are used with other isotopes for dating geological specimens. Some rocks, for example, contain the unstable potassium isotope ^{40}K , a beta emitter that decays to the stable nuclide ^{40}Ar with a half-life of $2.4 \times 10^8 \text{ y}$. The age of the rock can be determined by comparing the concentrations of ^{40}K and ^{40}Ar .

EXAMPLE 43.9 Radiocarbon dating

WITH VARIATION PROBLEMS

Before 1900 the activity per unit mass of atmospheric carbon due to the presence of ^{14}C averaged about 0.255 Bq per gram of carbon. (a) What fraction of carbon atoms were ^{14}C ? (b) In analyzing an archaeological specimen containing 500 mg of carbon, you observe 174 decays in one hour. What is the age of the specimen, assuming that its activity per unit mass of carbon when it died was that average value of the air?

IDENTIFY and SET UP The key idea is that the present-day activity of a biological sample containing ^{14}C is related to both the elapsed time since it stopped taking in atmospheric carbon and its activity at that time. We use Eqs. (43.16) and (43.17) to solve for the age t of the specimen. In part (a) we determine the number of ^{14}C atoms $N(t)$ from the activity $-dN(t)/dt$ by using Eq. (43.16). We find the total number of carbon atoms in 500 mg by using the molar mass of carbon (12.011 g/mol, given in Appendix D), and we use the result to calculate the fraction of carbon atoms that are ^{14}C . The activity decays at the same rate as the number of ^{14}C nuclei; we use this and Eq. (43.17) to solve for the age t of the specimen.

EXECUTE (a) To use Eq. (43.16), we must first find the decay constant λ from Eq. (43.18):

$$\begin{aligned} T_{1/2} &= 5730 \text{ y} = (5730 \text{ y})(3.156 \times 10^7 \text{ s/y}) = 1.808 \times 10^{11} \text{ s} \\ \lambda &= \frac{\ln 2}{T_{1/2}} = \frac{0.693}{1.808 \times 10^{11} \text{ s}} = 3.83 \times 10^{-12} \text{ s}^{-1} \end{aligned}$$

Then, from Eq. (43.16),

$$N(t) = \frac{-dN/dt}{\lambda} = \frac{0.255 \text{ s}^{-1}}{3.83 \times 10^{-12} \text{ s}^{-1}} = 6.65 \times 10^{10} \text{ atoms}$$

The *total* number of C atoms in 1 gram ($1/12.011 \text{ mol}$) is $(1/12.011)(6.022 \times 10^{23}) = 5.01 \times 10^{22}$. The ratio of ^{14}C atoms to all C atoms is

$$\frac{6.65 \times 10^{10}}{5.01 \times 10^{22}} = 1.33 \times 10^{-12}$$

Only four carbon atoms in every 3×10^{12} are ^{14}C .

Continued

(b) Assuming that the activity per gram of carbon in the specimen when it died ($t = 0$) was $0.255 \text{ Bq/g} = (0.255 \text{ s}^{-1} \cdot \text{g}^{-1}) \times (3600 \text{ s/h}) = 918 \text{ h}^{-1} \cdot \text{g}^{-1}$, the activity of 500 mg of carbon then was $(0.500 \text{ g})(918 \text{ h}^{-1} \cdot \text{g}^{-1}) = 459 \text{ h}^{-1}$. The observed activity now, at time t , is 174 h^{-1} . Since the activity is proportional to the number of radioactive nuclei, the activity ratio $174/459 = 0.379$ equals the number ratio $N(t)/N_0$.

Now we solve Eq. (43.17) for t and insert values for $N(t)/N_0$ and λ :

$$t = \frac{\ln(N(t)/N_0)}{-\lambda} = \frac{\ln 0.379}{-3.83 \times 10^{-12} \text{ s}^{-1}} = 2.53 \times 10^{11} \text{ s} = 8020 \text{ y}$$

EVALUATE After 8020 y the ^{14}C activity has decreased from 459 to 174 decays per hour. The specimen died and stopped taking CO_2 out of the air about 8000 years ago.

KEYCONCEPT To find the age of a sample using radioactive dating, measure its present activity and determine what its activity would have been when it formed. The age is proportional to the logarithm of the ratio of these two activities.

Radiation in the Home

A serious health hazard in some areas is the accumulation in houses of ^{222}Rn , an inert, colorless, odorless radioactive gas. Looking at the ^{238}U decay chain in Fig. 43.7, we see that the half-life of ^{222}Rn is 3.82 days. If so, why not just move out of the house for a while and let it decay away? The answer is that ^{222}Rn is continuously being *produced* by the decay of ^{226}Ra , which is found in minute quantities in the rocks and soil on which some houses are built. It's a dynamic equilibrium situation, in which the rate of production equals the rate of decay. The reason ^{222}Rn is a bigger hazard than the other elements in the ^{238}U decay series is that it's a gas. During its short half-life of 3.82 days it can migrate from the soil into your house. If a ^{222}Rn nucleus decays in your lungs, it emits a damaging α particle and its daughter nucleus ^{218}Po , which is *not* chemically inert and is likely to stay in your lungs until it decays, emits another damaging α particle and so on down the ^{238}U decay series.

How much of a hazard is radon? Although reports indicate values as high as 3500 pCi/L, the average activity per volume in the air inside American homes due to ^{222}Rn is about 1.5 pCi/L (over a thousand decays each second in an average-sized room). *If* your environment has this level of activity, it has been estimated that a lifetime exposure would reduce your life expectancy by about 40 days. For comparison, smoking one pack of cigarettes per day reduces life expectancy by 6 years, and it is estimated that the average emission from all the nuclear power plants in the world reduces life expectancy by anywhere from 0.01 day to 5 days. These figures include catastrophes such as the nuclear reactor disasters at Chernobyl, Ukraine (1986), and Fukushima, Japan (2011), for which the *local* effect on life expectancy is much greater.

TEST YOUR UNDERSTANDING OF SECTION 43.4 Which sample contains a greater number of nuclei: a $5.00 \mu\text{Ci}$ sample of ^{240}Pu (half-life 6560 y) or a $4.45 \mu\text{Ci}$ sample of ^{243}Am (half-life 7370 y)? (i) The ^{240}Pu sample; (ii) the ^{243}Am sample; (iii) both have the same number of nuclei.

ANSWER

The two samples contain *equal* numbers of nuclei. The ^{243}Am sample has a longer half-life and hence a slower decay rate, so it has a lower activity than the ^{240}Pu sample.

$$\frac{N_{\text{Pu}}}{N_{\text{Am}}} = \frac{(-dN_{\text{Pu}}/dt) T_{1/2-\text{Pu}}}{(-dN_{\text{Am}}/dt) T_{1/2-\text{Am}}} = \frac{(5.00 \mu\text{Ci})(6560 \text{ y})}{(4.45 \mu\text{Ci})(7370 \text{ y})} = 1.00$$

and we get
ratio of this expression for ^{240}Pu to this same expression for ^{243}Am , the factors of $\ln 2$ cancel
The activity $-dN(t)/dt$ of a sample is the product of the number of nuclei in the sample $N(t)$ and the decay constant $\lambda = (\ln 2)/T_{1/2}$. Hence $N(t) = [-dN(t)/dt] T_{1/2}/(\ln 2)$. Taking the

43.5 BIOLOGICAL EFFECTS OF RADIATION

The above discussion of radon introduced the interaction of radiation with living organisms, a topic of vital interest and importance. Under *radiation* we include radioactivity (alpha, beta, gamma, and neutrons) and electromagnetic radiation such as x rays. As these particles pass through matter, they lose energy, breaking molecular bonds and creating ions—hence the term *ionizing radiation*. Charged particles interact directly with the electrons in the material. X rays and γ rays interact by the photoelectric effect, in which an electron absorbs a photon and breaks loose from its site, or by Compton scattering (see Section 38.3). Neutrons cause ionization indirectly through collisions with nuclei or absorption by nuclei with subsequent radioactive decay of the resulting nuclei.

These interactions are extremely complex. It is well known that excessive exposure to radiation, including sunlight, x rays, and all the nuclear radiations, can destroy tissues. In mild cases it results in a burn, as with common sunburn. Greater exposure can cause very severe illness or death by a variety of mechanisms, including massive destruction of tissue cells, alterations of genetic material, and destruction of the components in bone marrow that produce red blood cells.

Calculating Radiation Doses

Radiation dosimetry is the quantitative description of the effect of radiation on living tissue. The *absorbed dose* of radiation is defined as the energy delivered to the tissue per unit mass. The SI unit of absorbed dose, the joule per kilogram, is called the *gray* (Gy); $1 \text{ Gy} = 1 \text{ J/kg}$. Another unit is the *rad*, defined as

$$1 \text{ rad} = 0.01 \text{ J/kg} = 0.01 \text{ Gy}$$

Absorbed dose by itself is not an adequate measure of biological effect because equal energies of different kinds of radiation cause different extents of biological effect. This variation is described by a numerical factor called the **relative biological effectiveness (RBE)**, also called the *quality factor* (QF), of each specific radiation. X rays with 200 keV of energy are defined to have an RBE of unity, and the effects of other radiations can be compared experimentally. **Table 43.3** shows approximate values of RBE for several radiations. All these values depend somewhat on the kind of tissue in which the radiation is absorbed and on the energy of the radiation.

The biological effect is described by the product of the absorbed dose and the RBE of the radiation; this quantity is called the *biologically equivalent dose*, or simply the equivalent dose. The SI unit of equivalent dose for humans is the sievert (Sv):

$$\text{Equivalent dose (Sv)} = \text{RBE} \times \text{Absorbed dose (Gy)} \quad (43.20)$$

A more common unit, corresponding to the rad, is the rem (an abbreviation of *röntgen equivalent for man*):

$$\text{Equivalent dose (rem)} = \text{RBE} \times \text{Absorbed dose (rad)} \quad (43.21)$$

Thus the unit of the RBE is 1 Sv/Gy or 1 rem/rad , and $1 \text{ rem} = 0.01 \text{ Sv}$.

EXAMPLE 43.10 Dose from a medical x ray

During a diagnostic x-ray examination a 1.2 kg portion of a broken leg receives an equivalent dose of 0.40 mSv. (a) What is the equivalent dose in mrem? (b) What is the absorbed dose in mrad and in mGy? (c) If the x-ray energy is 50 keV, how many x-ray photons are absorbed?

IDENTIFY and SET UP We are asked to relate the equivalent dose (the biological effect of the radiation, measured in sieverts or rems) to the absorbed dose (the energy absorbed per mass, measured in grays or rads). In part (a) we use the conversion factor $1 \text{ rem} = 0.01 \text{ Sv}$ for equivalent dose. Table 43.3 gives the RBE for x rays; we use this

TABLE 43.3 Relative Biological Effectiveness (RBE) for Several Types of Radiation

Radiation	RBE (Sv/Gy or rem/rad)
X rays and γ rays	1
Electrons	1.0–1.5
Slow neutrons	3–5
Protons	10
α particles	20
Heavy ions	20

Continued

value in part (b) to determine the absorbed dose from Eqs. (43.20) and (43.21). Finally, in part (c) we use the mass and the definition of absorbed dose to find the total energy absorbed and the total number of photons absorbed.

EXECUTE (a) The equivalent dose in mrem is

$$\frac{0.40 \text{ mSv}}{0.01 \text{ Sv/rem}} = 40 \text{ mrem}$$

(b) For x rays, $\text{RBE} = 1 \text{ rem/rad}$ or 1 Sv/Gy , so the absorbed dose is

$$\frac{40 \text{ mrem}}{1 \text{ rem/rad}} = 40 \text{ mrad}$$

$$\frac{0.40 \text{ mSv}}{1 \text{ Sv/Gy}} = 0.40 \text{ mGy} = 4.0 \times 10^{-4} \text{ J/kg}$$

(c) The total energy absorbed is

$$(4.0 \times 10^{-4} \text{ J/kg})(1.2 \text{ kg}) = 4.8 \times 10^{-4} \text{ J} = 3.0 \times 10^{15} \text{ eV}$$

The number of x-ray photons is

$$\frac{3.0 \times 10^{15} \text{ eV}}{5.0 \times 10^4 \text{ eV/photon}} = 6.0 \times 10^{10} \text{ photons}$$

EVALUATE The absorbed dose is relatively large because x rays have a low RBE. If the ionizing radiation had been a beam of α particles, for which $\text{RBE} = 20$, the absorbed dose needed for an equivalent dose of 0.40 mSv would be only 0.020 mGy, corresponding to a smaller total absorbed energy of $2.4 \times 10^{-5} \text{ J}$.

KEYCONCEPT The absorbed dose for a sample of tissue exposed to radiation is the amount of energy delivered per kilogram to the tissue. The equivalent dose equals the absorbed dose multiplied by the relative biological effectiveness (RBE), which depends on the type of radiation used.


Radiation Hazards

Here are a few numbers for perspective. To convert from Sv to rem, simply multiply by 100. An ordinary chest x-ray exam delivers about 0.20–0.40 mSv to about 5 kg of tissue. Radiation exposure from cosmic rays and natural radioactivity in soil, building materials, and so on is of the order of 2–3 mSv per year at sea level and twice that at an elevation of 1500 m (5000 ft). A whole-body dose of up to about 0.20 Sv causes no immediately detectable effect. A short-term whole-body dose of 5 Sv or more usually causes death within a few days or weeks. A localized dose of 100 Sv causes complete destruction of the exposed tissues.

The long-term hazards of radiation exposure in causing various cancers and genetic defects have been widely publicized, and the question of whether there is any “safe” level of radiation exposure has been hotly debated. U.S. government regulations are based on a maximum *yearly* exposure, from all except natural resources, of 2 to 5 mSv. Workers with occupational exposure to radiation are permitted an average of 20 mSv per year. Recent studies suggest that these limits are too high and that even extremely small exposures carry hazards, but it is very difficult to gather reliable statistics on the effects of low doses. It has become clear that any use of x rays for medical diagnosis should be preceded by a very careful estimation of the relationship of risk to possible benefit.

Another sharply debated question is that of radiation hazards from nuclear power plants. The radiation level from these plants is *not* negligible. However, to make a meaningful evaluation of hazards, we must compare these levels with the alternatives, such as coal-powered plants. The health hazards of coal smoke are serious and well documented, and the natural radioactivity in the smoke from a coal-fired power plant is believed to be roughly 100 times as great as that from a properly operating nuclear plant with equal capacity. But the comparison is not this simple; the possibility of a nuclear accident and the very serious problem of safe disposal of radioactive waste from nuclear plants must also be considered. **Figure 43.9** shows one estimate of the various sources of radiation exposure for the U.S. population. Ionizing radiation is a two-edged sword; it poses very serious health hazards, yet it also provides many benefits to humanity, including the diagnosis and treatments of disease and a wide variety of analytical techniques.

Figure 43.9 Contribution of various sources to the total average radiation exposure in the U.S. population, expressed as percentages of the total.



Beneficial Uses of Radiation

Radiation is widely used in medicine for intentional selective destruction of tissue such as tumors. The hazards are considerable, but if the disease would be fatal without treatment, any hazard may be preferable. Artificially produced isotopes are often used as radiation sources. Such isotopes have several advantages over naturally radioactive isotopes. They may have shorter half-lives and correspondingly greater activity. Isotopes can be chosen

that emit the type and energy of radiation desired. Some artificial isotopes have been replaced by photon, proton, and electron beams from linear accelerators.

Nuclear medicine is an expanding field of application. Radioactive isotopes have virtually the same electron configurations and resulting chemical behavior as stable isotopes of the same element. But the location and concentration of radioactive isotopes can easily be detected by measurements of the radiation they emit. A familiar example is the use of radioactive iodine for thyroid studies. Nearly all the iodine ingested is either eliminated or stored in the thyroid, and the body's chemical reactions do not discriminate between the unstable isotope ^{131}I and the stable isotope ^{127}I . A minute quantity of ^{131}I is fed or injected into the patient, and the speed with which it becomes concentrated in the thyroid provides a measure of thyroid function. The half-life is 8.02 days, so there are no long-lasting radiation hazards. By use of more sophisticated scanning detectors, one can also obtain a "picture" of the thyroid, which shows enlargement and other abnormalities. This procedure, a type of *autoradiography*, is comparable to photographing the glowing filament of an incandescent light bulb by using the light emitted by the filament itself. If this process discovers cancerous thyroid nodules, they can be destroyed by much larger quantities of ^{131}I .

Another useful nuclide for nuclear medicine is technetium-99 (^{99}Tc), which is formed in an excited state by the β^- decay of molybdenum (^{99}Mo). The technetium then decays to its ground state by emitting a γ -ray photon with energy 143 keV. The half-life is 6.01 hours, unusually long for γ emission. (The ground state of ^{99}Tc is also unstable, with a half-life of 2.11×10^5 y; it decays by β^- emission to the stable ruthenium nuclide ^{99}Ru .) The chemistry of technetium is such that it can readily be attached to organic molecules that are taken up by various organs of the body. A small quantity of such technetium-bearing molecules is injected into a patient, and a scanning detector or *gamma camera* is used to produce an image, or *scintigram*, that reveals which parts of the body take up these γ -emitting molecules. This technique, in which ^{99}Tc acts as a radioactive *tracer*, plays an important role in locating cancers, embolisms, and other pathologies (**Fig. 43.10**).

Tracer techniques have many other applications. Tritium (^3H), a radioactive hydrogen isotope, is used to tag molecules in complex organic reactions; radioactive tags on pesticide molecules, for example, can be used to trace their passage through food chains. In the world of machinery, radioactive iron can be used to study piston-ring wear. Laundry detergent manufacturers have even used radioactive dirt to test the effectiveness of their products.

Many direct effects of radiation are also useful, such as strengthening polymers by cross-linking, sterilizing surgical tools, dispersing unwanted static electricity in the air, and intentionally ionizing the air in smoke detectors. Gamma rays are also used to sterilize and preserve some food products.

TEST YOUR UNDERSTANDING OF SECTION 43.5 Alpha particles have 20 times the relative biological effectiveness of 200 keV x rays. Which would be better to use to radiate tissue deep inside the body? (i) A beam of alpha particles; (ii) a beam of 200 keV x rays; (iii) both are equally effective.

ANSWER

(ii) We saw in Section 43.3 that alpha particles can travel only a very short distance before they are stopped. By contrast, x-ray photons are very penetrating, so they can easily pass into the body.

43.6 NUCLEAR REACTIONS


In the preceding sections we studied the decay of unstable nuclei, especially spontaneous emission of an α or β particle, sometimes followed by γ emission. Nothing needs to be done to initiate this decay, and nothing can be done to control it. This section examines some *nuclear reactions*, rearrangements of nuclear components that result from a bombardment by a particle rather than a spontaneous natural process. Rutherford suggested

BIO APPLICATION A Radioactive Building

The United States Capitol building in Washington, DC, is made of granite that contains a small amount of naturally radioactive uranium. As a result, the radiation exposure to someone working in the Capitol is 0.85 mSv per year. The health effects of this are negligible; it is estimated that a person who spent 20 years inside the Capitol would have an extra 0.1% chance of getting cancer, above the 10% chance due to all other causes during the same 20 year period.



Figure 43.10 This colored scintigram shows where a chemical containing radioactive ^{99}Tc was taken up by a patient's lungs. The orange color in the lung on the left indicates strong γ -ray emission by the ^{99}Tc , which shows that the chemical was able to pass into this lung through the bloodstream. The lung on the right shows weaker emission, indicating the presence of an embolism (a blood clot or other obstruction in an artery) that is restricting the flow of blood to this lung.



in 1919 that a massive particle with sufficient kinetic energy might be able to penetrate a nucleus. The result would be either a new nucleus with greater atomic number and mass number or a decay of the original nucleus. Rutherford verified this when he bombarded nitrogen (^{14}N) with α particles and obtained an oxygen (^{17}O) nucleus and a proton:



Rutherford used alpha particles from naturally radioactive sources. In Chapter 44 we'll describe some of the particle accelerators that are now used to initiate nuclear reactions.

Nuclear reactions are subject to several *conservation laws*. The classical conservation principles for charge, momentum, angular momentum, and energy (including rest energies) are obeyed in all nuclear reactions. An additional conservation law, not anticipated by classical physics, is conservation of the total number of nucleons. The numbers of protons and neutrons need not be conserved separately; in β decay, neutrons and protons change into one another. We'll study the basis of the conservation of nucleon number in Chapter 44.

When two nuclei interact, charge conservation requires that the sum of the initial atomic numbers must equal the sum of the final atomic numbers. Because of conservation of nucleon number, the sum of the initial mass numbers must also equal the sum of the final mass numbers. In general, these are *not* elastic collisions, and the total initial mass does *not* equal the total final mass.

Reaction Energy

The difference between the masses before and after the reaction corresponds to the **reaction energy**, according to the mass–energy relationship $E = mc^2$. If initial particles A and B interact to produce final particles C and D , the reaction energy Q is defined as

$$Q = (M_A + M_B - M_C - M_D)c^2 \quad (\text{reaction energy}) \quad (43.23)$$

To balance the electrons, we use the neutral atomic masses in Eq. (43.23). That is, we use the mass of ${}^1_1\text{H}$ for a proton, ${}^2_1\text{H}$ for a deuteron, ${}^4_2\text{He}$ for an α particle, and so on. When Q is positive, the total mass decreases and the total kinetic energy increases. Such a reaction is called an *exoergic reaction*. When Q is negative, the mass increases and the kinetic energy decreases, and the reaction is called an *endoergic reaction*. The terms *exothermal* and *endothermal*, borrowed from chemistry, are also used. In an endoergic reaction the reaction cannot occur at all unless the initial kinetic energy in the center-of-mass reference frame is at least as great as $|Q|$. That is, there is a **threshold energy**, the minimum kinetic energy to make an endoergic reaction go.

EXAMPLE 43.11 Exoergic and endoergic reactions

(a) When a lithium-7 nucleus is bombarded by a proton, two alpha particles (${}^4\text{He}$) are produced. Find the reaction energy. (b) Calculate the reaction energy for the reaction ${}^4_2\text{He} + {}^{14}_7\text{N} \rightarrow {}^{17}_8\text{O} + {}^1_1\text{H}$.

IDENTIFY and SET UP The reaction energy Q for any nuclear reaction equals c^2 times the difference between the total initial mass and the total final mass, as in Eq. (43.23). Table 43.2 gives the required masses.

EXECUTE (a) The reaction is ${}^1_1\text{H} + {}^7_3\text{Li} \rightarrow {}^4_2\text{He} + {}^4_2\text{He}$. The initial and final masses and their respective sums are

A: ${}^1_1\text{H}$	1.007825 u	C: ${}^4_2\text{He}$	4.002603 u
B: ${}^7_3\text{Li}$	7.016003 u	D: ${}^4_2\text{He}$	4.002603 u
	8.023828 u		8.005206 u

The mass decreases by 0.018622 u. From Eq. (43.23), the reaction energy is

$$Q = (0.018622 \text{ u})(931.5 \text{ MeV/u}) = +17.35 \text{ MeV}$$

(b) The initial and final masses are

A: ${}^4_2\text{He}$	4.002603 u	C: ${}^{17}_8\text{O}$	16.999132 u
B: ${}^{14}_7\text{N}$	14.003074 u	D: ${}^1_1\text{H}$	1.007825 u
	18.005677 u		18.006957 u

The mass increases by 0.001280 u, and the corresponding reaction energy is

$$Q = (-0.001280 \text{ u})(931.5 \text{ MeV/u}) = -1.192 \text{ MeV}$$

EVALUATE The reaction in part (a) is *exoergic*: The final total kinetic energy of the two separating alpha particles is 17.35 MeV greater than the initial total kinetic energy of the proton and the lithium nucleus. The reaction in part (b) is *endoergic*: In the center-of-mass system—that is, in a head-on collision with zero total momentum—the minimum total initial kinetic energy required for this reaction to occur is 1.192 MeV.

KEYCONCEPT To find the energy released in a nuclear reaction (the reaction energy), calculate the difference between the total mass of the initial nuclei and the total mass of the final nuclei and then multiply the result by c^2 .

Ordinarily, the endoergic reaction of part (b) of Example 43.11 would be produced by bombarding stationary ^{14}N nuclei with alpha particles from an accelerator. In this case an alpha's kinetic energy must be *greater than* 1.192 MeV. If all the alpha's kinetic energy went solely to increasing the rest energy, the final kinetic energy would be zero, and momentum would not be conserved. When a particle with mass m and kinetic energy K collides with a stationary particle with mass M , the total kinetic energy K_{cm} in the center-of-mass coordinate system (the energy available to cause reactions) is

$$K_{\text{cm}} = \frac{M}{M + m}K \quad (43.24)$$

This expression assumes that the kinetic energies of the particles and nuclei are much less than their rest energies. We leave the derivation of Eq. (43.24) to you. In part (b) of Example 43.11, $M = 14.003074 \text{ u}$ and $m = 4.002603 \text{ u}$, so $M/(M + m) = (14.003074 \text{ u})/(18.005677 \text{ u}) = 0.7777$ and $K_{\text{cm}} = 0.7777K$. Since K_{cm} must be at least 1.192 MeV, the α particle's kinetic energy K must be at least $(1.192 \text{ MeV})/0.7777 = 1.533 \text{ MeV}$.

For a charged particle such as a proton or an α particle to penetrate the nucleus of another atom and cause a reaction, it must usually have enough initial kinetic energy to overcome the potential-energy barrier caused by the repulsive electrostatic forces. In the reaction of part (a) of Example 43.11, if we treat the proton and the ^7Li nucleus as spherically symmetric charges with radii given by Eq. (43.1), their centers will be $3.5 \times 10^{-15} \text{ m}$ apart when they touch. The repulsive potential energy of the proton (charge $+e$) and the ^7Li nucleus (charge $+3e$) at this separation r is

$$\begin{aligned} U &= \frac{1}{4\pi\epsilon_0} \frac{(e)(3e)}{r} = (9.0 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2) \frac{(3)(1.6 \times 10^{-19} \text{ C})^2}{3.5 \times 10^{-15} \text{ m}} \\ &= 2.0 \times 10^{-13} \text{ J} = 1.2 \text{ MeV} \end{aligned}$$

Even though the reaction is exoergic, the proton must have a minimum kinetic energy of about 1.2 MeV for the reaction to occur, unless the proton *tunnels* through the barrier (see Section 40.4).

Neutron Absorption

Absorption of *neutrons* by nuclei forms an important class of nuclear reactions. Heavy nuclei bombarded by neutrons can undergo a series of neutron absorptions alternating with beta decays, in which the mass number A increases by as much as 25. Some of the *transuranic elements*, elements having Z larger than 92, are produced in this way. These elements have not been found in nature. Many transuranic elements, having Z as high as 118, have been identified.

The analytical technique of *neutron activation analysis* uses similar reactions. When bombarded by neutrons, many stable nuclides absorb a neutron to become unstable and then undergo β^- decay. The energies of the β^- and associated γ emissions depend on the unstable nuclide and provide a means of identifying it and the original stable nuclide. Quantities of elements that are far too small for conventional chemical analysis can be detected in this way.


TEST YOUR UNDERSTANDING OF SECTION 43.6 The reaction described in part (a) of Example 43.11 is exoergic. Can it happen naturally when a sample of solid lithium is placed in a flask of hydrogen gas?

ANSWER

The reaction ${}^7_3\text{Li} + {}^2_1\text{H} \rightarrow {}^4_2\text{He} + {}^4_2\text{He}$ is a nuclear reaction that can take place only if a proton (a hydrogen nucleus) comes into contact with a lithium nucleus. If the hydrogen is in atomic form, the interaction between its electron cloud and the electron cloud of a lithium atom keeps the two nuclei from getting close to each other. Even if isolated protons are used, they must be fired at the lithium atoms with enough kinetic energy to overcome the electrical repulsion between the protons and the lithium nuclei. The statement that the reaction is exoergic means that more energy is released by the reaction than had to be put in to make the reaction occur.

43.7 NUCLEAR FISSION

Figure 43.11 Mass distribution of fission fragments from the fission of ${}^{236}\text{U}^*$ (an excited state of ${}^{235}\text{U}$), which is produced when ${}^{235}\text{U}$ absorbs a neutron. The vertical scale is logarithmic.




Nuclear fission is a decay process in which an unstable nucleus splits into two fragments of comparable mass. Fission was discovered in 1938 through the experiments of Otto Hahn and Fritz Strassmann in Germany. Pursuing earlier work by Enrico Fermi, they bombarded uranium ($Z = 92$) with neutrons. The resulting radiation did not coincide with that of any known radioactive nuclide. Urged on by their colleague Lise Meitner, they used meticulous chemical analysis to reach the astonishing but inescapable conclusion that they had found a radioactive isotope of barium ($Z = 56$). Later, radioactive krypton ($Z = 36$) was also found. Meitner and Otto Frisch correctly interpreted these results as showing that uranium nuclei were splitting into two massive fragments called *fission fragments*. Two or three free neutrons usually appear along with the fission fragments and, very occasionally, a light nuclide such as ${}^3_1\text{H}$.

Both the common isotope (99.3%) ${}^{238}\text{U}$ and the uncommon isotope (0.7%) ${}^{235}\text{U}$ (as well as several other nuclides) can be easily split by neutron bombardment: ${}^{235}\text{U}$ by slow neutrons (kinetic energy less than 1 eV) but ${}^{238}\text{U}$ only by fast neutrons with a minimum of about 1 MeV of kinetic energy. Fission resulting from neutron absorption is called *induced fission*. Some nuclides can also undergo *spontaneous fission* without initial neutron absorption, but this is quite rare. When ${}^{235}\text{U}$ absorbs a neutron, the resulting nuclide ${}^{236}\text{U}^*$ is in a highly excited state and splits into two fragments almost instantaneously. Strictly speaking, it is ${}^{236}\text{U}^*$, not ${}^{235}\text{U}$, that undergoes fission, but it's usual to speak of the fission of ${}^{235}\text{U}$.

Over 100 different nuclides, representing more than 20 different elements, have been found among the fission products. **Figure 43.11** shows the distribution of mass numbers for fission fragments from the fission of ${}^{235}\text{U}$.

Fission Reactions


You should check the following two typical fission reactions for conservation of nucleon number and charge:



The total kinetic energy of the fission fragments is enormous, about 200 MeV (compared to typical α and β energies of a few MeV). The reason for this is that nuclides at the high end of the mass spectrum (near $A = 240$) are less tightly bound than those nearer the middle ($A = 90$ to 145). Referring to Fig. 43.2, we see that the average binding energy per nucleon is about 7.6 MeV at $A = 240$ but about 8.5 MeV at $A = 120$. Therefore a rough estimate of the expected *increase* in binding energy during fission is about $8.5 \text{ MeV} - 7.6 \text{ MeV} = 0.9 \text{ MeV}$ per nucleon, or a total of $(235)(0.9 \text{ MeV}) \approx 200 \text{ MeV}$.

CAUTION Binding energy and rest energy It may seem to be a violation of conservation of energy to have an increase in both the binding energy and the kinetic energy during a fission reaction. But relative to the total rest energy E_0 of the separated nucleons, the rest energy of the nucleus is E_0 minus E_B . Thus an *increase* in binding energy corresponds to a *decrease* in rest energy as rest energy is converted to the kinetic energy of the fission fragments. ■

Figure 43.12 A liquid-drop model of fission.



Fission fragments always have too many neutrons to be stable. We noted in Section 43.3 that the neutron-to-proton ratio (N/Z) for stable nuclides is about 1 for light nuclides but almost 1.6 for the heaviest nuclides because of the increasing influence of the electrical repulsion of the protons. The N/Z value for stable nuclides is about 1.3 at $A = 100$ and 1.4 at $A = 150$. The fragments have about the same N/Z as ^{235}U , about 1.55. They usually respond to this surplus of neutrons by undergoing a series of β^- decays (each of which increases Z by 1 and decreases N by 1) until a stable value of N/Z is reached. A typical example is



The nuclide ^{140}Ce is stable. This series of β^- decays produces, on average, about 15 MeV of additional kinetic energy. The neutron excess of fission fragments also explains why two or three free neutrons are released during the fission.

Fission appears to set an upper limit on the production of transuranic nuclei, mentioned in Section 43.6, that are relatively stable. There are theoretical reasons to expect that nuclei near $Z = 114$, $N = 184$ or 196, might be stable with respect to spontaneous fission. In the shell model (see Section 43.2), these numbers correspond to filled shells and subshells in the nuclear energy-level structure. Such *superheavy nuclei* would still be unstable with respect to alpha emission. Physicists have found evidence for at least seven isotopes with $Z = 114$, the longest-lived of which has a half-life due to alpha decay of about 2.6 s.

Liquid-Drop Model

We can understand fission qualitatively on the basis of the liquid-drop model of the nucleus (see Section 43.2). The process is shown in **Fig. 43.12** in terms of an electrically charged liquid drop. These sketches shouldn't be taken too literally, but they may help to develop your intuition about fission. A ^{235}U nucleus absorbs a neutron (Fig. 43.12a), becoming a $^{236}\text{U}^*$ nucleus with excess energy (Fig. 43.12b). This excess energy causes violent oscillations, during which a neck between two lobes develops (Fig. 43.12c). The electrical repulsion of these two lobes stretches the neck farther (Fig. 43.12d), and finally two smaller fragments are formed (Fig. 43.12e) that move rapidly apart.

This qualitative picture has been developed into a more quantitative theory to explain why some nuclei undergo fission and others don't. **Figure 43.13** shows a hypothetical potential-energy function for two possible fission fragments. If neutron absorption results in an excitation energy greater than the energy barrier height U_B , fission occurs immediately. Even when there isn't quite enough energy to surmount the barrier, fission can take place by quantum-mechanical *tunneling*, discussed in Section 40.4. In principle, many stable heavy nuclei can fission by tunneling. But the probability depends very critically on the height and width of the barrier. For most nuclei this process is so unlikely that it is never observed.

Figure 43.13 Hypothetical potential-energy function for two fission fragments in a fissionable nucleus. At distances r beyond the range of the nuclear force, the potential energy varies approximately as $1/r$. Fission occurs if there is an excitation energy greater than U_B or an appreciable probability for tunneling through the potential-energy barrier.





Figure 43.14 Schematic diagram of a nuclear fission chain reaction.




BIO APPLICATION Making Radioactive Isotopes for Medicine

The fragments that result from nuclear fission are typically unstable, neutron-rich isotopes. A number of these are useful for medical diagnosis and cancer radiotherapy (see Section 43.5). This photograph shows a nuclear fission reactor used for producing such isotopes. The uranium fuel is kept in a large tank of water for cooling. Some of the neutron-rich fission fragments undergo beta decay and emit electrons that move faster than the speed of light in water (about $0.75c$). Like an airplane that produces an intense sonic boom when it flies faster than sound (see Section 16.9), these ultrafast electrons produce a “light boom” called *Čerenkov radiation* that has a characteristic blue color.



Chain Reactions

Fission of a uranium nucleus, triggered by neutron bombardment, releases other neutrons that can trigger more fissions, suggesting the possibility of a **chain reaction** (Fig. 43.14). The chain reaction may be made to proceed slowly and in a controlled manner in a nuclear reactor or explosively in a bomb. The energy release in a nuclear chain reaction is enormous, far greater than that in any chemical reaction. (In a sense, *fire* is a chemical chain reaction.) For example, when uranium is “burned” to uranium dioxide in the chemical reaction



the heat of combustion is about 4500 J/g . Expressed as energy per atom, this is about 11 eV per atom. By contrast, fission liberates about 200 MeV per atom, nearly 20 million times as much energy.

Nuclear Reactors

A *nuclear reactor* is a system in which a controlled nuclear chain reaction is used to liberate energy. In a nuclear power plant, this energy is used to generate steam, which operates a turbine and turns an electrical generator.

On average, each fission of a ${}^{235}\text{U}$ nucleus produces about 2.5 free neutrons, so 40% of the neutrons are needed to sustain a chain reaction. A ${}^{235}\text{U}$ nucleus is much more likely to absorb a low-energy neutron (less than 1 eV) than one of the higher-energy neutrons (1 MeV or so) that are liberated during fission. In a nuclear reactor the higher-energy neutrons are slowed down by collisions with nuclei in the surrounding material, called the *moderator*, so they are much more likely to cause further fissions. In nuclear power plants, the moderator is often




Figure 43.15 Schematic diagram of a nuclear power plant.

water, occasionally graphite. The *rate* of the reaction is controlled by inserting or withdrawing *control rods* made of elements (such as boron or cadmium) whose nuclei *absorb* neutrons without undergoing any additional reaction. The isotope ^{238}U can also absorb neutrons, leading to $^{239}\text{U}^*$, but not with high enough probability for it to sustain a chain reaction by itself. Thus uranium that is used in reactors is often “enriched” by increasing the proportion of ^{235}U above the natural value of 0.7%, typically to 3% or so, by isotope-separation processing.

The most familiar application of nuclear reactors is for the generation of electrical power. As was noted above, the fission energy appears as kinetic energy of the fission fragments, and its immediate result is to increase the internal energy of the fuel elements and the surrounding moderator. This increase in internal energy is transferred as heat to generate steam to drive turbines, which spin the electrical generators. **Figure 43.15** is a schematic diagram of a nuclear power plant. The energetic fission fragments heat the water surrounding the reactor core. The steam generator is a heat exchanger that takes heat from this highly radioactive water and generates nonradioactive steam to run the turbines.

A typical nuclear plant has an electric-generating capacity of 1000 MW (or 10^9 W). The turbines are heat engines and are subject to the efficiency limitations imposed by the second law of thermodynamics, discussed in Chapter 20. In modern nuclear plants the overall efficiency is about one-third, so 3000 MW of thermal power from the fission reaction is needed to generate 1000 MW of electrical power.

EXAMPLE 43.12 Uranium consumption in a nuclear reactor

What mass of ^{235}U must undergo fission each day to provide 3000 MW of thermal power?

IDENTIFY and SET UP Fission of ^{235}U liberates about 200 MeV per atom. We use this and the mass of the ^{235}U atom to determine the required amount of uranium.

EXECUTE Each second, we need 3000 MJ or 3000×10^6 J. Each fission provides 200 MeV, or

$$(200 \text{ MeV/fission})(1.6 \times 10^{-13} \text{ J/MeV}) = 3.2 \times 10^{-11} \text{ J/fission}$$

The number of fissions needed each second is

$$\frac{3000 \times 10^6 \text{ J}}{3.2 \times 10^{-11} \text{ J/fission}} = 9.4 \times 10^{19} \text{ fissions}$$

Each ^{235}U atom has a mass of $(235 \text{ u})(1.66 \times 10^{-27} \text{ kg/u}) = 3.9 \times 10^{-25} \text{ kg}$, so the mass of ^{235}U that undergoes fission each second is

$$(9.4 \times 10^{19})(3.9 \times 10^{-25} \text{ kg}) = 3.7 \times 10^{-5} \text{ kg} = 37 \mu\text{g}$$

In one day (86,400 s), the total consumption of ^{235}U is

$$(3.7 \times 10^{-5} \text{ kg/s})(86,400 \text{ s}) = 3.2 \text{ kg}$$

EVALUATE For comparison, a 1000 MW coal-fired power plant burns 10,600 tons (about 10 million kg) of coal per day!

KEYCONCEPT The thermal power output of a power plant equals the energy released per reaction multiplied by the number of reactions that take place per second. Since the energy released per nuclear fission reaction is very large, a substantial power output can be obtained from relatively few fission reactions per second.

We mentioned above that about 15 MeV of the energy released after fission of a ^{235}U nucleus comes from the β^- decays of the fission fragments. This fact poses a serious problem with respect to control and safety of reactors. Even after the chain reaction has been completely stopped by insertion of control rods into the core, heat continues to be evolved by the β^- decays, which cannot be stopped. For a 3000 MW reactor this heat power is initially very large, about 200 MW. In the event of total loss of cooling water, this power is more than enough to cause a catastrophic meltdown of the reactor core and possible penetration of the containment vessel. The difficulty in achieving a “cold shutdown” following an accident at the Three Mile Island nuclear power plant in Pennsylvania in March 1979 was a result of the continued evolution of heat due to β^- decays.

The catastrophe of April 26, 1986, at Chernobyl reactor No. 4 in Ukraine resulted from a combination of an inherently unstable design and several human errors committed during a test of the emergency core cooling system. Too many control rods were withdrawn to compensate for a decrease in power caused by a buildup of neutron absorbers such as ^{135}Xe . The power level rose from 1% of normal to 100 times normal in 4 seconds; a steam explosion ruptured pipes in the core cooling system and blew the heavy concrete cover off the reactor. The graphite moderator caught fire and burned for several days, and there was a meltdown of the core. The total activity of the radioactive material released into the atmosphere has been estimated as about 10^8 Ci.

TEST YOUR UNDERSTANDING OF SECTION 43.7 The fission of ^{235}U can be triggered by the absorption of a slow neutron by a nucleus. Can a slow *proton* be used to trigger ^{235}U fission?


ANSWER

no Because the neutron has no electric charge, it experiences no electrical repulsion from a ^{235}U nucleus. Hence a slow-moving neutron can approach and enter a ^{235}U nucleus, thereby providing the excitation needed to trigger fission. By contrast, a slow-moving *proton* (charge $+e$) feels a strong electrical repulsion from a ^{235}U nucleus (charge $+92e$). It never gets close to the nucleus, so it cannot trigger fission.

43.8 NUCLEAR FUSION

In a **nuclear fusion** reaction, two or more small light nuclei come together, or *fuse*, to form a larger nucleus. Fusion reactions release energy for the same reason as fission reactions: The binding energy per nucleon after the reaction is greater than before. Referring to Fig. 43.2, we see that the binding energy per nucleon increases with A up to about $A = 60$, so fusion of nearly any two light nuclei to make a nucleus with A less than 60 is likely to be an exoergic reaction. In comparison to fission, we are moving toward the peak of this curve from the opposite side. Another way to express the energy relationships is that the total mass of the products is less than that of the initial particles.


Here are three examples of energy-liberating fusion reactions, written in terms of the neutral atoms:



In the first reaction, two protons combine to form a deuteron (${}^2_1\text{H}$), with the emission of a positron (β^+) and an electron neutrino. In the second, a proton and a deuteron combine to form the nucleus of the light isotope of helium, ${}^3_2\text{He}$, with the emission of a gamma ray. Now double the first two reactions to provide the two ${}^3_2\text{He}$ nuclei that fuse in the third reaction to form an alpha particle (${}^4_2\text{He}$) and two protons. Together the reactions make up the process called the *proton-proton chain* (**Fig. 43.16**).

The net effect of the chain is the conversion of four protons into one α particle, two positrons, two electron neutrinos, and two γ 's. We can calculate the energy release from

Figure 43.16 The proton-proton chain.



this part of the process: The mass of an α particle plus two positrons is the mass of neutral ${}^4\text{He}$, the neutrinos have zero (or negligible) mass, and the gammas have zero mass.

Mass of four protons	4.029106 u
Mass of ${}^4\text{He}$	4.002603 u
Mass difference and energy release	0.026503 u and 24.69 MeV

The two positrons that are produced during the first step of the proton-proton chain collide with two electrons; mutual annihilation of the four particles takes place, and their rest energy is converted into $4(0.511 \text{ MeV}) = 2.044 \text{ MeV}$ of gamma radiation. Thus the total energy released is $(24.69 + 2.044) \text{ MeV} = 26.73 \text{ MeV}$. The proton-proton chain takes place in the interior of the sun and other stars (Fig. 43.17). Each gram of the sun's mass contains about 4.5×10^{23} protons. If all of these protons were fused into helium, the energy released would be about 130,000 kWh. If the sun were to continue to radiate at its present rate, it would take about 75×10^9 years to exhaust its supply of protons. As we'll soon see, fusion reactions can occur only at extremely high temperatures; in the sun, these temperatures are found only deep within the interior. Hence the sun cannot fuse *all* of its protons and can sustain fusion for only about 10×10^9 years in total. The present age of the solar system (including the sun) is 4.54×10^9 years, so the sun is about halfway through its available store of protons.

Figure 43.17 The energy released as starlight comes from fusion reactions deep within a star's interior. When a star is first formed and for most of its life, it converts the hydrogen in its core into helium. As a star ages, the core temperature can become high enough for additional fusion reactions that convert helium into carbon, oxygen, and other elements.



EXAMPLE 43.13 A fusion reaction

Two deuterons fuse to form a *triton* (a nucleus of tritium, or ${}^3\text{H}$) and a proton. How much energy is liberated?

IDENTIFY and SET UP This is a nuclear reaction of the type discussed in Section 43.6. We use Eq. (43.23) to find the energy released.

EXECUTE Adding one electron to each nucleus makes each a neutral atom; we find their masses in Table 43.2. Substituting into Eq. (43.23), we find

$$Q = [2(2.014102 \text{ u}) - 3.016049 \text{ u} - 1.007825 \text{ u}](931.5 \text{ MeV/u}) = 4.03 \text{ MeV}$$

EVALUATE Thus 4.03 MeV is released in the reaction; the triton and proton together have 4.03 MeV more kinetic energy than the two deuterons had together.

KEYCONCEPT When two light nuclei undergo fusion to form a more massive nucleus, the binding energy per nucleon increases. Hence energy is released in such fusion reactions.

Achieving Fusion

For two nuclei to undergo fusion, they must come together to within the range of the nuclear force, typically of the order of $2 \times 10^{-15} \text{ m}$. To do this, they must overcome the electrical repulsion of their positive charges. For two protons at this distance, the corresponding potential

energy is about 1.2×10^{-13} J or 0.7 MeV; this represents the total initial *kinetic* energy that the fusion nuclei must have—for example, 0.6×10^{-13} J each in a head-on collision.



Atoms have this much energy only at extremely high temperatures. The discussion of Section 18.3 showed that the average translational kinetic energy of a gas molecule at temperature T is $\frac{3}{2}kT$, where k is Boltzmann's constant. The temperature at which this is equal to $E = 0.6 \times 10^{-13}$ J is determined by the relationship

$$E = \frac{3}{2}kT$$

$$T = \frac{2E}{3k} = \frac{2(0.6 \times 10^{-13} \text{ J})}{3(1.38 \times 10^{-23} \text{ J/K})} = 3 \times 10^9 \text{ K}$$

Fusion reactions are possible at lower temperatures because the Maxwell–Boltzmann distribution function (see Section 18.5) gives a small fraction of protons with kinetic energies much higher than the average value. The proton-proton reaction occurs at “only” 1.5×10^7 K at the center of the sun, making it an extremely low-probability process; but that's why the sun is expected to last so long. At these temperatures the fusion reactions are called *thermonuclear* reactions.

Intensive efforts are under way to achieve controlled fusion reactions, which potentially represent an enormous new resource of energy (see Fig. 24.11). At the temperatures mentioned, light atoms are fully ionized, and the resulting state of matter is called a *plasma*. In one kind of experiment using *magnetic confinement*, a plasma is heated to extremely high temperature by an electrical discharge, while being contained by appropriately shaped magnetic fields. In another, through *inertial confinement*, pellets of the material to be fused are heated by a high-intensity laser beam (see Fig. 43.18). Some of the reactions being studied are



We considered reaction (1) in Example 43.13; two deuterons fuse to form a triton and a proton. In reaction (2) a triton combines with another deuteron to form an alpha particle and a neutron. The result of these two reactions together is the conversion of three deuterons into an alpha particle, a proton, and a neutron, with 21.6 MeV of energy liberated. Reactions (3) and (4) together achieve the same conversion. In a plasma that contains deuterons, the two pairs of reactions occur with roughly equal probability. As yet, no one has succeeded in producing these reactions under controlled conditions in such a way as to yield a net surplus of usable energy.


Methods of achieving fusion that don't require high temperatures are also being studied; these are called *cold fusion*. One successful scheme uses an unusual hydrogen molecule ion. The usual H_2^+ ion consists of two protons bound by one shared electron; the nuclear spacing is about 0.1 nm. If the protons are replaced by a deuteron (${}^2\text{H}$) and a triton (${}^3\text{H}$) and the electron by a *muon*, which is 208 times as massive as the electron, the spacing is reduced by a factor of 208. The probability then becomes appreciable for the two nuclei to tunnel through the narrow repulsive potential-energy barrier and fuse in reaction (2) above. The prospect of making this process, called *muon-catalyzed fusion*, into a practical energy source is still distant.

TEST YOUR UNDERSTANDING OF SECTION 43.8 Are all fusion reactions exoergic?

ANSWER

no Fusion reactions between sufficiently light nuclei are exoergic because the binding energy per nucleon E_B/A increases. If the nuclei are too massive, however, E_B/A decreases and fusion is *endoergic* (i.e., it takes in energy rather than releasing it). As an example, imagine fusing together two nuclei of $A = 100$ to make a single nucleus with $A = 200$. From Fig. 43.2, E_B/A is more than 8.5 MeV for the $A = 100$ nuclei but is less than 8 MeV for the $A = 200$ nucleus. Such a fusion reaction is possible, but requires a substantial input of energy.

Figure 43.18 This target chamber at the National Ignition Facility in California has apertures for 192 powerful laser beams. The lasers deliver 5×10^{14} W of power for a few nanoseconds to a millimeter-sized pellet of deuterium and tritium at the center of the chamber, thus triggering thermonuclear fusion.




CHAPTER 43 SUMMARY

Nuclear properties: A nucleus is composed of A nucleons (Z protons and N neutrons). All nuclei have about the same density. The radius of a nucleus with mass number A is given approximately by Eq. (43.1). A single nuclear species of a given Z and N is called a nuclide. Isotopes are nuclides of the same element (same Z) that have different numbers of neutrons. Nuclear masses are measured in atomic mass units. Nucleons have angular momentum and a magnetic moment. (See Examples 43.1 and 43.2.)

$$R = R_0 A^{1/3} \quad (43.1)$$

$$(R_0 = 1.2 \times 10^{-15} \text{ m})$$


Nuclear binding and structure: The mass of a nucleus is always less than the mass of the protons and neutrons within it. The mass difference multiplied by c^2 gives the binding energy E_B . The binding energy for a given nuclide is determined by the nuclear force, which is short range and favors pairs of particles, and by the electrical repulsion between protons. A nucleus is unstable if A or Z is too large or if the ratio N/Z is wrong. Two widely used models of the nucleus are the liquid-drop model and the shell model; the latter is analogous to the central-field approximation for atomic structure. (See Examples 43.3 and 43.4.)




Radioactive decay: Unstable nuclides usually emit an alpha particle (a ${}^4_2\text{He}$ nucleus) or a beta particle (an electron) in the process of changing to another nuclide, sometimes followed by a gamma-ray photon. The rate of decay of an unstable nucleus is described by the decay constant λ , the half-life $T_{1/2}$, or the lifetime T_{mean} . If the number of nuclei at time $t = 0$ is N_0 and no more are produced, the number at time t is given by Eq. (43.17). (See Examples 43.5–43.9.)

$$N(t) = N_0 e^{-\lambda t} \quad (43.17)$$


$$T_{\text{mean}} = \frac{1}{\lambda} = \frac{T_{1/2}}{\ln 2} = \frac{T_{1/2}}{0.693} \quad (43.19)$$



Biological effects of radiation: The biological effect of any radiation depends on the product of the energy absorbed per unit mass and the relative biological effectiveness (RBE), which is different for different radiations. (See Example 43.10.)



Nuclear reactions: In a nuclear reaction, two nuclei or particles collide to produce two new nuclei or particles. Reactions can be exoergic or endoergic. Several conservation laws, including charge, energy, momentum, angular momentum, and nucleon number, are obeyed. Energy is released by the fission of a heavy nucleus into two lighter, always unstable, nuclei. Energy is also released by the fusion of two light nuclei into a heavier nucleus. (See Examples 43.11–43.13.)





GUIDED PRACTICE

For assigned homework and other learning materials, go to Mastering Physics.

KEY EXAMPLE VARIATION PROBLEMS

Be sure to review **EXAMPLES 43.3 and 43.4** (Section 43.2) before attempting these problems.

VP43.4.1 Find (a) the mass defect, (b) the total binding energy, and (c) the binding energy per nucleon of $^{16}_8\text{O}$, which has a neutral atomic mass of 15.994915 u.

VP43.4.2 The mass defect of $^{63}_{29}\text{Cu}$ is 0.5919378 u. Find (a) its binding energy, (b) its binding energy per nucleon, and (c) its neutral atomic mass.

VP43.4.3 Calculate the binding energy E_B/A for the nuclides (a) $^{75}_{33}\text{As}$ (mass defect 0.7005604 u), (b) $^{150}_{62}\text{Sm}$ (mass defect 1.330388 u), and (c) $^{225}_{88}\text{Ra}$ (mass defect 1.852094 u). (d) Figure 43.2 shows that for nuclei with nucleon numbers $A > 62$, the binding energy per nucleon E_B/A decreases as A increases. Do your results agree with this?

VP43.4.4 Use the semiempirical mass formula to calculate the total estimated binding energy E_B and the binding energy per nucleon E_B/A for the nuclides (a) ruthenium-100 ($^{100}_{44}\text{Ru}$) and (b) mercury-200 ($^{200}_{80}\text{Hg}$). (c) Are your results consistent with Fig. 43.2?

Be sure to review **EXAMPLES 43.5, 43.6, and 43.7** (Section 43.3) before attempting these problems.

VP43.7.1 Household smoke detectors contain a tiny amount of radioactive americium that decays by α emission to neptunium: $^{241}_{95}\text{Am} \rightarrow ^{237}_{93}\text{Np} + ^4_2\text{He}$. The atomic masses are 241.056827 u for $^{241}_{95}\text{Am}$, 237.048172 u for $^{237}_{93}\text{Np}$, and 4.002603 u for ^4_2He . If the $^{241}_{95}\text{Am}$ nucleus is originally at rest, find (a) the total energy released in the decay, (b) the kinetic energy of the emitted α particle, and (c) the speed of the emitted α particle.

VP43.7.2 The beryllium isotope ^8_4Be is very unstable, and rapidly decays into two α particles: $^8_4\text{Be} \rightarrow ^4_2\text{He} + ^4_2\text{He}$. The atomic masses are 8.0053051 u for ^8_4Be and 4.0026033 u for ^4_2He . If the ^8_4Be nucleus is originally at rest, find (a) the kinetic energy and (b) the speed of each emitted α particle.

VP43.7.3 State whether each of the following proposed β^- decays is possible or impossible. If it is possible, find the total energy released in the decay. The atomic mass is given for each nuclide. (a) $^{12}_4\text{Be}$

(12.026922 u) to $^{12}_5\text{B}$ (12.014353 u); (b) $^{33}_{17}\text{Cl}$ (32.977452 u) to $^{33}_{18}\text{Ar}$ (32.989926 u); (c) $^{82}_{35}\text{Br}$ (81.9168018 u) to $^{82}_{36}\text{Kr}$ (81.9134812 u).

VP43.7.4 State whether each of the following proposed β^+ decays is possible or impossible. If it is possible, find the total energy released in the decay; if it is impossible, state whether the decay can occur by electron capture. The atomic mass is given for each nuclide. The combined mass of two electrons is 0.001097 u. (a) $^{46}_{23}\text{V}$ (45.960198 u) to $^{46}_{22}\text{Ti}$ (45.952627 u); (b) $^{134}_{57}\text{La}$ (133.908514 u) to $^{134}_{56}\text{Ba}$ (133.904508 u); (c) $^{67}_{31}\text{Ga}$ (66.9282024 u) to $^{67}_{30}\text{Zn}$ (66.9271275 u).

Be sure to review **EXAMPLES 43.8 and 43.9** (Section 43.4) before attempting these problems.

VP43.9.1 The sodium isotope $^{24}_{11}\text{Na}$ undergoes β^- decay to $^{24}_{12}\text{Mg}$ (a stable isotope) with a half-life of 15.0 h. Initially a sample of $^{24}_{11}\text{Na}$ has β^- activity of 1.50 μCi . Find (a) the mean lifetime, (b) the decay constant, (c) the initial number of $^{24}_{11}\text{Na}$ nuclei in the sample, and (d) the activity of the sample after 24.0 h.

VP43.9.2 You are given a sample that was initially 100% $^{95}_{41}\text{Nb}$. This isotope undergoes β^- decay to $^{95}_{42}\text{Mo}$ (a stable isotope) with a half-life of 35.0 h. You find that the sample now contains 23.8% $^{95}_{41}\text{Nb}$ and 76.2% $^{95}_{42}\text{Mo}$ and has a β^- activity of 0.514 μCi . Find (a) the decay constant, (b) the present number of $^{95}_{41}\text{Nb}$ nuclei in the sample, (c) the initial number of $^{95}_{41}\text{Nb}$ nuclei in the sample, and (d) how much time has elapsed since the sample was 100% $^{95}_{41}\text{Nb}$.

VP43.9.3 A certain isotope undergoes γ decay from an excited state to its ground state. Initially a sample of this isotope has an activity of 0.425 μCi ; 45.0 s later, the activity has decreased to 0.121 μCi . Find (a) the decay constant and half-life of the excited state and (b) the number of nuclei in the excited state initially and 45.0 s later.

VP43.9.4 Before 1900 the activity per unit mass of atmospheric carbon due to the presence of ^{14}C (half-life 5730 y) averaged about 0.255 Bq per gram of carbon. You detect 113 decays of ^{14}C in 20.0 min from a sample of carbon of mass 0.510 g. Assuming that its activity per unit mass of carbon when it formed was the pre-1900 average value for the air, find (a) the age of the sample and (b) the current number of ^{14}C nuclei in the sample.

BRIDGING PROBLEM Saturation of ^{128}I Production

In an experiment, the iodine isotope ^{128}I is created by irradiating a sample of ^{127}I with a beam of neutrons, yielding 1.50×10^6 ^{128}I nuclei per second. Initially no ^{128}I nuclei are present. A ^{128}I nucleus decays by β^- emission with a half-life of 25.0 min. (a) To what nuclide does ^{128}I decay? (b) Could that nuclide decay back to ^{128}I by β^+ emission? Why or why not? (c) After the sample has been irradiated for a long time, what is the maximum number of ^{128}I atoms that can be present in the sample? What is the maximum activity that can be produced? (This steady-state situation is called *saturation*.) (d) Find an expression for the number of ^{128}I atoms present in the sample as a function of time.

SOLUTION GUIDE

IDENTIFY and SET UP

1. What happens to the values of Z , N , and A in β^- decay? What must be true for β^- decay to be possible? For β^+ decay to be possible?
2. You'll need to write an equation for the rate of change dN/dt of the number N of ^{128}I atoms in the sample, taking account of both the creation of ^{128}I by the neutron irradiation and the decay of

any ^{128}I present. In the steady state, how do the rates of these two processes compare?

3. List the unknown quantities for each part of the problem and identify your target variables.

EXECUTE

4. Find the values of Z and N of the nuclide produced by the decay of ^{128}I . What element is this?
5. Decide whether this nuclide can decay back to ^{128}I .
6. Inspect your equation for dN/dt . What is the value of dN/dt in the steady state? Use this to solve for the steady-state values of N and the activity.
7. Solve your dN/dt equation for the function $N(t)$. (Hint: See Section 26.4.)

EVALUATE

8. Your result from step 6 tells you the value of N after a long time (that is, for large values of t). Is this consistent with your result from step 7? What would constitute a "long time" under these conditions?

PROBLEMS

•, ••, •••: Difficulty levels. **CP**: Cumulative problems incorporating material from earlier chapters. **CALC**: Problems requiring calculus. **DATA**: Problems involving real data, scientific evidence, experimental design, and/or statistical reasoning. **BIO**: Biosciences problems.

DISCUSSION QUESTIONS

Q43.1 BIO Neutrons have a magnetic dipole moment and can undergo spin flips by absorbing electromagnetic radiation. Why, then, are protons rather than neutrons used in MRI of body tissues? (See Fig. 43.1.)

Q43.2 In Eq. (43.11), as the total number of nucleons becomes larger, the importance of the second term in the equation decreases relative to that of the first term. Does this make physical sense? Explain.

Q43.3 Why aren't the masses of all nuclei integer multiples of the mass of a single nucleon?

Q43.4 The only two stable nuclides with more protons than neutrons are ${}^1_1\text{H}$ and ${}^3_2\text{He}$. Why is $Z > N$ so uncommon?

Q43.5 What are the six known elements for which Z is a magic number? Discuss what properties these elements have as a consequence of their special values of Z .

Q43.6 The binding energy per nucleon for most nuclides doesn't vary much (see Fig. 43.2). Is there similar consistency in the *atomic* energy of atoms, on an "energy per electron" basis? If so, why? If not, why not?

Q43.7 Heavy, unstable nuclei usually decay by emitting an α or a β particle. Why don't they usually emit a single proton or neutron?

Q43.8 As stars age, they use up their supply of hydrogen and eventually begin producing energy by a reaction that involves the fusion of three helium nuclei to form a carbon nucleus. Would you expect the interiors of these old stars to be hotter or cooler than the interiors of younger stars? Explain.

Q43.9 Since lead is a stable element, why doesn't the ${}^{238}\text{U}$ decay series shown in Fig. 43.7 stop at lead, ${}^{214}\text{Pb}$?

Q43.10 In the ${}^{238}\text{U}$ decay series shown in Fig. 43.7, some nuclides in the series are found much more abundantly in nature than others, even though every ${}^{238}\text{U}$ nucleus goes through every step in the series before finally becoming ${}^{206}\text{Pb}$. Why don't the intermediate nuclides all have the same abundance?

Q43.11 Compared to α particles with the same energy, β particles can much more easily penetrate through matter. Why is this?

Q43.12 If ${}^A_Z\text{Ei}$ represents the initial nuclide, what is the decay process or processes if the final nuclide is (a) ${}_{Z+1}^A\text{Ei}$; (b) ${}_{Z-2}^A\text{Ei}$; (c) ${}_{Z-1}^A\text{Ei}$?

Q43.13 In a nuclear decay equation, why can we represent an electron as ${}^0_{-1}\beta^-$? What are the equivalent representations for a positron, a neutrino, and an antineutrino?

Q43.14 Why is the alpha, beta, or gamma decay of an unstable nucleus unaffected by the *chemical* situation of the atom, such as the nature of the molecule or solid in which it is bound? The chemical situation of the atom can, however, have an effect on the half-life in electron capture. Why is this?

Q43.15 In the process of *internal conversion*, a nucleus decays from an excited state to a ground state by giving the excitation energy directly to an atomic electron rather than emitting a gamma-ray photon. Why can this process also produce x-ray photons?

Q43.16 In Example 43.9 (Section 43.4), the activity of atmospheric carbon *before* 1900 was given. Discuss why this activity may have changed since 1900.

Q43.17 BIO One problem in radiocarbon dating of biological samples, especially very old ones, is that they can easily be contaminated with modern biological material during the measurement process. What effect would such contamination have on the estimated age? Why is such contamination a more serious problem for samples of older material than for samples of younger material?

Q43.18 The most common radium isotope found on earth, ${}^{226}\text{Ra}$, has a half-life of about 1600 years. If the earth was formed well over 10^9 years ago, why is there any radium left now?

Q43.19 Fission reactions occur only for nuclei with large nucleon numbers, while exoergic fusion reactions occur only for nuclei with small nucleon numbers. Why is this?

Q43.20 When a large nucleus splits during nuclear fission, the daughter nuclei of the fission fly apart with enormous kinetic energy. Why does this happen?

EXERCISES

Section 43.1 Properties of Nuclei

43.1 • How many protons and how many neutrons are there in a nucleus of the most common isotope of (a) silicon, ${}^{28}_{14}\text{Si}$; (b) rubidium, ${}^{85}_{37}\text{Rb}$; (c) thallium, ${}^{205}_{81}\text{Tl}$?

43.2 • Hydrogen atoms are placed in an external magnetic field. The protons can make transitions between states in which the nuclear spin component is parallel and antiparallel to the field by absorbing or emitting a photon. What magnetic-field magnitude is required for this transition to be induced by photons with frequency 22.7 MHz?

Section 43.2 Nuclear Binding and Nuclear Structure

43.3 • The most common isotope of boron is ${}^{11}_5\text{B}$. (a) Determine the total binding energy of ${}^{11}_5\text{B}$ from Table 43.2 in Section 43.1. (b) Calculate this binding energy from Eq. (43.11). (Why is the fifth term zero?) Compare to the result you obtained in part (a). What is the percent difference? Compare the accuracy of Eq. (43.11) for ${}^{11}_5\text{B}$ to its accuracy for ${}^{62}_{28}\text{Ni}$ (see Example 43.4).

43.4 •• The nuclei ${}^{11}_5\text{B}$ and ${}^{11}_6\text{C}$ are called *mirror nuclei*, because the number of protons in ${}^{11}_5\text{B}$ equals the number of neutrons in ${}^{11}_6\text{C}$ and the number of neutrons in ${}^{11}_5\text{B}$ equals the number of protons in ${}^{11}_6\text{C}$. The atomic mass of ${}^{11}_6\text{C}$ is 11.011434 u, and the atomic mass of ${}^{11}_5\text{B}$ is given in Table 43.2. (a) Calculate the binding energies of ${}^{11}_5\text{B}$ and ${}^{11}_6\text{C}$. (b) Which of the two nuclei has the larger binding energy? Why is this so?

43.5 • Calculate (a) the total binding energy and (b) the binding energy per nucleon of ${}^{12}_6\text{C}$. (c) What percent of the rest mass of this nucleus is its total binding energy?

43.6 • The most common isotope of uranium, ${}^{238}_{92}\text{U}$, has atomic mass 238.050788 u. Calculate (a) the mass defect; (b) the binding energy (in MeV); (c) the binding energy per nucleon.

43.7 • The binding energy per nucleon for ${}^{56}_{26}\text{Fe}$ is 8.79 MeV/nucleon. What is the mass, in amu, of a neutral ${}^{56}_{26}\text{Fe}$ atom? Give your answer to five significant figures.

43.8 •• An alpha particle is strongly bound. The ${}^{12}_6\text{C}$ nucleus might be modeled as a composite of three alpha particles. Compare the binding energy of ${}^{12}_6\text{C}$ with three times the binding energy of an alpha particle. Which of these quantities is larger, and why might this be so?

43.9 • CP A photon with a wavelength of 3.50×10^{-13} m strikes a deuteron, splitting it into a proton and a neutron. (a) Calculate the kinetic energy released in this interaction. (b) Assuming the two particles share the energy equally, and taking their masses to be 1.00 u, calculate their speeds after the photodisintegration.

43.10 • Calculate the mass defect, the binding energy (in MeV), and the binding energy per nucleon of (a) the nitrogen nucleus, ${}^{14}_7\text{N}$, and (b) the helium nucleus, ${}^4_2\text{He}$. (c) How does the binding energy per nucleon compare for these two nuclei?

Section 43.3 Nuclear Stability and Radioactivity

43.11 • The carbon isotope $^{11}_6\text{C}$ undergoes β^+ (positron) decay. The atomic mass of $^{11}_6\text{C}$ is 11.011433 u. (a) How many protons and how many neutrons are in the daughter nucleus produced by this decay? (b) How much energy, in MeV, is released in the decay of one $^{11}_6\text{C}$ nucleus?

43.12 • (a) Is the decay $n \rightarrow p + \beta^- + \bar{\nu}_e$ energetically possible? If not, explain why not. If so, calculate the total energy released. (b) Is the decay $p \rightarrow n + \beta^+ + \nu_e$ energetically possible? If not, explain why not. If so, calculate the total energy released.

43.13 • What nuclide is produced in the following radioactive decays? (a) α decay of $^{239}_{94}\text{Pu}$; (b) β^- decay of $^{24}_{11}\text{Na}$; (c) β^+ decay of $^{15}_8\text{O}$.

43.14 •• CP ^{238}U decays spontaneously by α emission to ^{234}Th . Calculate (a) the total energy released by this process and (b) the recoil velocity of the ^{234}Th nucleus. The atomic masses are 238.050788 u for ^{238}U and 234.043601 u for ^{234}Th .

43.15 • The radioactive isotope $^{12}_7\text{N}$ undergoes β^+ decay. The energy released in the decay of one nucleus is 16.316 MeV. What is the mass, in amu, of a neutral $^{12}_7\text{N}$ atom? Give your answer to eight significant figures.

43.16 • What particle (α particle, electron, or positron) is emitted in the following radioactive decays? (a) $^{27}_{14}\text{Si} \rightarrow ^{27}_{13}\text{Al}$; (b) $^{238}_{92}\text{U} \rightarrow ^{234}_{90}\text{Th}$; (c) $^{74}_{33}\text{As} \rightarrow ^{74}_{34}\text{Se}$.

43.17 • Tritium (^3_1H) is an unstable isotope of hydrogen; its mass, including one electron, is 3.016049 u. (a) Show that tritium must be unstable with respect to beta decay because the decay products (^3_2He plus an emitted electron) have less total mass than the tritium. (b) Determine the total kinetic energy (in MeV) of the decay products, taking care to account for the electron masses correctly.

Section 43.4 Activities and Half-Lives

43.18 • A sample of radioactive nuclei has N_0 nuclei at time $t = 0$. The half-life of the decay is $T_{1/2}$. In terms of N_0 , how many decays occur in the time period between $t = 0$ and $t = 0.500T_{1/2}$?

43.19 • If a 6.13 g sample of an isotope having a mass number of 124 decays at a rate of 0.350 Ci, what is its half-life?

43.20 • BIO Radioactive isotopes used in cancer therapy have a “shelf-life,” like pharmaceuticals used in chemotherapy. Just after it has been manufactured in a nuclear reactor, the activity of a sample of ^{60}Co is 5000 Ci. When its activity falls below 3500 Ci, it is considered too weak a source to use in treatment. You work in the radiology department of a large hospital. One of these ^{60}Co sources in your inventory was manufactured on October 6, 2011. It is now April 6, 2014. Is the source still usable? The half-life of ^{60}Co is 5.271 years.

43.21 •• The common isotope of uranium, ^{238}U , has a half-life of 4.47×10^9 years, decaying to ^{234}Th by alpha emission. (a) What is the decay constant? (b) What mass of uranium is required for an activity of 1.00 curie? (c) How many alpha particles are emitted per second by 10.0 g of uranium?

43.22 •• BIO Radiation Treatment of Prostate Cancer. In many cases, prostate cancer is treated by implanting 60 to 100 small seeds of radioactive material into the tumor. The energy released from the decays kills the tumor. One isotope that is used (there are others) is palladium (^{103}Pd) with a half-life of 17 days. If a typical grain contains 0.250 g of ^{103}Pd , (a) what is its initial activity rate in Bq, and (b) what is the rate 68 days later?

43.23 •• A 12.0 g sample of carbon from living matter decays at the rate of 184 decays/minute due to the radioactive ^{14}C in it. What will be the decay rate of this sample in (a) 1000 years and (b) 50,000 years?

43.24 •• BIO Radioactive Tracers. Radioactive isotopes are often introduced into the body through the bloodstream. Their spread through the body can then be monitored by detecting the appearance of radiation in different organs. One such tracer is ^{131}I , a β^- emitter with a half-life

of 8.0 d. Suppose a scientist introduces a sample with an activity of 325 Bq and watches it spread to the organs. (a) Assuming that all of the sample went to the thyroid gland, what will be the decay rate in that gland 24 d (about $3\frac{1}{2}$ weeks) later? (b) If the decay rate in the thyroid 24 d later is measured to be 17.0 Bq, what percentage of the tracer went to that gland? (c) What isotope remains after the I-131 decays?

43.25 • The isotope $^{90}_{39}\text{Y}$ undergoes β^- decay with a half-life of 64.0 h. You measure an activity of 8.0×10^{16} Bq. (a) How many $^{90}_{39}\text{Y}$ nuclei are present in the sample at the time you make this measurement? (b) How many will be present after 12.0 days?

43.26 • As a health physicist, you are being consulted about a spill in a radiochemistry lab. The isotope spilled was 400 μCi of ^{131}Ba , which has a half-life of 12 days. (a) What mass of ^{131}Ba was spilled? (b) Your recommendation is to clear the lab until the radiation level has fallen 1.00 μCi . How long will the lab have to be closed?

43.27 • Measurements on a certain isotope tell you that the decay rate decreases from 8318 decays/min to 3091 decays/min in 4.00 days. What is the half-life of this isotope?

43.28 •• Radiocarbon Dating. At an archeological site, a sample from timbers containing 500 g of carbon provides 2690 decays/min. What is the age of the sample?

43.29 • The radioactive nuclide ^{199}Pt has a half-life of 30.8 minutes. A sample is prepared that has an initial activity of 7.56×10^{11} Bq. (a) How many ^{199}Pt nuclei are initially present in the sample? (b) How many are present after 30.8 minutes? What is the activity at this time? (c) Repeat part (b) for a time 92.4 minutes after the sample is first prepared.

Section 43.5 Biological Effects of Radiation

43.30 • BIO Radiation Overdose. If a person's entire body is exposed to 5.0 J/kg of x rays, death usually follows within a few days. (a) Express this lethal radiation dose in Gy, rad, Sv, and rem. (b) How much total energy does a 70.0 kg person absorb from such a dose? (c) If the 5.0 J/kg came from a beam of protons instead of x rays, what would be the answers to parts (a) and (b)?

43.31 •• BIO A nuclear chemist receives an accidental radiation dose of 5.0 Gy from slow neutrons ($\text{RBE} = 4.0$). What does she receive in rad, rem, and J/kg?

43.32 •• BIO A person exposed to fast neutrons receives a radiation dose of 300 rem on part of his hand, affecting 25 g of tissue. The RBE of these neutrons is 10. (a) How many rad did he receive? (b) How many joules of energy did he receive? (c) Suppose the person received the same rad dosage, but from beta rays with an RBE of 1.0 instead of neutrons. How many rem would he have received?

43.33 • BIO Food Irradiation. Food is often irradiated with either x rays or electron beams to help prevent spoilage. A low dose of 5–75 kilorads (krad) helps to reduce and kill inactive parasites, a medium dose of 100–400 krad kills microorganisms and pathogens such as salmonella, and a high dose of 2300–5700 krad sterilizes food so that it can be stored without refrigeration. (a) A dose of 175 krad kills spoilage microorganisms in fish. If x rays are used, what would be the dose in Gy, Sv, and rem, and how much energy would a 220 g portion of fish absorb? (See Table 43.3.) (b) Repeat part (a) if electrons of RBE 1.50 are used instead of x rays.

43.34 • BIO To Scan or Not to Scan? It has become popular for some people to have yearly whole-body scans (CT scans, formerly called CAT scans) using x rays, just to see if they detect anything suspicious. A number of medical people have recently questioned the advisability of such scans, due in part to the radiation they impart. Typically, one such scan gives a dose of 12 mSv, applied to the *whole body*. By contrast, a chest x ray typically administers 0.20 mSv to only 5.0 kg of tissue. How many chest x rays would deliver the same *total* amount of energy to the body of a 75 kg person as one whole-body scan?

43.35 •• The radioactive isotope $^{90}_{38}\text{Sr}$ is found in the fallout from atmospheric tests of nuclear weapons (see Problem 43.49). It replaces calcium in items such as bone and milk. It has a half-life of 29 years, and the net energy absorbed from each decay averages about 1.1 MeV. (a) Calculate the absorbed dose from 1.0 microgram of $^{90}_{38}\text{Sr}$ in one year if the energy is absorbed in 50 kg of body tissue. (b) The radiation from $^{90}_{38}\text{Sr}$ involves beta-minus particles and gamma rays. Calculate the equivalent dose in part (a) if you assume the RBE for the electrons is 1.0.

43.36 • BIO In an industrial accident a 65 kg person receives a lethal whole-body equivalent dose of 5.4 Sv from x rays. (a) What is the equivalent dose in rem? (b) What is the absorbed dose in rad? (c) What is the total energy absorbed by the person's body? How does this amount of energy compare to the amount of energy required to raise the temperature of 65 kg of water 0.010 C°?

43.37 •• CP BIO In a diagnostic x-ray procedure, 5.00×10^{10} photons are absorbed by tissue with a mass of 0.600 kg. The x-ray wavelength is 0.0200 nm. (a) What is the total energy absorbed by the tissue? (b) What is the equivalent dose in rem?


Section 43.6 Nuclear Reactions

Section 43.7 Nuclear Fission

Section 43.8 Nuclear Fusion


43.38 •• Calculate the reaction energy Q for the reaction $p + {}^3_1\text{H} \rightarrow {}^2_1\text{H} + {}^2_1\text{H}$. Is this reaction exoergic or endoergic?

43.39 • Consider the nuclear reaction




where X is a nuclide. (a) What are the values of Z and A for the nuclide X? (b) How much energy is liberated? (c) Estimate the threshold energy for this reaction.

43.40 • Energy from Nuclear Fusion. Calculate the energy released in the fusion reaction




43.41 •• At the beginning of Section 43.7 the equation of a fission process is given in which $^{235}_{92}\text{U}$ is struck by a neutron and undergoes fission to produce $^{144}_{54}\text{Ba}$, $^{89}_{36}\text{Kr}$, and three neutrons. The measured masses of these isotopes are 235.043930 u ($^{235}_{92}\text{U}$), 143.922953 u ($^{144}_{54}\text{Ba}$), 88.917631 u ($^{89}_{36}\text{Kr}$), and 1.0086649 u (neutron). (a) Calculate the energy (in MeV) released by each fission reaction. (b) Calculate the energy released per gram of $^{235}_{92}\text{U}$, in MeV/g.

43.42 •• Consider the nuclear reaction



where X is a nuclide. (a) What are Z and A for the nuclide X? (b) Ignoring the effects of recoil, what minimum energy must the photon have for this reaction to occur? The mass of a ${}^{28}_{14}\text{Si}$ atom is 27.976927 u, and the mass of a ${}^{24}_{12}\text{Mg}$ atom is 23.985042 u.

43.43 • Consider the nuclear reaction



where X is a nuclide. (a) What are Z and A for the nuclide X? (b) Is energy absorbed or liberated? How much?

PROBLEMS

43.44 •• CP The nucleus $^{10}_4\text{Be}$ undergoes β^- decay. The atomic mass of $^{10}_4\text{Be}$ is 10.013535 u. (a) How much energy, in MeV, is released in this decay? (b) If you assume that all the energy released in the decay appears as kinetic energy of the β^- particle (a very accurate assumption), what is the speed of the emitted β^- ? Note that the energy released is nearly the same as the rest energy of the β^- .

43.45 • Comparison of Energy Released per Gram of Fuel.

(a) When gasoline is burned, it releases 1.3×10^8 J of energy per gallon (3.788 L). Given that the density of gasoline is 737 kg/m³, express the quantity of energy released in J/g of fuel. (b) During fission, when a neutron is absorbed by a $^{235}_{92}\text{U}$ nucleus, about 200 MeV of energy is released for each nucleus that undergoes fission. Express this quantity in J/g of fuel. (c) In the proton-proton chain that takes place in stars like our sun, the overall fusion reaction can be summarized as six protons fusing to form one ${}^4_2\text{He}$ nucleus with two leftover protons and the liberation of 26.7 MeV of energy. The fuel is the six protons. Express the energy produced here in units of J/g of fuel. Notice the huge difference between the two forms of nuclear energy, on the one hand, and the chemical energy from gasoline, on the other. (d) Our sun produces energy at a measured rate of 3.86×10^{26} W. If its mass of 1.99×10^{30} kg were all gasoline, how long could it last before consuming all its fuel? (*Historical note:* Before the discovery of nuclear fusion and the vast amounts of energy it releases, scientists were confused. They knew that the earth was at least many millions of years old, but could not explain how the sun could survive that long if its energy came from chemical burning.)

43.46 •• (a) Calculate the minimum energy required to remove one proton from the nucleus $^{12}_6\text{C}$. This is called the proton-removal energy. (*Hint:* Find the difference between the mass of a $^{12}_6\text{C}$ nucleus and the mass of a proton plus the mass of the nucleus formed when a proton is removed from $^{12}_6\text{C}$.) (b) How does the proton-removal energy for $^{12}_6\text{C}$ compare to the binding energy per nucleon for $^{12}_6\text{C}$, calculated using Eq. (43.10)?

43.47 •• (a) Calculate the minimum energy required to remove one neutron from the nucleus $^{17}_8\text{O}$. This is called the neutron-removal energy. (See Problem 43.46.) (b) How does the neutron-removal energy for $^{17}_8\text{O}$ compare to the binding energy per nucleon for $^{17}_8\text{O}$, calculated using Eq. (43.10)?

43.48 •• Heat flows to the surface of the earth from its interior. Two-thirds of this heat is radiogenic in origin, which means it comes from the decay of radioactive elements in the earth's mantle, primarily from the isotopes $^{238}_{92}\text{U}$ and $^{232}_{90}\text{Th}$. We can estimate the earth's internal energy supply due to the decay of uranium as follows: (a) There are 31 μg of $^{238}_{92}\text{U}$ in each kg of mantle. Estimate how many $^{238}_{92}\text{U}$ isotopes there are per kg of mantle. (b) The half-life of $^{238}_{92}\text{U}$ is 4.47 billion years. Calculate the decay constant in units of s⁻¹. (c) Multiply the results in parts (a) and (b) to obtain the activity in each kg of mantle. (d) Each decay chain $^{238}_{92}\text{U} \rightarrow {}^{206}_{82}\text{Pb}$ releases 52 MeV of energy. Use this value to convert the activity into watts per kg of mantle. (e) The mass of the earth is 6×10^{24} kg, and two-thirds of this is mantle mass. Use these values to estimate the power earth receives from uranium decays. (f) Uranium decays account for 39% of the radiogenic supply of heat. Estimate earth's total heat power related to radioactive decays. (g) Earth also releases heat that remains from its formation, with a power comparable with that due to radioactivity. Estimate the total heat power earth receives from its interior.

43.49 • BIO Radioactive Fallout. One of the problems of in-air testing of nuclear weapons (or, even worse, the use of such weapons!) is the danger of radioactive fallout. One of the most problematic nuclides in such fallout is strontium-90 ($^{90}_{38}\text{Sr}$), which breaks down by β^- decay with a half-life of 28 years. It is chemically similar to calcium and therefore can be incorporated into bones and teeth, where, due to its rather long half-life, it remains for years as an internal source of radiation. (a) What is the daughter nucleus of the $^{90}_{38}\text{Sr}$ decay? (b) What percentage of the original level of $^{90}_{38}\text{Sr}$ is left after 56 years? (c) How long would you have to wait for the original level to be reduced to 6.25% of its original value?

43.50 •• CP Thorium $^{230}_{90}\text{Th}$ decays to radium $^{226}_{88}\text{Ra}$ by α emission. The masses of the neutral atoms are 230.033134 u for $^{230}_{90}\text{Th}$ and 226.025410 u for $^{226}_{88}\text{Ra}$. If the parent thorium nucleus is at rest, what is the kinetic energy of the emitted α particle? (Be sure to account for the recoil of the daughter nucleus.)

43.51 •• The atomic mass of $^{25}_{12}\text{Mg}$ is 24.985837 u, and the atomic mass of $^{25}_{13}\text{Al}$ is 24.990428 u. (a) Which of these nuclei will decay into the other? (b) What type of decay will occur? Explain how you determined this. (c) How much energy (in MeV) is released in the decay?

43.52 •• The fusion reactions in the sun produce 26.73 MeV per alpha particle created. Each of these reactions produces two neutrinos. In total, the sun's power is about 3.8×10^{26} W. (a) Estimate how many solar fusion reactions take place every second. (b) Estimate the number of neutrinos that leave the sun each second. (c) The distance between the earth and the sun is 150 million km, and the radius of the earth is 6370 km. Using these figures, estimate the number of solar neutrinos that encounter the earth each second. (d) How many square centimeters does the earth present to the sun? (e) Divide your answer in part (c) by your answer in part (d) to obtain an estimate of the number of solar neutrinos that pass through every square centimeter on earth every second.

43.53 •• BIO Irradiating Ourselves! The radiocarbon in our bodies is one of the naturally occurring sources of radiation. Let's see how large a dose we receive. ^{14}C decays via β^- emission, and 18% of our body's mass is carbon. (a) Write out the decay scheme of carbon-14 and show the end product. (A neutrino is also produced.) (b) Neglecting the effects of the neutrino, how much kinetic energy (in MeV) is released per decay? The atomic mass of ^{14}C is 14.003242 u. (c) How many grams of carbon are there in a 75 kg person? How many decays per second does this carbon produce? (*Hint:* Use data from Example 43.9.) (d) Assuming that all the energy released in these decays is absorbed by the body, how many MeV/s and J/s does the ^{14}C release in this person's body? (e) Consult Table 43.3 and use the largest appropriate RBE for the particles involved. What radiation dose does the person give himself in a year, in Gy, rad, Sv, and rem?

43.54 •• BIO Pion Radiation Therapy. A neutral pion (π^0) has a mass of 264 times the electron mass and decays with a lifetime of 8.4×10^{-17} s to two photons. Such pions are used in the radiation treatment of some cancers. (a) Find the energy and wavelength of these photons. In which part of the electromagnetic spectrum do they lie? What is the RBE for these photons? (b) If you want to deliver a dose of 200 rem (which is typical) in a single treatment to 25 g of tumor tissue, how many π^0 mesons are needed?

43.55 •• Calculate the mass defect for the β^+ decay of $^{11}_6\text{C}$. Is this decay energetically possible? Why or why not? The atomic mass of $^{11}_6\text{C}$ is 11.011434 u.

43.56 •• BIO A person ingests an amount of a radioactive source that has a very long lifetime and activity $0.52 \mu\text{Ci}$. The radioactive material lodges in her lungs, where all of the emitted 4.0 MeV α particles are absorbed within a 0.50 kg mass of tissue. Calculate the absorbed dose and the equivalent dose for one year.

43.57 • We Are Stardust. In 1952 spectral lines of the element technetium-99 (^{99}Tc) were discovered in a red giant star. Red giants are very old stars, often around 10 billion years old, and near the end of their lives. Technetium has *no* stable isotopes, and the half-life of ^{99}Tc is 200,000 years. (a) For how many half-lives has the ^{99}Tc been in the red giant star if its age is 10 billion years? (b) What fraction of the original ^{99}Tc would be left at the end of that time? This discovery was extremely important because it provided convincing evidence for the theory (now essentially known to be true) that most of the atoms heavier than hydrogen and helium were made inside of stars by thermonuclear fusion and other nuclear processes. If the ^{99}Tc had been part of the star since it was born, the amount remaining after 10 billion years would have been so minute that it would not have been detectable. This knowledge is what led the late astronomer Carl Sagan to proclaim that "we are stardust."

43.58 • BIO A 70.0 kg person experiences a whole-body exposure to α radiation with energy 4.77 MeV. A total of 7.75×10^{12} α particles are absorbed. (a) What is the absorbed dose in rad? (b) What is the equivalent dose in rem? (c) If the source is 0.0320 g of ^{226}Ra (half-life 1600 y) somewhere in the body, what is the activity of this source? (d) If all of the alpha particles produced are absorbed, what time is required for this dose to be delivered?

43.59 •• Measurements indicate that 27.83% of all rubidium atoms currently on the earth are the radioactive ^{87}Rb isotope. The rest are the stable ^{85}Rb isotope. The half-life of ^{87}Rb is 4.75×10^{10} y. Assuming that no rubidium atoms have been formed since, what percentage of rubidium atoms were ^{87}Rb when our solar system was formed 4.6×10^9 y ago?

43.60 • The nucleus $^{15}_8\text{O}$ has a half-life of 122.2 s; $^{19}_8\text{O}$ has a half-life of 26.9 s. If at some time a sample contains equal amounts of $^{15}_8\text{O}$ and $^{19}_8\text{O}$, what is the ratio of $^{15}_8\text{O}$ to $^{19}_8\text{O}$ (a) after 3.0 min and (b) after 12.0 min?

43.61 •• BIO A ^{60}Co source with activity 2.6×10^{-4} Ci is embedded in a tumor that has mass 0.200 kg. The source emits γ photons with average energy 1.25 MeV. Half the photons are absorbed in the tumor, and half escape. (a) What energy is delivered to the tumor per second? (b) What absorbed dose (in rad) is delivered per second? (c) What equivalent dose (in rem) is delivered per second if the RBE for these γ rays is 0.70? (d) What exposure time is required for an equivalent dose of 200 rem?

43.62 ••• The proton-proton chain described in Section 43.8, known as the p-p I chain, accounts for 83.3% of the helium synthesis in the sun. This process uses six protons to produce one alpha particle, two protons, two neutrinos, and a least six gamma-ray photons. We can determine the energies of each of the products as follows: (a) The first reaction, $2^1_1\text{H} \rightarrow ^2_1\text{H} + \beta^+ + \nu_e$, is followed immediately by the annihilation of the positron by an electron into two photons, each with energy equal to the rest energy of an electron. What is the remaining energy E_1 carried by the deuteron and the neutrino? (b) Work in the frame where the net momentum of the deuteron and neutrino is zero. Denote the (relativistic) energy of the neutrino by E_ν and the (nonrelativistic) deuteron speed by v_d . Denote the mass of the deuteron by m_d . What are the conditions for energy and momentum conservation for these two particles, in terms of E_1 ? Treat the neutrino as massless. (c) Solve the equations from part (b) and use the deuteron mass in Table 43.2 to determine the percentage of the energy E_1 carried by the neutrino. (d) What is the total energy carried away by neutrinos in the p-p I chain? (*Note:* This chain creates two neutrinos by the above mechanism.) (e) What percentage of the energy generated by the p-p I chain is carried away by neutrinos? (*Note:* The total energy of the p-p I chain is computed in Section 43.8.)

43.63 ••• The sun produces a minority 16.68% of its helium-4 using the p-p II chain, in which four protons and an ambient alpha particle produce two alpha particles, two neutrinos, and at least four gamma-ray photons. The first two steps are the same as the first two steps of the p-p I chain (see Problem 43.62). However, the third step is $^3_2\text{He} + ^4_2\text{He} \rightarrow ^7_4\text{Be} + \gamma$, where ^7_4Be is an unstable beryllium nuclide with mass 7.016930 u. (a) What is the energy of the photon? (b) The beryllium nuclide then captures an electron, which changes one of its protons into a neutron, resulting in $^7_4\text{Be} + e^- \rightarrow ^7_3\text{Li} + \nu_e$, where ^7_3Li is the common (and stable) nuclide of lithium, with mass 7.016003 u. The excess energy is carried away by the neutrino. What is the energy of that neutrino? (c) A proton then collides with the lithium nucleus, resulting in two alpha particles via $^1_1\text{H} + ^7_3\text{Li} \rightarrow 2^4_2\text{He}$. How much energy is released in this step? (d) If each alpha particle has the same speed, then in the rest frame of the lithium disintegration, what is the speed of each alpha particle? (e) How much energy is released in total by the processes in this p-p II chain? (f) How does this compare with the energy released in the p-p I chain? (g) What percentage of the energy in the p-p II chain is carried away by neutrinos?

43.64 ••• In addition to the p-p I and p-p II chains (see Problems 43.62 and 43.63), the sun produces helium-4 in a relatively rare reaction called the p-p III chain. This produces only 0.1% of the sun's alpha particles but results in neutrinos with substantially higher energy, which have proved crucial to the study of neutrino physics. In the p-p III chain, an ambient ${}^3_2\text{He}$ fuses with ambient ${}^4_2\text{He}$ (alpha particles) to produce ${}^7_4\text{Be}$, precisely as in the p-p II chain. But in the p-p III chain, the ${}^7_4\text{Be}$ then absorbs a proton to produce ${}^8_5\text{B}$. The ${}^8_5\text{B}$ then decays to ${}^8_4\text{Be}$ along with a positron and a neutrino. (a) The total energy released in this process can be ascertained from its net input and output products $4\,{}^1_1\text{H} \rightarrow {}^4_2\text{He}$. How much energy is released? (b) Given the isotope masses 8.024607 u for ${}^8_5\text{B}$ and 8.0053051 u for ${}^8_4\text{Be}$, determine the energy of the neutrino emitted in the ${}^8_5\text{B}$ to ${}^8_4\text{Be}$ decay. (c) What percentage of the energy in the p-p III chain is carried away by neutrinos? (Note: This chain also produces a low-energy neutrino.)

43.65 • A bone fragment found in a cave believed to have been inhabited by early humans contains 0.29 times as much ${}^{14}\text{C}$ as an equal amount of carbon in the atmosphere when the organism containing the bone died. (See Example 43.9 in Section 43.4.) Find the approximate age of the fragment.

43.66 •• BIO In the 1986 disaster at the Chernobyl reactor in eastern Europe, about $\frac{1}{8}$ of the ${}^{137}\text{Cs}$ present in the reactor was released. The isotope ${}^{137}\text{Cs}$ has a half-life of 30.07 y for β decay, with the emission of a total of 1.17 MeV of energy per decay. Of this, 0.51 MeV goes to the emitted electron; the remaining 0.66 MeV goes to a γ ray. The radioactive ${}^{137}\text{Cs}$ is absorbed by plants, which are eaten by livestock and humans. How many ${}^{137}\text{Cs}$ atoms would need to be present in each kilogram of body tissue if an equivalent dose for one week is 3.5 Sv? Assume that all of the energy from the decay is deposited in 1.0 kg of tissue and that the RBE of the electrons is 1.5.

43.67 •• Consider the fusion reaction ${}^2_1\text{H} + {}^2_1\text{H} \rightarrow {}^3_2\text{He} + {}^1_0\text{n}$. (a) Estimate the barrier energy by calculating the repulsive electrostatic potential energy of the two ${}^2_1\text{H}$ nuclei when they touch. (b) Compute the energy liberated in this reaction in MeV and in joules. (c) Compute the energy liberated *per mole* of deuterium, remembering that the gas is diatomic, and compare with the heat of combustion of hydrogen, about 2.9×10^5 J/mol.

43.68 •• DATA As a scientist in a nuclear physics research lab, you are conducting a photodisintegration experiment to verify the binding energy of a deuteron. A photon with wavelength λ in air is absorbed by a deuteron, which breaks apart into a neutron and a proton. The two fragments share the released kinetic energy equally, and the deuteron can be assumed to be initially at rest. You measure the speed of the proton after the disintegration as a function of the wavelength λ of the photon. Your experimental results are given in the table.

λ (10^{-13} m)	3.50	3.75	4.00	4.25	4.50	4.75	5.00
v (10^6 m/s)	11.3	10.2	9.1	8.1	7.2	6.1	4.9

(a) Graph the data as v^2 versus $1/\lambda$. Explain why the data points, when graphed this way, should follow close to a straight line. Find the slope and y-intercept of the straight line that gives the best fit to the data. (b) Assume that h and c have their accepted values. Use your results from part (a) to calculate the mass of the proton and the binding energy (in MeV) of the deuteron.

43.69 •• DATA Your company develops radioactive isotopes for medical applications. In your work there, you measure the activity of a radioactive sample. Your results are given in the table.

Time (h)	Decays/s
0	20,000
0.5	14,800
1.0	11,000
1.5	8130
2.0	6020
2.5	4460
3.0	3300
4.0	1810
5.0	1000
6.0	550
7.0	300

(a) Find the half-life of the sample. (b) How many radioactive nuclei were present in the sample at $t = 0$? (c) How many were present after 7.0 h?

43.70 ••• DATA In your job as a health physicist, you measure the activity of a mixed sample of radioactive elements. Your results are given in the table.

Time (h)	Decays/s
0	7500
0.5	4120
1.0	2570
1.5	1790
2.0	1350
2.5	1070
3.0	872
4.0	596
5.0	404
6.0	288
7.0	201
8.0	140
9.0	98
10.0	68
12.0	33

(a) What minimum number of different nuclides are present in the mixture? (b) What are their half-lives? (c) How many nuclei of each type are initially present in the sample? (d) How many of each type are present at $t = 5.0$ h?

CHALLENGE PROBLEMS

43.71 ••• Industrial Radioactivity. Radioisotopes are used in a variety of manufacturing and testing techniques. Wear measurements can be made using the following method. An automobile engine is produced using piston rings with a total mass of 100 g, which includes $9.4\,\mu\text{Ci}$ of ${}^{59}\text{Fe}$ whose half-life is 45 days. The engine is test-run for 1000 hours, after which the oil is drained and its activity is measured. If the activity of the engine oil is 84 decays/s, how much mass was worn from the piston rings per hour of operation?

43.72 ••• Many radioactive decays occur within a sequence of decays—for example, ${}^{234}_{92}\text{U} \rightarrow {}^{230}_{88}\text{Th} \rightarrow {}^{226}_{84}\text{Ra}$. The half-life for the ${}^{234}_{92}\text{U} \rightarrow {}^{230}_{88}\text{Th}$ decay is 2.46×10^5 y, and the half-life for the ${}^{230}_{88}\text{Th} \rightarrow {}^{226}_{84}\text{Ra}$ decay is 7.54×10^4 y. Let 1 refer to ${}^{234}_{92}\text{U}$, 2 to ${}^{230}_{88}\text{Th}$, and 3 to ${}^{226}_{84}\text{Ra}$; let λ_1 be the decay constant for the ${}^{234}_{92}\text{U} \rightarrow {}^{230}_{88}\text{Th}$ decay and λ_2 be the decay constant for the ${}^{230}_{88}\text{Th} \rightarrow {}^{226}_{84}\text{Ra}$ decay. The amount of ${}^{230}_{88}\text{Th}$ present at any time depends on the rate at which it is produced by the decay of ${}^{234}_{92}\text{U}$ and the rate by which it is depleted by its decay to ${}^{226}_{84}\text{Ra}$. Therefore, $dN_2(t)/dt = \lambda_1 N_1(t) - \lambda_2 N_2(t)$. If we start with a sample that contains N_{10} nuclei of ${}^{234}_{92}\text{U}$ and nothing else, then $N(t) = N_{10}e^{-\lambda_1 t}$. Thus $dN_2(t)/dt = \lambda_1 N_{10}e^{-\lambda_1 t} - \lambda_2 N_2(t)$. This differential equation for $N_2(t)$ can be solved as follows. Assume a trial solution of the form $N_2(t) = N_{10}[h_1 e^{-\lambda_1 t} + h_2 e^{-\lambda_2 t}]$, where h_1 and h_2 are constants. (a) Since $N_2(0) = 0$, what must be the relationship between h_1 and h_2 ? (b) Use the trial solution to calculate $dN_2(t)/dt$, and substitute that into the differential equation for $N_2(t)$. Collect the coefficients of $e^{-\lambda_1 t}$ and $e^{-\lambda_2 t}$. Since the equation must hold at all t , each of these coefficients must be zero. Use this requirement to solve for h_1 and thereby complete the determination of $N_2(t)$. (c) At time $t = 0$, you have a pure sample containing 30.0 g of ${}^{234}_{92}\text{U}$ and nothing else. What mass of ${}^{230}_{88}\text{Th}$ is present at time $t = 2.46 \times 10^5$ y, the half-life for the ${}^{234}_{92}\text{U}$ decay?

MCAT-STYLE PASSAGE PROBLEMS

BIO Radioactive Iodine in Medicine. Iodine in the body is preferentially taken up by the thyroid gland. Therefore, radioactive iodine in small doses is used to image the thyroid and in large doses is used to kill thyroid cells to treat some types of cancer or thyroid disease. The iodine isotopes used have relatively short half-lives, so they must be produced in a nuclear reactor or accelerator. One isotope frequently used for imaging is ${}^{123}_{53}\text{I}$; it has a half-life of 13.2 h and emits a 0.16 MeV

gamma-ray photon. One method of producing ${}^{123}_{53}\text{I}$ is in the nuclear reaction ${}^{123}_{52}\text{Te} + p \rightarrow {}^{123}_{53}\text{I} + n$. The atomic masses relevant to this reaction are ${}^{123}_{52}\text{Te}$, 122.904270 u; ${}^{123}_{53}\text{I}$, 122.905589 u; n, 1.008665 u; and ${}^1_1\text{H}$, 1.007825 u.

The iodine isotope commonly used for treatment of disease is ${}^{131}_{53}\text{I}$, which is produced by irradiating ${}^{130}_{52}\text{Te}$ in a nuclear reactor to form ${}^{131}_{52}\text{Te}$. The ${}^{131}_{52}\text{Te}$ then decays to ${}^{131}_{53}\text{I}$. ${}^{131}_{53}\text{I}$ undergoes β^- decay with a half-life of 8.04 d, emitting electrons with energies up to 0.61 MeV and gamma-ray photons of energy 0.36 MeV. A typical thyroid cancer treatment might involve administration of 3.7 GBq of ${}^{131}_{53}\text{I}$.

43.73 Which reaction produces ${}^{131}_{53}\text{I}$ in the nuclear reactor? (a) ${}^{130}_{52}\text{Te} + n \rightarrow {}^{131}_{52}\text{Te}$; (b) ${}^{130}_{53}\text{I} + n \rightarrow {}^{131}_{53}\text{I}$; (c) ${}^{132}_{52}\text{Te} + n \rightarrow {}^{131}_{53}\text{I}$; (d) ${}^{132}_{53}\text{I} + n \rightarrow {}^{131}_{53}\text{I}$.

43.74 Which type of radioactive decay produces ${}^{131}_{53}\text{I}$ from ${}^{131}_{52}\text{Te}$? (a) Alpha decay; (b) β^- decay; (c) β^+ decay; (d) gamma decay.

43.75 How many ${}^{131}_{53}\text{I}$ atoms are administered in a typical thyroid cancer treatment? (a) 4.2×10^{10} ; (b) 1.0×10^{12} ; (c) 2.5×10^{14} ; (d) 3.7×10^{15} .

43.76 In the reaction that produces ${}^{123}_{53}\text{I}$, is there a minimum kinetic energy the protons need to make the reaction go? (a) No, because the proton has a smaller mass than the neutron. (b) No, because the total initial mass is smaller than the total final mass. (c) Yes, because the proton has a smaller mass than the neutron. (d) Yes, because the total initial mass is smaller than the total final mass.

43.77 Why might ${}^{123}_{53}\text{I}$ be preferred for imaging over ${}^{131}_{53}\text{I}$? (a) The atomic mass of ${}^{123}_{53}\text{I}$ is smaller, so the ${}^{123}_{53}\text{I}$ particles travel farther through tissue. (b) Because ${}^{123}_{53}\text{I}$ emits only gamma-ray photons, the radiation dose to the body is lower with that isotope. (c) The beta particles emitted by ${}^{131}_{53}\text{I}$ can leave the body, whereas the gamma-ray photons emitted by ${}^{123}_{53}\text{I}$ cannot. (d) ${}^{123}_{53}\text{I}$ is radioactive, whereas ${}^{131}_{53}\text{I}$ is not.

ANSWERS

Chapter Opening Question ?

(iv) When an organism dies, it stops taking in carbon from atmospheric CO_2 . Some of this carbon is radioactive ${}^{14}_6\text{C}$, which decays with a half-life of 5730 years. By measuring the proportion of ${}^{14}_6\text{C}$ that remains in the specimen, scientists can determine how long ago the organism died. (See Section 43.4.)

Key Example VARIATION Problems

VP43.4.1 (a) 0.137005 u (b) 127.6 MeV (c) 7.976 MeV

VP43.4.2 (a) 551.4 MeV (b) 8.752 MeV (c) 62.929597 u

VP43.4.3 (a) 8.701 MeV (b) 8.262 MeV (c) 7.668 MeV (d) yes

VP43.4.4 (a) $E_B = 868.1$ MeV, $E_B/A = 8.681$ MeV

(b) $E_B = 1584$ MeV, $E_B/A = 7.922$ MeV (c) yes

VP43.7.1 (a) 5.637 MeV (b) 5.543 MeV (c) 1.63×10^7 m/s

VP43.7.2 (a) 0.0459 MeV = 45.9 keV (b) 1.49×10^6 m/s

VP43.7.3 (a) possible, 11.71 MeV (b) impossible (c) possible, 3.093 MeV

VP43.7.4 (a) possible, 6.030 MeV (b) possible, 2.710 MeV (c) impossible, yes

VP43.9.1 (a) 7.79×10^4 s (b) $1.28 \times 10^{-5} \text{ s}^{-1}$ (c) 4.32×10^9 (d) 0.495 μCi

VP43.9.2 (a) $5.50 \times 10^{-6} \text{ s}^{-1}$ (b) 3.46×10^9 (c) 1.45×10^{10} (d) 72.5 h

VP43.9.3 (a) $\lambda = 0.0279 \text{ s}^{-1}$, $T_{1/2} = 24.8$ s (b) 5.63×10^5 initially, 1.60×10^5 after 45.0 s

VP43.9.4 (a) 2.67×10^3 y (b) 2.46×10^{10}

Bridging Problem

(a) ${}^{128}_{54}\text{Xe}$

(b) no; β^+ emission would be endoergic

(c) 3.25×10^9 atoms, 1.50×10^6 Bq

(d) $N(t) = (3.25 \times 10^9 \text{ atoms})(1 - e^{-(4.62 \times 10^{-4} \text{ s}^{-1})t})$