




? This image shows a portion of the Tarantula Nebula, a region some 170,000 light-years away where new stars are forming. The luminous stars, glowing gas, and opaque dust clouds are all made of “normal” matter—that is, atoms and their constituents. What percentage of the mass and energy in the universe is composed of “normal” matter? (i) 75% to 100%; (ii) 50% to 75%; (iii) 25% to 50%; (iv) 5% to 25%; (v) less than 5%.

44 Particle Physics and Cosmology

What are the most fundamental constituents of matter? How did the universe begin? And what is the fate of our universe? In this chapter we’ll explore what physicists and astronomers have learned in their quest to answer these questions.

The chapter title, “Particle Physics and Cosmology,” may seem strange. Fundamental particles are the *smallest* things in the universe, and cosmology deals with the *biggest* thing there is—the universe itself. Nonetheless, we’ll see in this chapter that physics on the most microscopic scale plays an essential role in determining the nature of the universe on the largest scale.

The development of high-energy accelerators and associated detectors has been crucial in our emerging understanding of particles. We can classify particles and their interactions in several ways in terms of conservation laws and symmetries, some of which are absolute and others of which are obeyed only in certain kinds of interactions. We’ll conclude by discussing our present understanding of the nature and evolution of the universe as a whole.

44.1 FUNDAMENTAL PARTICLES—A HISTORY

The idea that the world is made of fundamental particles has a long history. In about 400 B.C. the Greek philosophers Democritus and Leucippus suggested that matter is made of indivisible particles that they called *atoms*, a word derived from *a-* (not) and *tomos* (cut or divided). This idea lay dormant until about 1804, when the English scientist John Dalton (1766–1844), often called the father of modern chemistry, discovered that many chemical phenomena could be explained if atoms of each element are the basic, indivisible building blocks of matter.

The Electron and the Proton

Toward the end of the 19th century it became clear that atoms are *not* indivisible. The characteristic spectra of elements suggested that atoms have internal structure, and

LEARNING OUTCOMES

In this chapter, you'll learn...

- 44.1** The key varieties of fundamental subatomic particles and how they were discovered.
- 44.2** How physicists use accelerators and detectors to probe the properties of subatomic particles.
- 44.3** The four ways in which subatomic particles interact with each other.
- 44.4** How the structure of protons, neutrons, and other particles can be explained in terms of quarks.
- 44.5** The standard model of particles and interactions.
- 44.6** The evidence that the universe is expanding and that the expansion is speeding up.
- 44.7** The history of the first 380,000 years after the Big Bang.

You'll need to review...

- 13.3** Escape speed.
- 27.4** Motion of charged particles in a magnetic field.
- 32.1** Radiation from accelerated charges.
- 38.1, 38.3, 38.4** Photons; electron–positron annihilation; uncertainty principle.
- 39.1, 39.2** Electron waves; discovery of the nucleus.
- 41.5, 41.6** Electron spin; exclusion principle.
- 42.6** Valence bands and holes.
- 43.1, 43.3** Neutrons and protons; β^+ decay.

J. J. Thomson's discovery of the negatively charged *electron* in 1897 showed that atoms could be taken apart into charged particles. Rutherford's experiments in 1910–1911 (see Section 39.2) revealed that an atom's positive charge resides in a small, dense nucleus. In 1919 Rutherford made an additional discovery: When alpha particles are fired into nitrogen, one product is hydrogen gas. He reasoned that the hydrogen nucleus is a constituent of the nuclei of heavier atoms such as nitrogen, and that a collision with a fast-moving alpha particle can dislodge one of those hydrogen nuclei. Thus the hydrogen nucleus is an elementary particle that Rutherford named the *proton*. The following decade saw the blossoming of quantum mechanics, including the Schrödinger equation. Physicists were on their way to understanding the principles that underlie atomic structure.

The Photon

Einstein explained the photoelectric effect in 1905 by assuming that the energy of electromagnetic waves is quantized; that is, it comes in little bundles called *photons* with energy $E = hf$. Atoms and nuclei can emit (create) and absorb (destroy) photons (see Section 38.1). Considered as particles, photons have zero charge and zero rest mass. (Note that any discussions of a particle's mass in this chapter will refer to its rest mass.) In particle physics, a photon is denoted by the symbol γ (the Greek letter gamma).

The Neutron

In 1930 the German physicists Walther Bothe and Herbert Becker observed that when beryllium, boron, or lithium was bombarded by alpha particles, the target material emitted a radiation that had much greater penetrating power than the original alpha particles. Experiments by the English physicist James Chadwick in 1932 showed that the emitted particles were electrically neutral, with mass approximately equal to that of the proton. Chadwick christened these particles *neutrons* (symbol n or ${}_0^1n$). A typical reaction of the type studied by Bothe and Becker, with a beryllium target, is



Elementary particles are usually detected by their electromagnetic effects—for instance, by the ionization that they cause when they pass through matter. (This is the principle of the cloud chamber, described below.) Because neutrons have no charge, they are difficult to detect directly; they interact hardly at all with electrons and produce little ionization when they pass through matter. However, neutrons can be slowed down by scattering from nuclei, and they can penetrate a nucleus. Hence slow neutrons can be detected by means of a nuclear reaction in which a neutron is absorbed and an alpha particle is emitted. An example is



The ejected alpha particle is easy to detect because it is charged. Later experiments showed that neutrons and protons, like electrons, are spin- $\frac{1}{2}$ particles (see Section 43.1).

The discovery of the neutron cleared up a mystery about the composition of the nucleus. Before 1930 the mass of a nucleus was thought to be due only to protons, but no one understood why the charge-to-mass ratio was not the same for all nuclides. It soon became clear that all nuclides (except ${}_1^1\text{H}$) contain both protons and neutrons. Hence the proton, the neutron, and the electron are the building blocks of atoms. However, that is not the end of the particle story; these are not the only particles, and particles can do more than build atoms.

The Positron

The positive electron, or positron, was discovered by the American physicist Carl D. Anderson in 1932, during an investigation of particles bombarding the earth

from space. **Figure 44.1** shows a historic photograph made with a *cloud chamber*, an instrument used to visualize the tracks of charged particles. The chamber contained a supercooled vapor; a charged particle passing through the vapor causes ionization, and the ions trigger the condensation of liquid droplets from the vapor. The droplets make a visible track showing the charged particle's path.

The cloud chamber in Fig. 44.1 is in a magnetic field directed into the plane of the photograph. The particle has passed through a thin lead plate (which extends from left to right in the figure) that lies within the chamber. The track is more tightly curved above the plate than below it, showing that the speed was less above the plate than below it. Therefore the particle had to be moving upward; it could not have gained energy passing through the lead. The thickness and curvature of the track suggested that its mass and the magnitude of its charge equaled those of the electron. But the directions of the magnetic field and the velocity in the magnetic-force equation $\vec{F} = q\vec{v} \times \vec{B}$ showed that the particle had *positive* charge. Anderson christened this particle the *positron*.

To theorists, the appearance of the positron was a welcome development. In 1928 the English physicist Paul Dirac had developed a relativistic generalization of the Schrödinger equation for the electron. In Section 41.5 we discussed how Dirac's ideas helped explain the spin magnetic moment of the electron.

One of the puzzling features of the Dirac equation was that for a *free* electron it predicted not only a continuum of energy states greater than its rest energy $m_e c^2$, as expected, but also a continuum of *negative* energy states *less than* $-m_e c^2$ (**Fig. 44.2a**). That posed a problem. What was to prevent an electron from emitting a photon with energy $2m_e c^2$ or greater and hopping from a positive state to a negative state? It wasn't clear what these negative-energy states meant, and there was no obvious way to get rid of them. Dirac's ingenious interpretation was that all the negative-energy states were filled with electrons, and that these electrons were for some reason unobservable. The exclusion principle (see Section 41.6) would forbid a transition to a state that was already occupied.

A vacancy in a negative-energy state would act like a positive charge, just as a hole in the valence band of a semiconductor (see Section 42.6) acts like a positive charge. Initially, Dirac tried to argue that such vacancies were protons. But after Anderson's discovery it became clear that the vacancies were observed physically as *positrons*. Furthermore, the Dirac energy-state picture provides a mechanism for the *creation* of positrons. When an electron in a negative-energy state absorbs a photon with energy greater than $2m_e c^2$, it goes to a positive state (Fig. 44.2b), in which it becomes observable. The vacancy that it leaves behind is observed as a positron; the result is the creation of an electron–positron pair. Similarly, when an electron in a positive-energy state falls into a vacancy, both the electron and the vacancy (that is, the positron) disappear, and photons are emitted (Fig. 44.2c). Thus the Dirac theory leads naturally to the conclusion that, like photons, *electrons can be created and destroyed*. While photons can be created and destroyed singly, electrons can be produced or destroyed only in electron–positron pairs or in association with other particles. (Creating or destroying an electron alone would mean creating or destroying an amount of charge $-e$, which would violate the conservation of electric charge.)

In 1949 the American physicist Richard Feynman showed that a positron could be described mathematically as an electron traveling backward in time. His reformulation of the Dirac theory eliminated difficult calculations involving the infinite sea of negative-energy

Figure 44.1 Photograph of the cloud-chamber track made by the first positron ever identified. The photograph was made by Carl D. Anderson in 1932.

The positron follows a curved path owing to the presence of a magnetic field.

The track is more strongly curved above the lead plate, showing that the positron was traveling upward and lost energy and speed as it passed through the plate.







Figure 44.2 (a) Energy states for a free electron predicted by the Dirac equation. (b) Raising an electron from an $E < 0$ state to an $E > 0$ state corresponds to electron–positron pair production. (c) An electron dropping from an $E > 0$ state to a vacant $E < 0$ state corresponds to electron–positron pair annihilation.

Figure 44.3 (a) Photograph of bubble-chamber tracks of electron–positron pairs that are produced when 300 MeV photons strike a lead sheet. A magnetic field directed out of the photograph made the electrons and positrons curve in opposite directions. (b) Diagram showing the pair-production process for two of the photons.



BIO APPLICATION Pair Annihilation in Medical Diagnosis

A technique called positron emission tomography (PET) can be used to identify the early stages of Alzheimer's disease. A patient is administered a glucose-like compound called FDG in which one oxygen atom is replaced by radioactive ^{18}F . FDG accumulates in active areas of the brain, where glucose metabolism is high. The ^{18}F undergoes β^+ decay (positron emission) with a half-life of 110 minutes, and the emitted positron immediately annihilates with an atomic electron to produce two gamma-ray photons. A scanner detects both photons, then calculates where the annihilation took place—the site of FDG accumulation. These PET images—which show areas of strongest emission, and hence greatest glucose metabolism, in red—reveal changes in the brains of patients.



states and put electrons and positrons on the same footing. But the creation and destruction of electron–positron pairs remain. The Dirac theory provides the beginning of a theoretical framework for creation and destruction of all fundamental particles.

Experiment and theory tell us that the masses of the positron and electron are identical and that their charges are equal in magnitude but opposite in sign. The positron's spin angular momentum $\vec{S}$ and magnetic moment $\vec{\mu}$ are parallel; they are opposite for the electron. However, $\vec{S}$ and $\vec{\mu}$ have the same magnitude for both particles because they have the same spin. We use the term **antiparticle** for a particle that is related to another particle as the positron is to the electron. Each kind of particle has a corresponding antiparticle. For a few kinds of particles (necessarily all neutral) the particle and antiparticle are identical, and we can say that they are their own antiparticles. The photon is an example; there is no way to distinguish a photon from an antiphoton. We'll use the standard symbols e^- for the electron and e^+ for the positron. The generic term “electron” often includes both electrons and positrons. Other antiparticles may be denoted by a bar over the particle's symbol; for example, an antiproton is $\bar{p}$. We'll see other examples of antiparticles later.

Positrons do not occur in ordinary matter. Electron–positron pairs are produced during high-energy collisions of charged particles or γ rays with matter. This process is called e^+e^- pair production (Fig. 44.3). The minimum available energy required for electron–positron pair production equals the rest energy $2m_e c^2$ of the two particles:

$$\begin{aligned} E_{\min} &= 2m_e c^2 = 2(9.109 \times 10^{-31} \text{ kg})(2.998 \times 10^8 \text{ m/s})^2 \\ &= 1.637 \times 10^{-13} \text{ J} = 1.022 \text{ MeV} \end{aligned}$$

The inverse process, e^+e^- pair annihilation, occurs when a positron and an electron collide (see Example 38.6 in Section 38.3). Both particles disappear, and two (or occasionally three) photons can appear, with total energy of at least $2m_e c^2 = 1.022 \text{ MeV}$. Decay into a single photon is impossible: Such a process could not conserve both energy and momentum.

Positrons also occur in the decay of some unstable nuclei, in which they are called beta-plus particles (β^+). We discussed β^+ decay in Section 43.3.

We'll frequently represent particle masses in terms of the equivalent rest energy by using $m = E/c^2$. Then typical mass units are MeV/c^2 ; for example, $m = 0.511 \text{ MeV}/c^2$ for an electron or positron.

Particles as Force Mediators

In classical physics we describe the interaction of charged particles in terms of electric and magnetic forces. In quantum mechanics we can describe this interaction in terms of emission and absorption of photons. Two electrons repel each other as one emits a photon and the other absorbs it, just as two skaters can push each other apart by tossing a heavy ball back and forth between them (Fig. 44.4a). For an electron and a proton, in which the charges are opposite and the force is attractive, we imagine the skaters trying to grab the ball away from each other (Fig. 44.4b). The electromagnetic interaction between two charged particles is *mediated* or transmitted by photons.

If charged-particle interactions are mediated by photons, where does the energy to create the photons come from? Recall from our discussion of the uncertainty principle (see Sections 38.4 and 39.6) that a state that exists for a short time Δt has an uncertainty ΔE in its energy such that

$$\Delta E \Delta t \geq \frac{\hbar}{2} \quad (44.3)$$

This uncertainty permits the creation of a photon with energy ΔE , provided that it lives no longer than the time Δt given by Eq. (44.3). A photon that can exist for a short time because of this energy uncertainty is called a *virtual photon*. It's as though there were an energy bank; you can borrow energy, provided that you pay it back within the time limit. According to Eq. (44.3), the more you borrow, the sooner you have to pay it back.

Mesons

Is there a particle that mediates the *nuclear* force? By the mid-1930s the nuclear force between two nucleons (neutrons or protons) appeared to be described by a potential energy $U(r)$ with the general form

$$U(r) = -f^2 \left(\frac{e^{-r/r_0}}{r} \right) \quad (\text{nuclear potential energy}) \quad (44.4)$$

The constant f characterizes the strength of the interaction, and r_0 describes its range. **Figure 44.5** compares the absolute value of this function with the function f^2/r , which would be analogous to the *electric* interaction of two protons:

$$U(r) = \frac{1}{4\pi\epsilon_0} \frac{e^2}{r} \quad (\text{electric potential energy}) \quad (44.5)$$

In 1935 the Japanese physicist Hideki Yukawa suggested that a hypothetical particle that he called a **meson** might mediate the nuclear force. He showed that the range of the force was related to the mass of the particle. Yukawa argued that the particle must live for a time Δt long enough to travel a distance comparable to the range r_0 of the nuclear force. This range was known from the sizes of nuclei and other information to be about $1.5 \times 10^{-15} \text{ m} = 1.5 \text{ fm}$. If we assume that an average particle's speed is comparable to c and travels about half the range, its lifetime Δt must be about

$$\Delta t = \frac{r_0}{2c} = \frac{1.5 \times 10^{-15} \text{ m}}{2(3.0 \times 10^8 \text{ m/s})} = 2.5 \times 10^{-24} \text{ s}$$

From Eq. (44.3), the minimum necessary uncertainty ΔE in energy is

$$\Delta E = \frac{\hbar}{2\Delta t} = \frac{1.05 \times 10^{-34} \text{ J} \cdot \text{s}}{2(2.5 \times 10^{-24} \text{ s})} = 2.1 \times 10^{-11} \text{ J} = 130 \text{ MeV}$$

The mass equivalent Δm of this energy is about 250 times the electron mass:

$$\Delta m = \frac{\Delta E}{c^2} = \frac{2.1 \times 10^{-11} \text{ J}}{(3.00 \times 10^8 \text{ m/s})^2} = 2.3 \times 10^{-28} \text{ kg} = 130 \text{ MeV}/c^2$$


Yukawa postulated that an as yet undiscovered particle with this mass serves as the messenger for the nuclear force.

A year later, Carl Anderson and his colleague Seth Neddermeyer discovered in cosmic radiation two new particles, now called **muons**. The μ^- has charge equal to that of the electron, and its antiparticle the μ^+ has a positive charge with equal magnitude. The two particles have equal mass, about 207 times the electron mass. But it soon became clear that muons were *not* Yukawa's particles because they interacted with nuclei only very weakly.

In 1947 a family of three particles, called π *mesons* or **pions**, were discovered. Their charges are $+e$, $-e$, and zero, and their masses are about 270 times the electron mass. The pions interact strongly with nuclei, and they *are* the particles predicted by Yukawa. Other, heavier mesons, the ω and ρ , evidently also act as shorter-range messengers of the nuclear force. The complexity of this explanation suggests that the nuclear force has simpler underpinnings; these involve the quarks and gluons that we'll discuss in Section 44.4. Before discussing mesons further, we'll describe some particle accelerators and detectors to see how mesons and other particles are created in a controlled fashion and observed.

Figure 44.4 An analogy for how particles act as force mediators.

(a) Two skaters exert repulsive forces on each other by tossing a ball back and forth.



(b) Two skaters exert attractive forces on each other when one tries to grab the ball out of the other's hands.





Figure 44.5 Graph of the magnitude of the Yukawa potential-energy function for nuclear forces, $|U(r)| = f^2 e^{-r/r_0}/r$. The function $U(r) = f^2/r$, proportional to the potential energy for Coulomb's law, is also shown. The two functions are similar at small r , but the Yukawa potential energy drops off much more quickly at large r .



TEST YOUR UNDERSTANDING OF SECTION 44.1 Each of the following particles can be exchanged between two protons, two neutrons, or a neutron and a proton as part of the nuclear force. Rank the particles in order of the range of the interaction that they mediate, from largest to smallest range. (i) The π^+ (pi-plus) meson, mass $140 \text{ MeV}/c^2$; (ii) the ρ^+ (rho-plus) meson, mass $776 \text{ MeV}/c^2$; (iii) the η^0 (eta-zero) meson, mass $548 \text{ MeV}/c^2$; (iv) the ω^0 (omega-zero) meson, mass $783 \text{ MeV}/c^2$.

ANSWER (i), (iii), (ii), (iv). The more massive the virtual particle, the shorter its lifetime and the shorter the distance that it can travel during its lifetime.

44.2 PARTICLE ACCELERATORS AND DETECTORS

BIO APPLICATION Linear Accelerators in Medicine

Electron linear accelerators that provide a kinetic energy of 4–20 MeV are important tools in the treatment of many cancers. The electrons themselves are used to irradiate superficial tumors. Alternatively, the electrons can be directed at a metal target; then bremsstrahlung (see Section 38.2) produces x rays that are used to irradiate tumors that lie deeper inside the patient.



Early nuclear physicists used alpha and beta particles from naturally occurring radioactive elements for their experiments, but they were restricted in energy to the few MeV that are available in such random decays. Present-day particle accelerators can produce precisely controlled beams of particles, from electrons and positrons up to heavy ions, with a wide range of energies. These beams have three main uses. First, high-energy particles can collide to produce new particles, just as a collision of an electron and a positron can produce photons. Second, a high-energy particle has a short de Broglie wavelength and so can probe the small-scale interior structure of other particles, just as electron microscopes (see Section 39.1) can give better resolution than optical microscopes. Third, they can be used to produce nuclear reactions of scientific or medical use.

Linear Accelerators


Particle accelerators use electric and magnetic fields to accelerate and guide beams of charged particles. A *linear accelerator* (linac) accelerates particles in a straight line. J. J. Thomson's cathode-ray tubes were early examples of linacs. Modern linacs use a series of electrodes with gaps to give the particles a series of boosts. Most present-day high-energy linear accelerators use a traveling electromagnetic wave; the charged particles “ride” the wave in more or less the way that a surfer rides an incoming ocean wave. In the highest-energy linac in the world today, at the SLAC National Accelerator Laboratory, electrons and positrons can be accelerated to 50 GeV in a tube 3 km long. At this energy their de Broglie wavelengths are 0.025 fm, much smaller than the size of a proton or a neutron.

The Cyclotron


Many accelerators use magnets to deflect the charged particles into circular paths. The first was the *cyclotron*, invented in 1931 by E. O. Lawrence and M. Stanley Livingston at the University of California (Fig. 44.6a). Particles with mass m and charge q move inside a vacuum chamber in a uniform magnetic field $\vec{B}$ that is perpendicular to the plane of

Figure 44.6 Layout and operation of a cyclotron.


(a) Schematic diagram of a cyclotron



(b) As the positive particle reaches the gap, it is accelerated by the electric-field force ...



(c) ... and the next semicircular orbit has a larger radius.



By the time the particle reaches the gap again, the dee voltage has reversed and the particle is again accelerated.

their paths. In Section 27.4 we showed that in such a field, a particle with speed v moves in a circular path with radius r given by

$$r = \frac{mv}{|q|B} \quad (44.6)$$

and with angular speed (angular frequency) ω given by

$$\omega = \frac{v}{r} = \frac{|q|B}{m} \quad (44.7)$$

An alternating potential difference is applied between the two hollow electrodes D_1 and D_2 (called *dees*), creating an electric field in the gap between them. The polarity of the potential difference and electric field is changed precisely twice each revolution (Figs. 44.6b and 44.6c), so that the particles get a push each time they cross the gap. The pushes increase their speed and kinetic energy, boosting them into paths of larger radius. The maximum speed $v_{\max}$ and kinetic energy $K_{\max}$ are determined by the radius R of the largest possible path. Solving Eq. (44.6) for v , we find $v = |q|Br/m$ and $v_{\max} = |q|BR/m$. Assuming nonrelativistic speeds, we have

$$K_{\max} = \frac{1}{2}mv_{\max}^2 = \frac{q^2B^2R^2}{2m} \quad (44.8)$$

EXAMPLE 44.1 Frequency and energy in a proton cyclotron

WITH VARIATION PROBLEMS

One cyclotron built during the 1930s has a path of maximum radius 0.500 m and a magnetic field of magnitude 1.50 T. If it is used to accelerate protons, find (a) the frequency of the alternating voltage applied to the dees and (b) the maximum particle energy.

IDENTIFY and SET UP The frequency f of the applied voltage must equal the frequency of the proton orbital motion. Equation (44.7) gives the *angular* frequency ω of the proton orbital motion; we find f from $f = \omega/2\pi$. The proton reaches its maximum energy $K_{\max}$, given by Eq. (44.8), when the radius of its orbit equals the radius of the dees.

EXECUTE (a) For protons, $q = 1.60 \times 10^{-19}$ C and $m = 1.67 \times 10^{-27}$ kg. From Eq. (44.7),

$$\begin{aligned} f &= \frac{\omega}{2\pi} = \frac{|q|B}{2\pi m} = \frac{(1.60 \times 10^{-19} \text{ C})(1.50 \text{ T})}{2\pi(1.67 \times 10^{-27} \text{ kg})} \\ &= 2.3 \times 10^7 \text{ Hz} = 23 \text{ MHz} \end{aligned}$$

(b) From Eq. (44.8) the maximum kinetic energy is

$$\begin{aligned} K_{\max} &= \frac{(1.60 \times 10^{-19} \text{ C})^2(1.50 \text{ T})^2(0.50 \text{ m})^2}{2(1.67 \times 10^{-27} \text{ kg})} \\ &= 4.3 \times 10^{-12} \text{ J} = 2.7 \times 10^7 \text{ eV} = 27 \text{ MeV} \end{aligned}$$

This proton kinetic energy is much larger than that available from natural radioactive sources.

EVALUATE From Eq. (44.6) or Eq. (44.7), the proton speed is $v = 7.2 \times 10^7$ m/s, which is about 25% of the speed of light. At such speeds, relativistic effects are beginning to become important. Since we ignored these effects in our calculation, the results for f and $K_{\max}$ are in error by a few percent; this is why we kept only two significant figures.

KEYCONCEPT In the type of particle accelerator called a cyclotron, particles follow a circular path in a plane perpendicular to a uniform magnetic field. Every one-half orbit the particles pass through a potential difference that increases their kinetic energy and the radius of their circular path. The maximum radius of the cyclotron path determines the maximum kinetic energy attainable.


The maximum energy that can be attained with a cyclotron is limited by relativistic effects. The relativistic version of Eq. (44.7) is

$$\omega = \frac{|q|B}{m} \sqrt{1 - v^2/c^2}$$


As the particles speed up, their angular frequency ω *decreases*, and their motion gets out of phase with the alternating dee voltage. In the *synchrocyclotron* the particles are accelerated in bursts. For each burst, the frequency of the alternating voltage is decreased as the particles speed up, maintaining the correct phase relationship with the particles' motion.

Figure 44.7 (a) The Large Hadron Collider at the European Organization for Nuclear Research (CERN). The underground accelerating ring (shown by the red circle) is 100 m underground and 8.5 km in diameter, so large that it spans the border between Switzerland and France. (Note the Alps in the background.) When accelerated to 7 TeV, protons travel around the ring more than 11,000 times per second. (b) An engineer working on one of the 9593 superconducting electromagnets around the LHC ring.

(a)



(b)



Another limitation of the cyclotron is the difficulty of building very large electromagnets. The largest synchrocyclotron ever built has a vacuum chamber that is about 8 m in diameter and accelerates protons to energies of about 600 MeV.

The Synchrotron

To attain higher energies, a type of machine called the *synchrotron* is more practical. Particles move in a vacuum chamber in the form of a thin doughnut called the *accelerating ring*. The particle beam is bent to follow the ring by a series of electromagnets placed around the ring. As the particles speed up, the magnetic field is increased so that the particles retrace the same trajectory over and over. The Large Hadron Collider (LHC) near Geneva, Switzerland, is the highest-energy accelerator in the world (Fig. 44.7). It is designed to accelerate protons to a maximum energy of 7 TeV, or 7×10^{12} eV. (As we'll see in Section 44.3, *hadrons* are a class of elementary particles that includes protons and neutrons.)

As we pointed out in Section 32.1, accelerated charges radiate electromagnetic energy. In an accelerator in which the particles move in curved paths, this radiation is often called *synchrotron radiation*. High-energy accelerators are typically constructed underground to provide protection from this radiation. From the accelerator standpoint, synchrotron radiation is undesirable, since the energy given to an accelerated particle is radiated right back out. It can be minimized by making the accelerator radius r large so that the centripetal acceleration v^2/r is small. On the positive side, synchrotron radiation is used as a source of well-controlled high-frequency electromagnetic waves.

Available Energy

When a beam of high-energy particles collides with a stationary target, not all of the energy of the incident particles is *available* to form new particle states. Because momentum must be conserved, the particles emerging from the collision must have some net motion and thus some kinetic energy. The discussion following Example 43.11 (Section 43.6) presented a nonrelativistic example of this principle. The available energy is greatest in the frame of reference in which the total momentum is zero. We call this the *center-of-momentum system*; it is the relativistic generalization of the center-of-mass system that we discussed in Section 8.5. In this system the total kinetic energy after the collision can be zero, so that the maximum amount of the initial energy (the sum of the rest energies and kinetic energies of the colliding particles) becomes available to cause the reaction being studied.

Consider the *laboratory system*, in which a target particle with mass M is initially at rest and is bombarded by a particle with mass m and total energy (including rest energy) E_m . The total available energy E_a in the center-of-momentum system (including rest energies of all the particles) can be shown to be

$$E_a^2 = 2Mc^2E_m + (Mc^2)^2 + (mc^2)^2 \quad (\text{available energy}) \quad (44.9)$$

When the masses of the target and projectile particles are equal, we can simplify:

$$E_a^2 = 2mc^2(E_m + mc^2) \quad (\text{available energy, equal masses}) \quad (44.10)$$

If in addition E_m is much greater than mc^2 , we can ignore the second term in the parentheses in Eq (44.10). Then E_a is

$$E_a = \sqrt{2mc^2E_m} \quad (\text{available energy, equal masses, } E_m \gg mc^2) \quad (44.11)$$

The square root in Eq. (44.11) means that with a stationary target, doubling the energy E_m of the bombarding particle increases the available energy E_a by only a factor of $\sqrt{2} = 1.414$. Examples 44.2 and 44.3 explore the limitations of having a stationary target particle.

CAUTION Available energy includes rest energies Note that the available energy given by Eqs. (44.9), (44.10), and (44.11) is the energy that goes into *all* of the particles that are present after the collision. If the initial particles survive the collision (see Example 44.2), some of the available energy goes back to these survivors; the rest goes to whatever additional particles are produced in the collision. ■

EXAMPLE 44.2 Threshold energy for pion production**WITH VARIATION PROBLEMS**

A proton (rest energy 938 MeV) with kinetic energy K collides with a proton at rest. Both protons survive the collision, and a neutral pion (π^0 , rest energy 135 MeV) is produced. What is the threshold energy (minimum value of K) for this process?

IDENTIFY and SET UP The final state includes the two original protons (mass m) and the pion (mass m_π). The threshold energy corresponds to the minimum-energy case in which all three particles are at rest in the center-of-momentum system. The total available energy E_a in that system must be at least the total rest energy, $2mc^2 + m_\pi c^2$. We use this to solve Eq. (44.10) for the total energy E_m of the bombarding proton; the kinetic energy K (our target variable) is then E_m minus the proton rest energy mc^2 .

EXECUTE We substitute $E_a = 2mc^2 + m_\pi c^2$ into Eq. (44.10), simplify, and solve for E_m :

$$4m^2c^4 + 4mm_\pi c^4 + m_\pi^2 c^4 = 2mc^2 E_m + 2(mc^2)^2$$

$$E_m = mc^2 + m_\pi c^2 \left(2 + \frac{m_\pi c^2}{2mc^2} \right) = mc^2 + K$$

$$K = m_\pi c^2 \left(2 + \frac{m_\pi c^2}{2mc^2} \right) = (135 \text{ MeV}) \left(2 + \frac{(135 \text{ MeV})}{2(938 \text{ MeV})} \right) = 280 \text{ MeV}$$

EVALUATE The bombarding proton's kinetic energy must be more than twice the pion rest energy of 135 MeV. By contrast, in Example 37.11 (Section 37.8) we found that a pion can be produced in a head-on collision of two protons, each with only 67.5 MeV of kinetic energy. We discuss the energy advantage of such collisions in the next subsection.

KEYCONCEPT The available energy E_a in a particle collision is the energy available to produce the particles that are present after the collision. This equals the total energy of the colliding particles in the center-of-momentum system, a frame in which the total momentum of the particles is zero. Given E_a , you can calculate the required kinetic energy in a different frame in which one of the particles is initially at rest.

EXAMPLE 44.3 Increasing the available energy**WITH VARIATION PROBLEMS**

The Fermilab accelerator in Illinois was designed to bombard stationary targets with 800 GeV protons. (a) What is the available energy E_a in a proton-proton collision? (b) What is E_a if the beam energy is increased to 980 GeV?

IDENTIFY and SET UP Our target variable is the available energy E_a in a stationary-target collision between identical particles. In both parts (a) and (b) the beam energy E_m is much larger than the proton rest energy $mc^2 = 938 \text{ MeV} = 0.938 \text{ GeV}$, so we can safely use the approximation of Eq. (44.11).

EXECUTE (a) For $E_m = 800 \text{ GeV}$, Eq. (44.11) gives

$$E_a = \sqrt{2(0.938 \text{ GeV})(800 \text{ GeV})} = 38.7 \text{ GeV}$$

(b) For $E_m = 980 \text{ GeV}$,

$$E_a = \sqrt{2(0.938 \text{ GeV})(980 \text{ GeV})} = 42.9 \text{ GeV}$$

EVALUATE With a stationary-proton target, increasing the proton beam energy by 180 GeV increases the available energy by only 4.2 GeV! This shows a major limitation of experiments in which one of the colliding particles is initially at rest. Below we describe how physicists can overcome this limitation.


KEYCONCEPT In experiments in which a fast-moving particle strikes a second target particle that is initially stationary, a large increase in the energy of the moving particle produces only a relatively small increase in the available energy.

Colliding Beams

The limitation illustrated by Example 44.3 doesn't apply to *colliding-beam* experiments. In these experiments there is no stationary target; instead, beams of particles moving in opposite directions are tightly focused onto one another so that head-on collisions can occur. Usually the two colliding particles have momenta of equal magnitude and opposite direction, so the total momentum is zero. Hence the laboratory system is also the center-of-momentum system, and the available energy is maximized.

The highest-energy colliding beams available are those at the Large Hadron Collider (see Fig. 44.7). In operation, 2808 bunches of 7 TeV protons circulate around the ring, half in one direction and half in the opposite direction. Each bunch contains about 10^{11} protons. Magnets steer the oppositely moving bunches to collide at interaction points. The available energy E_a in the resulting head-on collisions is the total energy of the two colliding particles: $E_a = 2 \times 7 \text{ TeV} = 14 \text{ TeV}$. The very large available energy at the Large Hadron Collider makes it possible to produce particles that have never been seen before (see Section 44.5).

Figure 44.8 This computer-generated image shows the result of a simulated collision between two protons (not shown) in one of the interaction regions at the Large Hadron Collider. The view is along the beampipe. The different color tracks show different types of particles emerging from the collision. A variety of different detectors surround the collision region. (Note the woman in a red dress, drawn for scale.)



Detectors

A wide variety of devices have been designed to measure the properties of subatomic particles. Many detectors use the ionization caused by charged particles as they move through a gas, liquid, or solid. The ions along the particle's path give rise to droplets of liquid in the supersaturated vapor of a cloud chamber (Fig. 44.1) or cause small volumes of vapor in the superheated liquid of a bubble chamber (Fig. 44.3a). In a semiconducting solid the ionization can take the form of electron–hole pairs. We discussed their detection in Section 42.7. *Wire chambers* contain arrays of closely spaced wires that detect the ions. The charge collected and time information from each wire are processed by using computers to reconstruct the particle trajectories. The detectors at the Large Hadron Collider use an array of devices to follow the tracks of particles produced by collisions between protons (Fig. 44.8). The giant solenoid in the photo that opens Chapter 28 is at the heart of one these detector arrays. The intense magnetic field of the solenoid helps identify newly produced particles, which curve in different directions and along paths of different radii depending on their charge and energy.

Cosmic-Ray Experiments

Large numbers of particles called *cosmic rays* continually bombard the earth from sources both within and beyond our galaxy. These particles consist mostly of neutrinos, protons, and heavier nuclei, with energies ranging from less than 1 MeV to more than 10^{20} eV. The earth's atmosphere and magnetic field protect us from much of this radiation. This means that cosmic-ray experimentation often must be carried out above all or most of the atmosphere by means of rockets or high-altitude balloons.

In contrast, neutrino detectors are buried below the earth's surface in tunnels or mines or submerged deep in the ocean. This is done to screen out all other types of particles so that only neutrinos, which interact only very weakly with matter, reach the detector. Because neutrino interactions with matter are so weak, neutrino detectors must consist of huge amounts of matter: The Super-Kamiokande detector looks for flashes of light produced when a neutrino interacts in a tank containing 5×10^7 kg of water (see Section 44.5).

Cosmic rays were important in early particle physics, and their study currently brings us important information about the rest of the universe. Although cosmic rays provide a source of high-energy particles that does not depend on expensive accelerators, most particle physicists use accelerators because the high-energy cosmic-ray particles they want are too few and too random.

TEST YOUR UNDERSTANDING OF SECTION 44.2 In a colliding-beam experiment, a 90 GeV electron collides head-on with a 90 GeV positron. The electron and the positron annihilate each other, forming a single virtual photon that then transforms into other particles. Does the virtual photon obey the same relationship $E = pc$ as real photons do?

ANSWER In a head-on collision between an electron and a positron of equal energy, the net momentum is zero. Since both momentum and energy are conserved in the collision, the virtual photon also has momentum $p = 0$ but has energy $E = 90 \text{ GeV} + 90 \text{ GeV} = 180 \text{ GeV}$. Hence the relationship $E = pc$ is definitely *not* true for this virtual photon.

44.3 PARTICLES AND INTERACTIONS

We have mentioned the array of subatomic particles that were known as of 1947: photons, electrons, positrons, protons, neutrons, muons, and pions. Since then, literally hundreds of additional particles have been discovered in accelerator experiments. The vast majority of known particles are *unstable* and decay spontaneously into other particles. Particles of all kinds, whether stable or unstable, can be created or destroyed in interactions between particles. Each such interaction involves the exchange of virtual particles, which exist on borrowed energy allowed by the uncertainty principle.

Although the world of subatomic particles and their interactions is complex, some key results bring order and simplicity to the seeming chaos. One key simplification is that there are only four fundamental types of interactions, each mediated or transmitted by the exchange of certain characteristic virtual particles. Furthermore, not all particles respond to all four kinds of interaction. In this section we'll examine the fundamental interactions more closely and see how physicists classify particles in terms of the ways in which they interact.

Four Forces and Their Mediating Particles

In Section 5.5 we first described the four fundamental types of forces or interactions (Fig. 44.9). They are, in order of decreasing strength:

1. The strong interaction
2. The electromagnetic interaction
3. The weak interaction
4. The gravitational interaction

The *electromagnetic* and *gravitational* interactions are familiar from classical physics. Both are characterized by a $1/r^2$ dependence on distance. In this scheme, the mediating particles for both interactions have mass zero and are stable as ordinary particles. The mediating particle for the electromagnetic interaction is the familiar photon, which has spin 1. (That means its spin quantum number is $s = 1$, so the magnitude of its spin angular momentum is $S = \sqrt{s(s+1)}\hbar = \sqrt{2}\hbar$.) The mediating particle for the gravitational force is the spin-2 *graviton* ($s = 2$, $S = \sqrt{s(s+1)}\hbar = \sqrt{6}\hbar$). The graviton has not yet been observed experimentally because the gravitational force is very much weaker than the electromagnetic force. For example, the gravitational attraction of two protons is smaller than their electrical repulsion by a factor of about 10^{36} . The gravitational force is of primary importance in the structure of stars and the large-scale behavior of the universe, but it is not believed to play a significant role in particle interactions at the energies that are currently attainable.

The other two forces are less familiar. One, usually called the *strong interaction*, is responsible for the nuclear force and also for the production of pions and several other particles in high-energy collisions. At the most fundamental level, the mediating particle for the strong interaction is called a *gluon*. However, the force between nucleons is more easily described in terms of mesons as the mediating particles. We'll discuss the spin-1, massless gluon in Section 44.4.

Equation (44.4) is a possible potential-energy function for the nuclear force. The strength of the interaction is described by the constant f^2 , which has units of energy times distance. A better basis for comparison with other forces is the dimensionless ratio $f^2/\hbar c$, called the *coupling constant* for the interaction. (We invite you to verify that this ratio is a pure number and so must have the same value in all systems of units.) The observed behavior of nuclear forces suggests that $f^2/\hbar c \approx 1$. The dimensionless coupling constant for *electromagnetic* interactions is the fine-structure constant, which we introduced in Section 41.5:

$$\frac{1}{4\pi\epsilon_0} \frac{e^2}{\hbar c} = 7.2974 \times 10^{-3} = \frac{1}{137.04} \quad (44.12)$$

Thus the strong interaction is roughly 100 times as strong as the electromagnetic interaction; however, it drops off with distance more quickly than $1/r^2$.

The fourth interaction is called the *weak* interaction. It is responsible for beta decay, such as the conversion of a neutron into a proton, an electron, and an antineutrino. It is also responsible for the decay of many unstable particles (pions into muons, muons into electrons, and so on). Its mediating particles are the short-lived particles W^+ , W^- , and Z^0 . The existence of these particles was confirmed in 1983 in experiments at CERN, for which Carlo Rubbia and Simon van der Meer were awarded the Nobel Prize in 1984.

Figure 44.9 The ties that bind us together originate in the fundamental interactions of nature. The nuclei within our bodies are held together by the strong interaction. The electromagnetic interaction binds nuclei and electrons together to form atoms, binds atoms together to form molecules, and binds molecules together to form us.



TABLE 44.1 Four Fundamental Interactions

Interaction	Relative Strength	Range	Mediating Particle			
			Name	Mass	Charge	Spin
Strong	1	Short (~ 1 fm)	Gluon	0	0	1
Electromagnetic	$\frac{1}{137.04}$	Long ($1/r^2$)	Photon	0	0	1
Weak	10^{-9}	Short (~ 0.001 fm)	$W^\pm, Z^0$	80.4, 91.2 GeV/c^2	$\pm e, 0$	1
Gravitational	10^{-38}	Long ($1/r^2$)	Graviton	0	0	2

The $W^\pm$ and Z^0 have spin 1 like the photon and the gluon, but they are *not* massless. In fact, they have enormous masses, $80.4 \text{ GeV}/c^2$ for the W 's and $91.2 \text{ GeV}/c^2$ for the Z^0 . With such massive mediating particles the weak interaction has a much shorter range than the strong interaction. It also lives up to its name by being weaker than the strong interaction by a factor of about 10^9 .

Table 44.1 compares the main features of these four fundamental interactions.

More Particles

In Section 44.1 we mentioned the discoveries of muons in 1937 and of pions in 1947. The electric charges of the muons and the charged pions have the same magnitude e as the electron charge. The positive muon μ^+ is the antiparticle of the negative muon μ^- . Each has spin $\frac{1}{2}$, like the electron, and a mass of about $207m_e = 106 \text{ MeV}/c^2$. Muons are unstable; each decays with a lifetime of $2.2 \times 10^{-6} \text{ s}$ into an electron of the same sign, a neutrino, and an antineutrino.

There are three kinds of pions, all with spin 0; they have *no* spin angular momentum. The π^+ and π^- have masses of $273m_e = 140 \text{ MeV}/c^2$. They are unstable; each $\pi^\pm$ decays with a lifetime of $2.6 \times 10^{-8} \text{ s}$ into a muon of the same sign along with a neutrino for the π^+ and an antineutrino for the π^- . The π^0 is somewhat less massive, $264m_e = 135 \text{ MeV}/c^2$, and it decays with a lifetime of $8.4 \times 10^{-17} \text{ s}$ into two photons. The π^+ and π^- are antiparticles of one another, while the π^0 is its own antiparticle. (That is, there is no distinction between particle and antiparticle for the π^0 .)

The existence of the *antiproton* $\bar{p}$ had been suspected ever since the discovery of the positron. The $\bar{p}$ was found in 1955, when proton–antiproton ($p\bar{p}$) pairs were created by use of a beam of 6 GeV protons from the Bevatron at the University of California, Berkeley. The *antineutron* $\bar{n}$ was found soon afterward. After 1960, as higher-energy accelerators and more sophisticated detectors were developed, a veritable blizzard of new unstable particles were identified. To describe and classify them, we need a small blizzard of new terms.

Initially, particles were classified by mass into three categories: (1) leptons (“light ones” such as electrons); (2) mesons (“intermediate ones” such as pions); and (3) baryons (“heavy ones” such as nucleons and more massive particles). But this scheme has been superseded by a more useful one in which particles are classified in terms of their *interactions*. For instance, *hadrons* (which include mesons and baryons) have strong interactions, and *leptons* do not.

In the following discussion we’ll also distinguish between **fermions**, which have half-integer spins, and **bosons**, which have zero or integer spins. Fermions obey the exclusion principle, on which the Fermi–Dirac distribution function (see Section 42.5) is based. Bosons do not obey the exclusion principle (there is no limit on how many bosons can occupy the same quantum state) and have a different distribution function, the Bose–Einstein distribution.

TABLE 44.2 The Six Leptons

Particle Name	Symbol	Anti-particle	Mass (MeV/c ²)	L_e	L_μ	L_τ	Lifetime (s)	Principal Decay Modes
Electron	e^-	e^+	0.511	+1	0	0	Stable	
Electron neutrino	ν_e	$\bar{\nu}_e$	$<2 \times 10^{-6}$	+1	0	0	Stable	
Muon	μ^-	μ^+	105.7	0	+1	0	2.20×10^{-6}	$e^- \bar{\nu}_e \nu_\mu$
Muon neutrino	ν_μ	$\bar{\nu}_\mu$	<0.19	0	+1	0	Stable	
Tau	τ^-	τ^+	1777	0	0	+1	2.9×10^{-13}	$\mu^- \bar{\nu}_\mu \nu_\tau$ or $e^- \bar{\nu}_e \nu_\tau$
Tau neutrino	ν_τ	$\bar{\nu}_\tau$	<18.2	0	0	+1	Stable	

Note: In addition to the limits on the individual neutrino masses, there is a much more stringent limit on the *sum* of the masses of the three types of neutrinos. Evidence suggests that this sum is less than about $3 \times 10^{-7} \text{ MeV}/c^2 = 0.3 \text{ eV}/c^2$.

Leptons

The **leptons**, which do not have strong interactions, include six particles: the electron (e^-) and its neutrino (ν_e), the muon (μ^-) and its neutrino (ν_μ), and the tau particle (τ^-) and its neutrino (ν_τ). Each of these has a distinct antiparticle. All leptons have spin $\frac{1}{2}$ and thus are fermions. **Table 44.2** shows the family of leptons. The taus have mass $3478m_e = 1777 \text{ MeV}/c^2$. Taus and muons are unstable; a τ^- decays into a μ^- plus a tau neutrino and a muon antineutrino, or an electron plus a tau neutrino and an electron antineutrino. A μ^- decays into an electron plus a muon neutrino and an electron antineutrino. They have relatively long lifetimes because their decays are mediated by the weak interaction. Despite their zero charge, a neutrino is distinct from an antineutrino; the spin angular momentum of a neutrino has a component that is opposite its linear momentum, while for an antineutrino that component is parallel to its linear momentum. Because neutrinos are so elusive, physicists have only been able to place upper limits on the rest masses of the ν_e , the ν_μ , and the ν_τ . It was thought that the rest masses of the neutrinos were zero; compelling recent evidence indicates that they have small but nonzero masses. We'll return to this point and its implications later.

Leptons obey a *conservation principle*. Corresponding to the three pairs of leptons are three lepton numbers L_e , L_μ , and L_τ . The electron e^- and the electron neutrino ν_e are assigned $L_e = 1$, and their antiparticles e^+ and $\bar{\nu}_e$ are given $L_e = -1$. Corresponding assignments of L_μ and L_τ are made for the μ and τ particles and their neutrinos.

In all interactions, each lepton number is separately conserved.

For example, in the decay of the μ^- , the lepton numbers are

$$\begin{array}{ccccccc} \mu^- & \rightarrow & e^- & + & \bar{\nu}_e & + & \nu_\mu \\ L_\mu = 1 & & L_e = 1 & & L_e = -1 & & L_\mu = 1 \end{array}$$

These conservation principles have no counterpart in classical physics.

EXAMPLE 44.4 Lepton number conservation

Check conservation of lepton numbers for these decay schemes:

- $\mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu$
- $\pi^- \rightarrow \mu^- + \bar{\nu}_\mu$
- $\pi^0 \rightarrow \mu^- + e^+ + \nu_e$

IDENTIFY and SET UP Lepton number conservation requires that L_e , L_μ , and L_τ (given in Table 44.2) separately have the same sums after the decay as before.

EXECUTE We tabulate L_e and L_μ for each decay scheme. An antiparticle has the opposite lepton number from its corresponding particle listed in

Table 44.2. No τ particles or τ neutrinos appear in any of the schemes, so $L_\tau = 0$ both before and after each decay and L_τ is conserved.

- $\mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu$
 $L_e: 0 = -1 + 1 + 0$
 $L_\mu: -1 = 0 + 0 + (-1)$
- $\pi^- \rightarrow \mu^- + \bar{\nu}_\mu$
 $L_e: 0 = 0 + 0$
 $L_\mu: 0 = 1 + (-1)$

Continued

(c) $\pi^0 \rightarrow \mu^- + e^+ + \nu_e$
 $L_e: 0 = 0 + (-1) + 1$
 $L_\mu: 0 \neq 1 + 0 + 0$

EVALUATE Decays (a) and (b) are consistent with lepton number conservation and are observed. Decay (c) violates the conservation of L_μ and has *never* been observed. Physicists used these and other

experimental results to deduce the principle that all three lepton numbers must separately be conserved.

KEYCONCEPT There are three types of lepton number: electron lepton number L_e , muon lepton number L_μ , and tau lepton number L_τ . Each of these is separately conserved in particle interactions; no violations of these conservation laws have ever been observed.

Hadrons

Hadrons, the strongly interacting particles, are a more complex family than leptons. Each hadron has an antiparticle, often denoted with an overbar, as with the antiproton $\bar{p}$. There are two subclasses of hadrons: *mesons* and *baryons*. **Table 44.3** shows some of the many hadrons that are currently known. (We'll explain *strangeness* and *quark content* later in this section and in the next one.)

Mesons include the pions that have already been mentioned, K mesons or *kaons*, η mesons, and others that we'll discuss later. Mesons have spin 0 or 1 and therefore are all bosons. There are no stable mesons; all mesons decay to less massive particles, obeying all the conservation laws for such decays.

Baryons include the nucleons and several particles called *hyperons*, including the Λ , Σ , Ξ , and Ω . These resemble nucleons but are more massive. Baryons have half-integer spin, and therefore all are fermions. The only stable baryon is the proton; a free neutron decays to a proton, and hyperons decay to other hyperons or to nucleons by various processes. Baryons obey the *conservation of baryon number*, analogous to conservation of lepton numbers, again with no counterpart in classical physics. We assign a baryon number $B = 1$ to each baryon (p, n, Λ , Σ , and so on) and $B = -1$ to each antibaryon ($\bar{p}$, $\bar{n}$, $\bar{\Lambda}$, $\bar{\Sigma}$, and so on).

In all interactions, the total baryon number is conserved.

This principle is the reason the mass number A was conserved in all of the nuclear reactions that we studied in Chapter 43.

TABLE 44.3 Some Hadrons and Their Properties

Particle	Mass (MeV/c ²)	Charge Ratio, Q/e	Spin	Baryon Number, B	Strangeness, S	Mean Lifetime (s)	Typical Decay Modes	Quark Content
<i>Mesons</i>								
π^0	135.0	0	0	0	0	8.5×10^{-17}	$\gamma \gamma$	$u\bar{u}, d\bar{d}$
π^+	139.6	+1	0	0	0	2.60×10^{-8}	$\mu^+ \nu_\mu$	$u\bar{d}$
π^-	139.6	-1	0	0	0	2.60×10^{-8}	$\mu^- \bar{\nu}_\mu$	$\bar{u}d$
K^+	493.7	+1	0	0	+1	1.24×10^{-8}	$\mu^+ \nu_\mu$	$u\bar{s}$
K^-	493.7	-1	0	0	-1	1.24×10^{-8}	$\mu^- \bar{\nu}_\mu$	$\bar{u}s$
η^0	547.9	0	0	0	0	$\approx 10^{-18}$	$\gamma \gamma$	$u\bar{u}, d\bar{d}, s\bar{s}$
<i>Baryons</i>								
p	938.3	+1	$\frac{1}{2}$	1	0	Stable	—	uud
n	939.6	0	$\frac{1}{2}$	1	0	880	$p e^- \bar{\nu}_e$	udd
Λ^0	1116	0	$\frac{1}{2}$	1	-1	2.63×10^{-10}	$p \pi^-$ or $n \pi^0$	uds
Σ^+	1189	+1	$\frac{1}{2}$	1	-1	8.02×10^{-11}	$p \pi^0$ or $n \pi^+$	uus
Σ^0	1193	0	$\frac{1}{2}$	1	-1	7.4×10^{-20}	$\Lambda^0 \gamma$	uds
Σ^-	1197	-1	$\frac{1}{2}$	1	-1	1.48×10^{-10}	$n \pi^-$	dds
Ξ^0	1315	0	$\frac{1}{2}$	1	-2	2.90×10^{-10}	$\Lambda^0 \pi^0$	uss
Ξ^-	1322	-1	$\frac{1}{2}$	1	-2	1.64×10^{-10}	$\Lambda^0 \pi^-$	dss
Δ^{++}	1232	+2	$\frac{3}{2}$	1	0	$\approx 10^{-23}$	$p \pi^+$	uuu
Ω^-	1672	-1	$\frac{3}{2}$	1	-3	8.2×10^{-11}	$\Lambda^0 K^-$	sss
Λ_c^+	2286	+1	$\frac{1}{2}$	1	0	2.0×10^{-13}	$p K^- \pi^+$	udc

EXAMPLE 44.5 Baryon number conservation

Check conservation of baryon number for these reactions:

- (a) $n + p \rightarrow n + p + p + \bar{p}$
 (b) $n + p \rightarrow n + p + \bar{n}$

IDENTIFY and SET UP This example is similar to Example 44.4. We compare the total baryon number before and after each reaction, using data from Table 44.3.

EXECUTE We tabulate the baryon numbers, noting that a baryon has $B = 1$ and an antibaryon has $B = -1$:

- (a) $n + p \rightarrow n + p + p + \bar{p}$: $1 + 1 = 1 + 1 + 1 + (-1)$
 (b) $n + p \rightarrow n + p + \bar{n}$: $1 + 1 \neq 1 + 1 + (-1)$

EVALUATE Reaction (a) is consistent with baryon number conservation. It can occur if enough energy is available in the $n + p$ collision. Reaction (b) violates baryon number conservation and has never been observed.


KEYCONCEPT Baryon number, like the three lepton numbers, is always conserved in particle interactions.

EXAMPLE 44.6 Antiproton creation

What is the minimum proton energy required to produce an antiproton in a collision with a stationary proton?

IDENTIFY and SET UP The reaction must conserve baryon number, charge, and energy. Since the target and bombarding protons are of equal mass and the target is at rest, we determine the minimum energy E_m of the bombarding proton from Eq. (44.10).

EXECUTE Conservation of charge and conservation of baryon number forbid the creation of an antiproton by itself; it must be created as part of a proton–antiproton pair. The complete reaction is



For this reaction to occur, the minimum available energy E_a in Eq. (44.10) is the final rest energy $4mc^2$ of three protons and an antiproton. Equation (44.10) then gives

$$(4mc^2)^2 = 2mc^2(E_m + mc^2)$$

$$E_m = 7mc^2$$

EVALUATE The energy E_m of the bombarding proton includes its rest energy mc^2 , so its minimum *kinetic* energy must be $6mc^2 = 6(938 \text{ MeV}) = 5.63 \text{ GeV}$.


The search for the antiproton was a principal reason for the construction of the Bevatron at the University of California, Berkeley, with beam energy of 6 GeV. The search succeeded in 1955, and Emilio Segrè and Owen Chamberlain were later awarded the Nobel Prize for this discovery.

KEYCONCEPT To calculate the minimum energy required to produce a new particle, you must first consider the conservation laws associated with the particle. These laws will tell you which other particles may have to be produced along with the desired one. Once you have done this, you can determine how much available energy is required.


Strangeness

The K mesons and the Λ and Σ hyperons were discovered during the late 1950s. Because of their unusual behavior they were called *strange particles*. They were produced in high-energy collisions such as $\pi^- + p$, and a K meson and a hyperon were always produced *together*. The relatively high rate of production of these particles suggested that it was a *strong*-interaction process, but their relatively long lifetimes suggested that their decay was a *weak*-interaction process. The K^0 appeared to have *two* lifetimes, one about $9 \times 10^{-11} \text{ s}$ and another nearly 600 times longer. Were the K mesons strongly interacting hadrons or not?

The search for the answer to this question led physicists to introduce a new quantity called **strangeness**. The hyperons Λ^0 and $\Sigma^{\pm,0}$ were assigned a strangeness quantum number $S = -1$, and the associated K^0 and K^+ mesons were assigned $S = +1$. The corresponding antiparticles had opposite strangeness, $S = +1$ for $\bar{\Lambda}^0$ and $\bar{\Sigma}^{\pm,0}$ and $S = -1$ for $\bar{K}^0$ and K^- . Then strangeness was *conserved* in production processes such as




The process



does not conserve strangeness and it does not occur.

When strange particles decay individually, strangeness is usually *not* conserved. Typical processes include



In each of these decays, the initial strangeness is 1 or -1 , and the final value is zero. All observations of these particles are consistent with the conclusion that *strangeness is conserved in strong interactions but it can change by zero or one unit in weak interactions*. There is no counterpart to the strangeness quantum number in classical physics.

CAUTION **Strangeness vs. spin** Take care not to confuse the symbol S for strangeness with the identical symbol for the magnitude of the spin angular momentum. ■

Conservation Laws

The decay of strange particles provides our first example of a *conditional conservation law*, one that is obeyed in some interactions and not in others. By contrast, several conservation laws are obeyed in *all* interactions. These include the familiar conservation laws: energy, momentum, angular momentum, and electric charge. These are called *absolute conservation laws*. Baryon number and the three lepton numbers are also conserved in all interactions. Strangeness is conserved in strong and electromagnetic interactions but *not* in all weak interactions.

Two other quantities, which are conserved in some but not all interactions, are useful in classifying particles and their interactions. One is *isospin*, a quantity that is used to describe the charge independence of the strong interactions. The other is *parity*, which describes the comparative behavior of two systems that are mirror images of each other. Isospin is conserved in strong interactions, which are charge independent, but not in electromagnetic or weak interactions. (The electromagnetic interaction is certainly *not* charge independent.) Parity is conserved in strong and electromagnetic interactions but not in weak ones. The Chinese-American physicists T. D. Lee and C. N. Yang received the Nobel Prize in 1957 for laying the theoretical foundations for nonconservation of parity in weak interactions.

This discussion shows that conservation laws provide another basis for classifying particles and their interactions. Each conservation law is also associated with a *symmetry* property of the system. A familiar example is angular momentum. If a system is in an environment that has spherical symmetry, no torque can act on it because the direction of the torque would violate the symmetry. In such a system, total angular momentum is *conserved*. When a conservation law is violated, the interaction may be described as a *symmetry-breaking interaction*.

TEST YOUR UNDERSTANDING OF SECTION 44.3 From conservation of energy, a particle of mass m and rest energy mc^2 can decay only if the decay products have a total mass less than m . (The remaining energy goes into the kinetic energy of the decay products.) Can a proton decay into less massive mesons?

ANSWER Mesons all have baryon number $B = 0$, while a proton has $B = 1$. The decay of a proton into one or more mesons would require that baryon number *not* be conserved. No violation of this conservation principle has ever been observed, so the proposed decay is impossible.

44.4 QUARKS AND GLUONS

The leptons form a fairly neat package: three particles and three neutrinos, each with its antiparticle, and a conservation law relating their numbers. Physicists believe that leptons are genuinely fundamental particles. The hadron family, by comparison, is a mess. Table 44.3 (in Section 44.3) contains only a sample of well over 100 hadrons that have been discovered since 1960, and it has become clear that these particles *do not* represent the most fundamental level of the structure of matter.

Our present understanding of the structure of hadrons is based on a proposal made initially in 1964 by the American physicist Murray Gell-Mann and his collaborators. In this proposal, hadrons are not fundamental particles but are composite structures whose constituents are spin- $\frac{1}{2}$ fermions called **quarks**. (The name is from the line “Three quarks for Muster Mark!” from *Finnegans Wake*, by James Joyce.) Each baryon is composed of three quarks (qqq), each antibaryon of three antiquarks ($\bar{q}\bar{q}\bar{q}$), and each meson of a quark–antiquark pair ($q\bar{q}$). Table 44.3 gives the quark content of many hadrons. No other compositions seem to be necessary. This scheme requires that quarks have electric charges with magnitudes $\frac{1}{3}$ and $\frac{2}{3}$ of the electron charge e , which had been thought to be the smallest unit of charge. Each quark also has a fractional value $\frac{1}{3}$ for its baryon number B , and each antiquark has a baryon-number value $-\frac{1}{3}$. In a meson, a quark and antiquark combine with net baryon number 0 and can have their spin angular momentum components parallel to form a spin-1 meson or antiparallel to form a spin-0 meson. Similarly, the three quarks in a baryon combine with net baryon number 1 and can form a spin- $\frac{1}{2}$ baryon or a spin- $\frac{3}{2}$ baryon.


The Three Original Quarks

The first (1964) quark theory included three types (called *flavors*) of quarks, labeled **u** (up), **d** (down), and **s** (strange). Their principal properties are listed in **Table 44.4**. The corresponding antiquarks $\bar{u}$, $\bar{d}$, and $\bar{s}$ have opposite values of charge Q , B , and S . Protons, neutrons, π and K mesons, and several hyperons can be constructed from these three quarks. For example, the proton quark content is **uud**. Checking Table 44.4, we see that the values of Q/e add to 1 and that the values of the baryon number B also add to 1, as we should expect. The neutron is **udd**, with total $Q = 0$ and $B = 1$. The π^+ meson is **u $\bar{d}$** , with $Q/e = 1$ and $B = 0$, and the K^+ meson is **u $\bar{s}$** . Checking the values of S for the quark content, we see that the proton, neutron, and π^+ have strangeness 0 and that the K^+ has strangeness 1, in agreement with Table 44.3. The antiproton is $\bar{p} = \bar{u}\bar{u}\bar{d}$, the negative pion is $\pi^- = \bar{u}\bar{d}$, and so on. The quark content can also be used to explain hadron excited states and magnetic moments. **Figure 44.10** shows the quark content of two baryons and two mesons.

TABLE 44.4 Properties of the Three Original Quarks

Symbol	Q/e	Spin	Baryon Number, B	Strangeness, S	Charm, C	Bottomness, B'	Topness, T
u	$\frac{2}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	0	0	0	0
d	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	0	0	0	0
s	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	-1	0	0	0

Figure 44.10 Quark content of four hadrons. The various color combinations that are needed for color neutrality are not shown.



EXAMPLE 44.7 Determining the quark content of baryons

Given that they contain only **u**, **d**, **s**, $\bar{u}$, $\bar{d}$, and/or $\bar{s}$, find the quark content of (a) Σ^+ and (b) Λ^0 . The Σ^+ and Λ^0 (the antiparticle of the $\bar{\Lambda}^0$) are both baryons with strangeness $S = -1$.

IDENTIFY and SET UP We use the idea that the total charge of each baryon is the sum of the individual quark charges, and similarly for the baryon number and strangeness. We use the quark properties given in Table 44.4.

EXECUTE Baryons contain three quarks. If $S = -1$, exactly *one* of the three must be an **s** quark, which has $S = -1$ and $Q/e = -\frac{1}{3}$.

(a) The Σ^+ has $Q/e = +1$, so the other two quarks must both be **u** quarks (each of which has $Q/e = +\frac{2}{3}$). Hence the quark content of Σ^+ is **uus**.

(b) First we find the quark content of the Λ^0 . To yield zero total charge, the other two quarks must be **u** ($Q/e = +\frac{2}{3}$) and **d** ($Q/e = -\frac{1}{3}$), so the quark content of the Λ^0 is **uds**. The quark content of the $\bar{\Lambda}^0$ is therefore $\bar{u}\bar{d}\bar{s}$.

EVALUATE Although the Λ^0 and $\bar{\Lambda}^0$ are both electrically neutral and have the same mass, they are different particles: Λ^0 has $B = 1$ and $S = -1$, while $\bar{\Lambda}^0$ has $B = -1$ and $S = 1$.

KEYCONCEPT All baryons are composed of three quarks, and all antibaryons are composed of three antiquarks. For a baryon or antibaryon, the total electric charge equals the sum of the charges of its constituent quarks or antiquarks; the same is true of the total baryon number and total strangeness.

Motivating the Quark Model

What caused physicists to suspect that hadrons were made up of something smaller? The magnetic moment of the neutron (see Section 43.1) was one of the first reasons. In Section 27.7 we learned that a magnetic moment results from a circulating current (a motion of electric charge). But the neutron has *no* charge or, to be more accurate, no *total* charge. It could be made up of smaller particles whose charges add to zero. The quantum motion of these particles within the neutron would then give its surprising nonzero magnetic moment. To verify this hypothesis by “seeing” inside a neutron, we need a probe with a wavelength that is much smaller than the neutron’s size of about a femtometer. This probe should not be affected by the strong interaction, so that it won’t interact with the neutron as a whole but will penetrate into it and interact electromagnetically with these supposed smaller charged particles. A probe with these properties is an electron with energy greater than 10 GeV. In experiments carried out at SLAC, such electrons were scattered from neutrons and protons to help show that nucleons are indeed made up of fractionally charged, spin- $\frac{1}{2}$ pointlike particles.

The Eightfold Way

Symmetry considerations play a very prominent role in particle theory. Here are two examples. Consider the eight spin- $\frac{1}{2}$ baryons we’ve mentioned: the familiar p and n; the strange Λ^0 , Σ^+ , Σ^0 , and Σ^- ; and the doubly strange Ξ^0 and Ξ^- . For each we plot the value of strangeness S versus the value of charge Q in **Fig. 44.11**. The result is a hexagonal pattern. A similar plot for the nine spin-0 mesons (six shown in Table 44.3 plus three others not included in that table) is shown in **Fig. 44.12**; the particles fall in exactly the same hexagonal pattern! In each plot, all the particles have masses that are within about $\pm 200 \text{ MeV}/c^2$ of the median mass value of that plot, with variations due to differences in quark masses and internal potential energies.

The symmetries that lead to these and similar patterns are collectively called the **eightfold way**. They were discovered in 1961 by Murray Gell-Mann and independently by Yuval Néeman. (The name is a slightly irreverent reference to the Noble Eightfold Path, a set of principles for right living in Buddhism.) A similar pattern for the spin- $\frac{3}{2}$ baryons contains *ten* particles, arranged in a triangular pattern like pins in a bowling alley. When this pattern was first discovered, one of the particles was missing. But Gell-Mann gave

Figure 44.11 (a) Plot of S and Q values for spin- $\frac{1}{2}$ baryons, showing the symmetry pattern of the eightfold way. (b) Quark content of each spin- $\frac{1}{2}$ baryon. The quark contents of the Σ^0 and Λ^0 are the same; the Σ^0 is an excited state of the Λ^0 and can decay into it by photon emission.





Figure 44.12 (a) Plot of S and Q values for nine spin-0 mesons, showing the symmetry pattern of the eightfold way. Each particle is on the opposite side of the hexagon from its antiparticle; each of the three particles in the center is its own antiparticle. (b) Quark content of each spin-0 meson. The particles in the center are different mixtures of the three quark–antiquark pairs shown.



it a name anyway (Ω^-), predicted the properties it should have, and told experimenters what they should look for. Three years later, the particle was found during an experiment at Brookhaven National Laboratory, a spectacular success for Gell-Mann's theory. The whole series of events is reminiscent of the way in which Mendeleev used gaps in the periodic table of the elements to predict properties of undiscovered elements and to guide chemists in their search for these elements.

What binds quarks to one another? The attractive interactions among quarks are mediated by massless spin-1 bosons called **gluons** in much the same way that photons mediate the electromagnetic interaction or that pions mediated the nucleon–nucleon force in the old Yukawa theory.

Color

Quarks, having spin $\frac{1}{2}$, are fermions and so are subject to the exclusion principle. This would seem to forbid a baryon having two or three quarks with the same flavor and same spin component. To avoid this difficulty, it is assumed that each quark comes in three varieties, which are whimsically called *colors*. Red, green, and blue are the usual choices. The exclusion principle applies separately to each color. A baryon always contains one red, one green, and one blue quark, so the baryon itself has no net color. Each gluon has a color–anticolor combination (for example, blue–antired) that allows it to transmit color when exchanged, and color is conserved during emission and absorption of a gluon by a quark. The gluon-exchange process changes the colors of the quarks in such a way that there is always one quark of each color in every baryon. The color of an individual quark changes continually as gluons are exchanged.

Similar processes occur in mesons such as pions. The quark–antiquark pairs of mesons have canceling color and anticolor (for example, blue and antiblue), so mesons also have no net color. Suppose a pion initially consists of a blue quark and an antiblue antiquark. The blue quark can become a red quark by emitting a blue–antired virtual gluon. The gluon is then absorbed by the antiblue antiquark, converting it to an antired antiquark (**Fig. 44.13**). Color is conserved in each emission and absorption, but a blue–antiblue pair has become a red–antired pair. Such changes occur continually, so we have to think of a pion as a superposition of three quantum states: blue–antiblue, green–antigreen, and red–antired. On a larger scale, the strong interaction between nucleons was described in Section 44.3 as due to the exchange of virtual mesons. In terms of quarks and gluons, these mediating virtual mesons are quark–antiquark systems bound together by the exchange of gluons.

The theory of strong interactions is known as *quantum chromodynamics* (QCD). No one has been able to isolate an individual quark, and indeed QCD predicts that quarks are bound in such a way that it is impossible to obtain a free quark. An impressive body of experimental evidence supports the correctness of the quark model and the idea that quantum chromodynamics is the key to understanding the strong interactions.

Three More Quarks

Before the tau particles were discovered, there were four known leptons. This fact, together with some puzzling decay rates, led to the speculation that there might be a fourth quark flavor. This quark is labeled *c* (the *charmed* quark); it has $Q/e = \frac{2}{3}$, $B = \frac{1}{3}$, $S = 0$, and a new quantum number **charm** $C = +1$. This was confirmed in 1974 by the observation at both SLAC and the Brookhaven National Laboratory of a meson, now named ψ , with mass $3097 \text{ MeV}/c^2$. This meson was found to have several decay modes, decaying into e^+e^- , $\mu^+\mu^-$, or hadrons. The mean lifetime was found to be about 10^{-20} s . These results are consistent with ψ being a spin-1 $c\bar{c}$ system. Almost immediately after this, similar mesons of greater mass were observed and identified as excited states of the $c\bar{c}$ system. A few years later, individual mesons with a nonzero net charm quantum number, $D^0 (c\bar{u})$ and $D^+ (c\bar{d})$, and a charmed baryon, $\Lambda_c^+ (udc)$, were also observed.

In 1977 a meson with mass $9460 \text{ MeV}/c^2$, called *upsilon* (Υ), was discovered at Brookhaven. Because it had properties similar to ψ , it was conjectured that the meson was really the bound system of a new quark, *b* (the *bottom* quark), and its antiquark, $\bar{b}$.

Figure 44.13 (a) A pion containing a blue quark and an antiblue antiquark. (b) The blue quark emits a blue–antired gluon, changing to a red quark. (c) The gluon is absorbed by the antiblue antiquark, which becomes an antired antiquark. The pion now consists of a red–antired quark–antiquark pair. The actual quantum state of the pion is an equal superposition of red–antired, green–antigreen, and blue–antiblue pairs.




TABLE 44.5 Properties of the Six Quarks

Symbol	Q/e	Spin	Baryon Number, B	Strangeness, S	Charm, C	Bottomness, B'	Topness, T
u	$\frac{2}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	0	0	0	0
d	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	0	0	0	0
s	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	-1	0	0	0
c	$\frac{2}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	0	+1	0	0
b	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	0	0	-1	0
t	$\frac{2}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	0	0	0	+1

The bottom quark has the value -1 of a new quantum number B' called *bottomness*. Excited states of the Y were soon observed, as were the B^+ ($\bar{b}u$) and B^0 ($\bar{b}d$) mesons.

With the five flavors of quarks (u , d , s , c , and b) and the six flavors of leptons (e , μ , τ , ν_e , ν_μ , and ν_τ) it was an appealing conjecture that nature is symmetric in its building blocks and that therefore there should be a *sixth* quark. This quark, labeled t (top), would have $Q/e = \frac{2}{3}$, $B = \frac{1}{3}$, and a new quantum number, $T = 1$. In 1995, groups using two different detectors at Fermilab’s Tevatron announced the discovery of the top quark. The groups collided 0.9 TeV protons with 0.9 TeV antiprotons, but even with 1.8 TeV of available energy, a top–antitop ($t\bar{t}$) pair was detected in fewer than two of every 10^{11} collisions! Table 44.5 lists some properties of the six quarks. Each has a corresponding antiquark with opposite values of Q , B , S , C , B' , and T .

CAUTION Bottomness vs. baryon number Don’t confuse the bottomness quantum number B' with baryon number B . For example, the proton (which has zero bottomness and is a baryon) has $B' = 0$ and $B = +1$; the B^+ meson (which includes an antibottom quark but is not a baryon) has $B' = +1$ and $B = 0$.

TEST YOUR UNDERSTANDING OF SECTION 44.4 Is it possible to have a baryon with charge $Q = +e$ and strangeness $S = -2$?

ANSWER **no** Only the s quark, with $S = -1$, has nonzero strangeness. For a baryon to have $S = -2$, it must have two s quarks and one quark of a different flavor. Since each s quark has charge $-\frac{1}{3}e$, the nonstrange quark must have charge $+\frac{5}{3}e$ to make the net charge equal to $+e$. But *no* quark has charge $+\frac{5}{3}e$, so the proposed baryon is impossible.

44.5 THE STANDARD MODEL AND BEYOND

The particles and interactions that we’ve discussed in this chapter provide a reasonably comprehensive picture of the fundamental building blocks of nature. There is enough confidence in the basic correctness of this picture that it is called the **standard model**.

The standard model includes three families of particles: (1) the six leptons, which have no strong interactions; (2) the six quarks, from which all hadrons are made; and (3) the particles that mediate the various interactions. These mediators are gluons for the strong interaction among quarks, photons for the electromagnetic interaction, the $W^\pm$ and Z^0 particles for the weak interaction, and the graviton for the gravitational interaction.

Electroweak Unification

Theoretical physicists have long dreamed of combining all the interactions of nature into a single unified theory. As a first step, Einstein spent much of his later life trying to develop a field theory that would unify gravitation and electromagnetism. He was only partly successful.

Between 1961 and 1967, Sheldon Glashow, Abdus Salam, and Steven Weinberg developed a theory that unifies the weak and electromagnetic forces. One outcome of their **electroweak theory** is a prediction of the weak-force mediator particles, the $W^\pm$ and Z^0

bosons, including their masses. The basic idea is that the mass difference between photons (zero mass) and the weak bosons ($\approx 100 \text{ GeV}/c^2$) makes the electromagnetic and weak interactions behave quite differently at low energies. At sufficiently high energies (well above 100 GeV), however, the distinction disappears, and the two merge into a single interaction. This prediction was verified in 1983 in experiments with proton-antiproton collisions at CERN. The weak bosons were found, again with the help provided by the theoretical description, and their observed masses agreed with the predictions of the electroweak theory, a wonderful convergence of theory and experiment. The electroweak theory and quantum chromodynamics form the backbone of the standard model. Glashow, Salam, and Weinberg received the Nobel Prize in 1979.

In the electroweak theory photons are massless but the weak bosons are very massive. To account for the broken symmetry among these interaction mediators, a field called the *Higgs field* was proposed by theoretical physicists in the 1960s. We use the symbol $\phi(\vec{r}, t)$ to denote the value of this field at position $\vec{r}$ and time t . (Unlike the electric and magnetic fields, which are vectors, the Higgs field is a scalar quantity.) According to the theory, the mass of the weak bosons is proportional to the absolute value of ϕ_{av} , where ϕ_{av} is the average value of $\phi(\vec{r}, t)$ over space. **Figure 44.14** shows a simplified model of how ϕ_{av} depends on energy. At very high energies, the value of the Higgs field $\phi(\vec{r}, t)$ oscillates between positive and negative values, so its average value is $\phi_{\text{av}} = 0$ (Fig. 44.14a). But at low energies, $\phi(\vec{r}, t)$ oscillates around either a positive average value $\phi_{\text{av}} = +\phi_0$ or a negative average value $\phi_{\text{av}} = -\phi_0$ (Fig. 44.14b). The oscillation is no longer symmetric around $\phi = 0$, so the symmetry has been broken. Hence at low energies, the weak bosons acquire a nonzero mass proportional to $|\phi_{\text{av}}|$, which is equal to ϕ_0 for either of the cases shown in Fig. 44.14b.

This theory also predicts that there should be a particle called the *Higgs boson* associated with the Higgs field itself. (In an analogous way, the photon is the particle associated with the electromagnetic field.) The Higgs boson was predicted to be unstable, have zero charge and spin 0, and have a large mass. An important mission of the Large Hadron Collider at CERN was to produce the Higgs boson from the available energy in proton-proton collisions and thereby verify the existence of the Higgs field. In 2012 the first Higgs bosons were detected in such collisions, with the predicted properties. This suggests that the concept of the Higgs field—that all of space is filled with a field that gives mass to the weak bosons—is indeed correct. The Nobel Prize was awarded in 2013 to François Englert and Peter Higgs, two of the theorists who first proposed the idea of the Higgs field in 1964. Current experiments show that the mass of the Higgs boson is about $125 \text{ GeV}/c^2$, even greater than the masses of the $W^\pm$ and Z^0 weak bosons.

Grand Unified Theories


Perhaps at sufficiently high energies the strong interaction and the electroweak interaction have a convergence similar to that between the electromagnetic and weak interactions. If so, they can be unified to give a comprehensive theory of strong, weak, and electromagnetic interactions. Such schemes, called **grand unified theories** (GUTs), are still speculative.

Some grand unified theories predict the decay of the proton (in violation of conservation of baryon number), with an estimated lifetime of more than 10^{28} years. (For comparison the age of the universe is known to be 1.38×10^{10} years.) With a lifetime of 10^{28} years, six metric tons of protons would be expected to have only one decay per day, so huge amounts of material must be examined. Some of the neutrino detectors that we mentioned in Section 44.2 originally looked for, and failed to find, evidence of proton decay. Current estimates set the proton lifetime well over 10^{33} years. Some GUTs also predict the existence of magnetic monopoles, which we mentioned in Chapter 27. At present there is no confirmed experimental evidence that magnetic monopoles exist.

In the standard model, the neutrinos have zero mass. Nonzero values are controversial because experiments to determine neutrino masses are difficult both to perform

Figure 44.14 The Higgs-field value ϕ can oscillate, much like the value of the coordinate of a ball rolling within a trough with two minima. The average value of ϕ determines the masses of the $W^\pm$ and Z^0 weak bosons. (a) At high energies, the average value of ϕ is zero and the $W^\pm$ and Z^0 are massless (like the photon). (b) At low energies, the symmetry is broken. The average value of ϕ is nonzero, and the $W^\pm$ and Z^0 acquire nonzero masses.

(a) When the energy E of the system is high, the ball can oscillate between these two limits ...



(b) When the energy E of the system is low, the ball is trapped near one of the two minima ...





Figure 44.15 This photo shows the interior of the Super-Kamiokande neutrino detector in Japan. When in operation, the detector is filled with 5×10^7 kg of water. A neutrino passing through the detector can produce a faint flash of light, which is detected by the 13,000 photomultiplier tubes lining the detector walls. Data from this detector were the first to indicate that neutrinos have mass.



and to analyze. In most GUTs the neutrinos *must* have nonzero masses. If neutrinos do have mass, transitions called *neutrino oscillations* can occur, in which one type of neutrino (ν_e , ν_μ , or ν_τ) changes into another type. In 1998, scientists using the Super-Kamiokande neutrino detector in Japan (Fig. 44.15) reported the discovery of oscillations between muon neutrinos and tau neutrinos. Subsequent measurements at the Sudbury Neutrino Observatory in Canada have confirmed the existence of neutrino oscillations. This discovery is evidence for exciting physics beyond that predicted by the standard model.

The discovery of neutrino oscillations cleared up a long-standing mystery. Since the 1960s, physicists have been using sensitive detectors to look for electron neutrinos produced by nuclear fusion reactions in the sun's core (see Section 43.8). However, the observed flux of solar electron neutrinos is only one-third of the predicted value. The explanation was provided in 2002 by the Sudbury Neutrino Observatory, which can detect neutrinos of all three flavors. The results showed that the combined flux of solar neutrinos of *all* flavors is equal to the theoretical prediction for the flux of *electron* neutrinos. The explanation is that the sun is producing electron neutrinos at the predicted rate, but that two-thirds of these electron neutrinos are transformed into muon or tau neutrinos during their flight from the sun's core to a detector on earth.

Supersymmetric Theories and TOEs

The ultimate dream of theorists is to unify all four fundamental interactions, adding gravitation to the strong and electroweak interactions that are included in GUTs. Such a unified theory is whimsically called a Theory of Everything (TOE). It turns out that an essential ingredient of such theories is a spacetime continuum with more than four dimensions. The additional dimensions are “rolled up” into extremely tiny structures that we ordinarily do not notice. Depending on the scale of these structures, it may be possible for the next generation of particle accelerators to reveal the presence of extra dimensions.

Another ingredient of many theories is *supersymmetry*, which gives every boson and fermion a “superpartner” of the other spin type. For example, the proposed supersymmetric partner of the spin- $\frac{1}{2}$ electron is a spin-0 particle called the *selectron*, and that of the spin-1 photon is a spin- $\frac{1}{2}$ *photino*. As yet, no superpartner particles have been discovered, perhaps because they are too massive to be produced by the present generation of accelerators. Within a few years, new data from the Large Hadron Collider will help us decide whether these intriguing theories have merit.

TEST YOUR UNDERSTANDING OF SECTION 44.5 One aspect of the standard model is that a *d* quark can transform into a *u* quark, an electron, and an antineutrino by means of the weak interaction. If this happens to a *d* quark inside a neutron, what kind of particle remains afterward in addition to the electron and antineutrino? (i) A proton; (ii) a Σ^- ; (iii) a Σ^+ ; (iv) a Λ^0 or a Σ^0 ; (v) any of these.

ANSWER (i) If a *d* quark in a neutron (quark content uud) undergoes the process $d \rightarrow u + e^- + \bar{\nu}_e$, the remaining baryon has quark content uue and hence is a proton (see Fig. 44.11). An electron is the same as a $\bar{e}^+$ particle, so the net result is beta-minus decay: $n \rightarrow p + e^- + \bar{\nu}_e$.

44.6 THE EXPANDING UNIVERSE

In the last two sections of this chapter we'll explore briefly the connections between the early history of the universe and the interactions of fundamental particles. It is remarkable that there are such close ties between physics on the smallest scale that we've explored experimentally (the range of the weak interaction, of the order of 10^{-18} m) and physics on the largest scale (the universe itself, of the order of at least 10^{26} m).

Gravitational interactions play an essential role in the large-scale behavior of the universe. We saw in Chapter 13 how the law of gravitation explains the motions of planets in the solar system. Astronomical evidence shows that gravitational forces also dominate in larger systems such as galaxies and clusters of galaxies (**Fig. 44.16**).

Until early in the 20th century it was usually assumed that the universe was *static*; stars might move relative to each other, but there was not thought to be any overall expansion or contraction. But measurements that were begun in 1912 by Vesto Slipher at Lowell Observatory in Arizona, and continued in the 1920s by Edwin Hubble with the help of Milton Humason at Mount Wilson in California, indicated that the universe is *not* static. The motions of galaxies relative to the earth can be measured by observing the shifts in the wavelengths of their spectra. For distant galaxies these shifts are always toward longer wavelength, so they appear to be receding from us and from each other. Astronomers first assumed that these were Doppler shifts and used a relationship between the wavelength λ_0 of light measured now on earth from a source receding at speed v and the wavelength λ_S measured in the rest frame of the source when it was emitted. We can derive this relationship by inverting Eq. (37.25) for the Doppler effect, making subscript changes, and using $\lambda = c/f$; the result is

$$\lambda_0 = \lambda_S \sqrt{\frac{c + v}{c - v}} \quad (44.13)$$

Wavelengths from receding sources are always shifted toward longer wavelengths; this increase in λ is called the **redshift**. We can solve Eq. (44.13) for v :

$$v = \frac{(\lambda_0/\lambda_S)^2 - 1}{(\lambda_0/\lambda_S)^2 + 1} c \quad (44.14)$$

CAUTION Redshift, not Doppler shift Equations (44.13) and (44.14) are from the *special* theory of relativity and refer to the Doppler effect. As we'll see, the redshift from *distant* galaxies is caused by an effect that is explained by the *general* theory of relativity and is *not* a Doppler shift. However, as the ratio v/c and the fractional wavelength change $(\lambda_0 - \lambda_S)/\lambda_S$ become small, the general theory's equations approach Eqs. (44.13) and (44.14), and those equations may be used. ■

Figure 44.16 (a) The galaxy M101 is a larger version of the Milky Way galaxy of which our solar system is a part. Like all galaxies, M101 is held together by the mutual gravitational attraction of its stars, gas, dust, and other matter, all of which orbit around the galaxy's center of mass. M101 is 25 million light-years away. (b) This image shows part of the Coma cluster, an immense grouping of over 1000 galaxies that lies 300 million light-years from us. The galaxies within the cluster are all in motion. Gravitational forces between the galaxies prevent them from escaping from the cluster.

(a)



(b)



EXAMPLE 44.8 Recession speed of a galaxy

WITH ✓ VARIATION PROBLEMS

The spectral lines of various elements are detected in light from a galaxy in the constellation Ursa Major. An ultraviolet line from singly ionized calcium ($\lambda_S = 393 \text{ nm}$) is observed at wavelength $\lambda_0 = 414 \text{ nm}$, redshifted into the visible portion of the spectrum. At what speed is this galaxy receding from us?

IDENTIFY and SET UP This example uses the relationship between redshift and recession speed for a distant galaxy. We can use the wavelengths λ_S at which the light is emitted and λ_0 that we detect on earth in Eq. (44.14) to determine the galaxy's recession speed v if the fractional wavelength shift is not too great.


EXECUTE The fractional wavelength redshift for this galaxy is $\lambda_0/\lambda_S = (414 \text{ nm})/(393 \text{ nm}) = 1.053$. This is only a 5.3% increase, so we can use Eq. (44.14) with reasonable accuracy:

$$v = \frac{(1.053)^2 - 1}{(1.053)^2 + 1} c = 0.0516c = 1.55 \times 10^7 \text{ m/s}$$

EVALUATE The galaxy is receding from the earth at 5.16% of the speed of light. Rather than going through this calculation, astronomers often just state the *redshift* $z = (\lambda_0 - \lambda_S)/\lambda_S = (\lambda_0/\lambda_S) - 1$. This galaxy has redshift $z = 0.053$.

KEYCONCEPT The redshift of light from distant galaxies shows that they are moving away from us. The recession speed can be calculated from the ratio of the measured wavelength λ_0 of light from the galaxy to the wavelength λ_S measured in the rest frame of the galaxy.

Figure 44.17 Graph of recession speed versus distance for several galaxies. The best-fit straight line illustrates Hubble's law. The slope of the line is the Hubble constant, H_0 .



The Hubble Law

Analysis of redshifts from many distant galaxies led Edwin Hubble to a remarkable conclusion: The speed of recession v of a galaxy is proportional to its distance r from us (Fig. 44.17). This relationship is now called the **Hubble law**, expressed as an equation,

$$v = H_0 r \quad (44.15)$$

where H_0 is an experimental quantity commonly called the *Hubble constant*, since at any given time it is constant over space. Determining H_0 has been a key goal of the Hubble Space Telescope, which can measure distances to galaxies with unprecedented accuracy. The current best value is about $2.2 \times 10^{-18} \text{ s}^{-1}$, with an uncertainty of about 2%.

Astronomical distances are often measured in *parsecs* (pc); one parsec is the distance at which there is a one arcsecond ($1/3600^\circ$) angular separation between two objects $1.50 \times 10^{11} \text{ m}$ apart (the average distance from the earth to the sun). A distance of 1 pc is equal to 3.26 *light-years* (ly), where $1 \text{ ly} = 9.46 \times 10^{12} \text{ km}$ is the distance that light travels in one year. The Hubble constant is then commonly expressed in the mixed units (km/s)/Mpc (kilometers per second per megaparsec), where $1 \text{ Mpc} = 10^6 \text{ pc}$:

$$H_0 = (2.2 \times 10^{-18} \text{ s}^{-1}) \left(\frac{9.46 \times 10^{12} \text{ km}}{1 \text{ ly}} \right) \left(\frac{3.26 \text{ ly}}{1 \text{ pc}} \right) \left(\frac{10^6 \text{ pc}}{1 \text{ Mpc}} \right) = 68 \frac{\text{km/s}}{\text{Mpc}}$$

EXAMPLE 44.9 Determining distance with the Hubble law

WITH VARIATION PROBLEMS

Use the Hubble law to find the distance from earth to the galaxy in Ursa Major described in Example 44.8.

IDENTIFY and SET UP The Hubble law relates the redshift of a distant galaxy to its distance r from earth. We solve Eq. (44.15) for r and substitute the recession speed v from Example 44.8.

EXECUTE Using $H_0 = 68 \text{ (km/s)/Mpc} = 6.8 \times 10^4 \text{ (m/s)/Mpc}$,

$$\begin{aligned} r &= \frac{v}{H_0} = \frac{1.55 \times 10^7 \text{ m/s}}{6.8 \times 10^4 \text{ (m/s)/Mpc}} = 230 \text{ Mpc} \\ &= 2.3 \times 10^8 \text{ pc} = 7.5 \times 10^8 \text{ ly} = 7.1 \times 10^{24} \text{ m} \end{aligned}$$

EVALUATE A distance of 230 million parsecs (750 million light-years) is truly stupendous, but many galaxies lie much farther away. To appreciate the immensity of this distance, consider that our farthest-ranging unmanned spacecraft have traveled only about 0.002 ly from our planet.

KEYCONCEPT The Hubble law is the observation that the speed at which a distant galaxy moves away from us is directly proportional to its distance from us. The speed of recession can be determined from the galaxy's redshift, and the distance can then be calculated from the Hubble law.

Another aspect of Hubble's observations was that, *in all directions*, distant galaxies appeared to be receding from us. There is no particular reason to think that our galaxy is at the very center of the universe; if we lived in some other galaxy, every distant galaxy would still seem to be moving away. That is, at any given time, *the universe looks more or less the same, no matter where in the universe we are*. This important idea is called the **cosmological principle**. There are local fluctuations in density, but on average, the universe looks the same from all locations. Thus the Hubble constant is constant in space although not necessarily constant in time, and the laws of physics are the same everywhere.

The Big Bang

The Hubble law suggests that at some time in the past, all the matter in the universe was far more concentrated than it is today. It was then blown apart in a rapid expansion called the **Big Bang**, giving all observable matter more or less the velocities that we observe today. When did this happen? According to the Hubble law, matter at a distance r away from us is traveling with speed $v = H_0 r$. The time t needed to travel a distance r is

$$t = \frac{r}{v} = \frac{r}{H_0 r} = \frac{1}{H_0} = 4.5 \times 10^{17} \text{ s} = 1.4 \times 10^{10} \text{ y}$$

By this hypothesis the Big Bang occurred about 14 billion years ago. It assumes that all speeds are *constant* after the Big Bang; that is, it ignores any change in the expansion rate due to gravitational attraction or other effects. We'll return to this point later. For now, however, notice that the age of the earth determined from radioactive dating (see Section 43.4) is 4.54 billion (4.54×10^9) years. It's encouraging that our hypothesis tells us that the universe is older than the earth!

Expanding Space

The general theory of relativity takes a radically different view of the expansion just described. According to this theory, the increased wavelength is *not* caused by a Doppler shift as the universe expands into a previously empty void. Rather, the increase comes from *the expansion of space itself* and everything in intergalactic space, including the wavelengths of light traveling to us from distant sources. This is not an easy concept to grasp, and if you haven't encountered it before, it may sound like doubletalk.

An analogy may help you develop some intuition on this point. Imagine we are all bugs crawling around on a horizontal surface. We can't leave the surface, and we can see in any direction along the surface, but not up or down. We are then living in a two-dimensional world; some writers have called such a world *flatland*. If the surface is a plane, we can locate our position with two Cartesian coordinates (x, y). If the plane extends indefinitely in both the x - and y -directions, we described our space as having *infinite* extent, or as being *unbounded*. No matter how far we go, we never reach an edge or a boundary.

An alternative habitat for us bugs would be the surface of a sphere of radius R . The space would still seem infinite—we could crawl forever and never reach an edge or a boundary. Yet in this case the space is *finite* or *bounded*. To describe the location of a point in this space, we could still use two coordinates: latitude and longitude, or the spherical coordinates θ and ϕ shown in Fig. 41.5.

Now suppose the spherical surface is that of a balloon (Fig. 44.18). As we inflate the balloon more and more, increasing the radius R , the coordinates of a point don't change, yet the distance between any two points gets larger and larger. Furthermore, as R increases, the *rate of change* of distance between two points (their recession speed) is proportional to their distance apart. *The recession speed is proportional to the distance*, just as with the Hubble law. For example, the distance from Pittsburgh to Miami is twice as great as the distance from Pittsburgh to Boston. If the earth were to begin to swell, Miami would recede from Pittsburgh twice as fast as Boston would.

Although the quantity R isn't one of the two coordinates giving the position of a point on the balloon's surface, it nevertheless plays an essential role in any discussion of distance. It is the radius of curvature of our two-dimensional universe, and it is also a varying *scale factor* that changes as this universe expands.

Generalizing this picture to three dimensions isn't so easy. We have to think of our three-dimensional space as being embedded in a space with four or more dimensions, just as we visualized the two-dimensional spherical flatland as being embedded in a three-dimensional Cartesian space. Our real three-space is *not Cartesian*; to describe its characteristics in any small region requires at least one additional parameter, the curvature of space, which is analogous to the radius of the sphere. In a sense, this scale factor, which we'll continue to call R , describes the *size* of the universe, just as the radius of the sphere described the size of our two-dimensional spherical universe. We'll return later to the question of whether the universe is bounded or unbounded.


Any length that is measured in intergalactic space is proportional to R , so the wavelength of light traveling to us from a distant galaxy increases along with every other dimension as the universe expands. That is,

$$\frac{\lambda_0}{\lambda} = \frac{R_0}{R} \quad (44.16)$$


The zero subscripts refer to the values of the wavelength and scale factor *now*, just as H_0 is the current value of the Hubble constant. The quantities λ and R without subscripts are the values at *any* time—past, present, or future. For the galaxy described in Examples 44.8 and 44.9,

Figure 44.18 An inflating balloon as an analogy for an expanding universe.

(a) Points (representing galaxies) on the surface of a balloon are described by their latitude and longitude coordinates.



(b) The radius R of the balloon has increased. The coordinates of the points are the same, but the distance between them has increased.



The rate of recession for any two points is proportional to the distance between them.

we have $\lambda_0 = 414 \text{ nm}$ and $\lambda = \lambda_S = 393 \text{ nm}$, so Eq. (44.16) gives $R_0/R = 1.053$. That is, the scale factor *now* (R_0) is 5.3% larger than it was 750 million years ago when the light was emitted from that galaxy in Ursa Major. This increase of wavelength with time as the scale factor increases in our expanding universe is called the *cosmological redshift*. The farther away an object is, the longer its light takes to get to us and the greater the change in R and λ . The current largest measured wavelength ratio for galaxies is about 12.1, meaning that the volume of space itself is about $(12.1)^3 \approx 1770$ times larger than it was when the light was emitted. Do *not* attempt to substitute $\lambda_0/\lambda_S = 12.1$ into Eq. (44.14) to find the recession speed; that equation is accurate only for small cosmological redshifts and $v \ll c$. The actual value of v depends on the density of the universe, the value of H_0 , and the expansion history of the universe.

Here's a surprise: If the distance from us in the Hubble law is large enough, then the speed of recession is greater than the speed of light! This does *not* violate the special theory of relativity because the recession speed is *not* caused by the motion of the astronomical object relative to some coordinates in its region of space. Rather, $v > c$ when two sets of coordinates move apart fast enough as space itself expands. In other words, there are objects whose coordinates have been moving away from our coordinates so fast that light from them hasn't had enough time in the entire history of the universe to reach us. What we see is just the *observable* universe; we have no direct evidence about what lies beyond its horizon.

CAUTION **The universe isn't expanding into emptiness** The balloon shown in Fig. 44.18 is expanding into the empty space around it. It's a common misconception to picture the universe in the same way as a large but finite collection of galaxies that's expanding into unoccupied space. The reality is quite different! All evidence shows that our universe is *infinite*: It has no edges, so there is nothing "outside" it and it isn't "expanding into" anything. The expansion of the universe simply means that the scale factor of the universe is increasing. A good two-dimensional analogy is to think of the universe as a flat, infinitely large rubber sheet that's stretching and expanding much like the surface of the balloon in Fig. 44.18. In a sense, the infinite universe is becoming more infinite! ■

Critical Density

In an expanding universe, gravitational attractions between galaxies should slow the initial expansion. But by how much? If these attractions are strong enough, the universe should expand more and more slowly, eventually stop, and then begin to contract, perhaps all the way down to what's been called a *Big Crunch*. On the other hand, if gravitational forces are much weaker, they slow the expansion only a little, and the universe should continue to expand forever.

The situation is analogous to the problem of escape speed of a projectile launched from the earth. We studied this problem in Example 13.5 (Section 13.3). The total energy $E = K + U$ when a projectile of mass m and speed v is at a distance r from the center of the earth (mass m_E) is

$$E = \frac{1}{2}mv^2 - \frac{Gmm_E}{r}$$

If E is positive, the projectile has enough kinetic energy to move infinitely far from the earth ($r \rightarrow \infty$) and have some kinetic energy left over. If E is negative, the kinetic energy $K = \frac{1}{2}mv^2$ becomes zero and the projectile stops when $r = -Gmm_E/E$. In that case, no greater value of r is possible, and the projectile can't escape the earth's gravity.

We can carry out a similar analysis for the universe. Whether the universe continues to expand indefinitely should depend on the average *density* of matter. If matter is relatively dense, there is a lot of gravitational attraction to slow and eventually stop the expansion and make the universe contract again. If not, the expansion should continue indefinitely. We can derive an expression for the *critical density* ρ_c needed to just barely stop the expansion.

Here's a calculation based on Newtonian mechanics; it isn't relativistically correct, but it illustrates the idea. Consider a large sphere with radius R , containing many galaxies (Fig. 44.19), with total mass M . Suppose our own galaxy has mass m and is located at the surface of this sphere. According to the cosmological principle, the average distribution of matter within the sphere is uniform. The total gravitational force on our galaxy is just the force due to the mass M inside the sphere. The force on our galaxy and potential energy U due to this spherically symmetric distribution are the same as though m and M were both points, so $U = -GmM/R$, just as in Section 13.3. The net force from all the uniform distribution of mass *outside* the sphere is zero, so we'll ignore it.

The total energy E (kinetic plus potential) for our galaxy is

$$E = \frac{1}{2}mv^2 - \frac{GmM}{R} \quad (44.17)$$

If E is *positive*, our galaxy has enough energy to escape from the gravitational attraction of the mass M inside the sphere; in this case the universe should keep expanding forever. If E is *negative*, our galaxy cannot escape and the universe should eventually pull back together. The crossover between these two cases occurs when $E = 0$, so

$$\frac{1}{2}mv^2 = \frac{GmM}{R} \quad (44.18)$$

The total mass M inside the sphere is the volume $4\pi R^3/3$ times the density ρ_c :

$$M = \frac{4}{3}\pi R^3 \rho_c$$

We'll assume that the speed v of our galaxy relative to the center of the sphere is given by the Hubble law: $v = H_0 R$. Substituting these expressions for m and v into Eq. (44.18), we get

$$\begin{aligned} \frac{1}{2}m(H_0 R)^2 &= \frac{Gm}{R} \left(\frac{4}{3}\pi R^3 \rho_c \right) \quad \text{or} \\ \rho_c &= \frac{3H_0^2}{8\pi G} \quad (\text{critical density of the universe}) \end{aligned} \quad (44.19)$$

This is the *critical density*. If the average density is less than ρ_c , the universe should continue to expand indefinitely; if it is greater, the universe should eventually stop expanding and begin to contract.

Putting numbers into Eq. (44.19), we find

$$\rho_c = \frac{3(2.2 \times 10^{-18} \text{ s}^{-1})^2}{8\pi(6.67 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2)} = 8.7 \times 10^{-27} \text{ kg/m}^3$$

The mass of a hydrogen atom is $1.67 \times 10^{-27} \text{ kg}$, so this density is equivalent to about five hydrogen atoms per cubic meter.

Dark Matter, Dark Energy, and the Accelerating Universe

Astronomers have made extensive studies of the average density of matter in the universe. One way to do so is to count the number of galaxies in a patch of sky. Based on the mass of an average star and the number of stars in an average galaxy, this effort gives an estimate of the average density of *luminous* matter in the universe—that is, matter that emits electromagnetic radiation. (You are made of luminous matter because you emit infrared radiation as a consequence of your temperature; see Sections 17.7 and 39.5.) It's also necessary to take into account other luminous matter within a galaxy, including the tenuous gas and dust between the stars.

Figure 44.19 An imaginary sphere of galaxies. The net gravitational force exerted on our galaxy (at the surface of the sphere) by the other galaxies is the same as if all of their mass were concentrated at the center of the sphere. (Since the universe is infinite, there's also an infinity of galaxies outside this sphere.)




Figure 44.20 The bright spots in this image are not stars but entire galaxies. We see the most distant of these, magnified in the inset, as it was 13.3 billion years ago, when the universe was just 500 million years old. At that time the scale factor of the universe was only about 12% as large as it is now. (The red color of this galaxy is due to its very large redshift.) By comparison, we see the relatively nearby Coma cluster (see Fig. 44.16b) as it was 300 million years ago, when the scale factor was 98% of the present-day value.





Figure 44.21 The composition of our universe. Conventional matter includes all of the familiar sorts of matter that you see around you, including your body, our planet, and the sun and stars.



Another technique is to study the motions of galaxies within clusters of galaxies (see Fig. 44.16b). The motions are so slow that we can't actually see galaxies changing positions within a cluster. However, observations show that different galaxies within a cluster have somewhat different redshifts, which indicates that the galaxies are moving relative to the center of mass of the cluster. The speeds of these motions are related to the gravitational force exerted on each galaxy by the other members of the cluster, which in turn depends on the total mass of the cluster. By measuring these speeds, astronomers can determine the average density of *all* kinds of matter within the cluster, whether or not the matter emits electromagnetic radiation.

Observations using these and other techniques show that the average density of *all* matter in the universe is 31.0% of the critical density, but the average density of *luminous* matter is only 4.9% of the critical density. In other words, most of the matter in the universe is not luminous: It does not emit electromagnetic radiation of *any* kind. At present, the nature of this **dark matter** remains an outstanding mystery. Some proposed candidates for dark matter are WIMPs (weakly interacting massive particles, which are hypothetical subatomic particles far more massive than those produced in accelerator experiments) and MACHOs (massive compact halo objects, which include objects such as black holes that might form “halos” around galaxies). Whatever the true nature of dark matter, it is by far the dominant form of matter in the universe. For every kilogram of the conventional matter that has been our subject for most of this book—including electrons, protons, atoms, molecules, blocks on inclined planes, planets, and stars—there are about *five and one-third* kilograms of dark matter.

Since the average density of matter in the universe is less than the critical density, it might seem fair to conclude that the universe will continue to expand indefinitely, and that gravitational attraction between matter in different parts of the universe should slow the expansion down (albeit not enough to stop it). One way to test this prediction is to examine the redshifts of extremely distant objects. The more distant a galaxy is, the more time it takes that light to reach us from that galaxy, so the further back in time we look when we observe that galaxy. If the expansion of the universe has been slowing down, the expansion must have been more rapid in the distant past. Thus we would expect very distant galaxies to have *greater* redshifts than predicted by the Hubble law, Eq. (44.15).

Only since the 1990s has it become possible to measure accurately both the distances and the redshifts of extremely distant galaxies. The results have been totally surprising: Very distant galaxies, seen as they were when the universe was a small fraction of its present age (Fig. 44.20), have *smaller* redshifts than predicted by the Hubble law! The implication is that the expansion of the universe was slower in the past than it is now, so the expansion has been *speeding up* rather than slowing down.

If gravitational attraction should make the expansion slow down, why is it speeding up instead? Our best explanation is that space is suffused with a kind of energy that has no gravitational effect and emits no electromagnetic radiation, but rather acts as a kind of “antigravity” that produces a universal *repulsion*. This invisible, immaterial energy is called **dark energy**. As the name suggests, the nature of dark energy is poorly understood but is the subject of very active research.

Observations show that the *energy* density of dark energy (measured in, say, joules per cubic meter) is 69.0% of the critical density times c^2 ; that is, it is equal to $0.690\rho_c c^2$. As described above, the average density of matter of all kinds is 31.0% of the critical density. From the Einstein relationship $E = mc^2$, the average *energy* density of matter in the universe is therefore $0.310\rho_c c^2$. Because the energy density of dark energy is nearly three times greater than that of matter, the expansion of the universe will continue to accelerate. This expansion will never stop, and the universe will never contract.

If we account for energy of *all* kinds, the average energy density of the universe **?** is equal to $0.690\rho_c c^2 + 0.310\rho_c c^2 = 1.00\rho_c c^2$. Of this, 69.0% is the mysterious dark energy, 26.1% is the no less mysterious dark matter, and a mere 4.9% is well-understood conventional matter. How little we know about the contents of our universe (Fig. 44.21)! When we take account of the density of matter in the universe (which tends to slow the expansion of space) and the density of dark energy (which tends to speed up the expansion), the age of the universe turns out to be 13.8 billion (1.38×10^{10}) years.

What is the significance of the result that within observational error, the average energy density of the universe is equal to $\rho_c c^2$? It tells us that the universe is infinite and unbounded, but just barely so. If the average energy density were even slightly larger than $\rho_c c^2$, the universe would be finite like the surface of the balloon depicted in Fig. 44.18. As of this writing, the observational error in the average energy density is less than 1%, but we can't be totally sure that the universe *is* unbounded. Improving these measurements will be an important task for physicists and astronomers in the years ahead.

TEST YOUR UNDERSTANDING OF SECTION 44.6 Is it accurate to say that your body is made of “ordinary” matter?

ANSWER

yes... and no The material of which your body is made is ordinary to us on earth. But from a cosmic perspective your material is quite *extraordinary*: Only 4.9% of the mass and energy in the universe is in the form of atoms.

44.7 THE BEGINNING OF TIME

What an odd title for the very last section of a book! We'll describe in general terms some of the current theories about the very early history of the universe and their relationship to fundamental particle interactions. We'll find that an astonishing amount happened in the very first second.

Temperatures

The early universe was extremely dense and extremely hot, and the average particle energies were extremely large, all many orders of magnitude beyond anything that exists in the present universe. We can compare particle energy E and absolute temperature T by using the equipartition principle (see Section 18.4):

$$E = \frac{3}{2} kT \quad (44.20)$$

In this equation k is Boltzmann's constant, which we'll often express in eV/K:

$$k = 8.617 \times 10^{-5} \text{ eV/K}$$

Thus we can replace Eq. (44.20) by $E \approx (10^{-4} \text{ eV/K})T = (10^{-13} \text{ GeV/K})T$ when we're discussing orders of magnitude.

EXAMPLE 44.10 Temperature and energy

- (a) What is the average kinetic energy E (in eV) of particles at room temperature ($T = 290 \text{ K}$) and at the surface of the sun ($T = 5800 \text{ K}$)?
 (b) What approximate temperature corresponds to the ionization energy of the hydrogen atom and to the rest energies of the electron and the proton?

IDENTIFY and SET UP In this example we are to apply the equipartition principle. We use Eq. (44.20) to relate the target variables E and T .

EXECUTE (a) At room temperature, from Eq. (44.20),

$$E = \frac{3}{2} kT = \frac{3}{2} (8.617 \times 10^{-5} \text{ eV/K})(290 \text{ K}) = 0.0375 \text{ eV}$$

The temperature at the sun's surface is higher than room temperature by a factor of $(5800 \text{ K})/(290 \text{ K}) = 20$, so the average kinetic energy there is $20(0.0375 \text{ eV}) = 0.75 \text{ eV}$.

(b) The ionization energy of hydrogen is 13.6 eV. Using the approximation $E \approx (10^{-4} \text{ eV/K})T$, we have

$$T \approx \frac{E}{10^{-4} \text{ eV/K}} = \frac{13.6 \text{ eV}}{10^{-4} \text{ eV/K}} \approx 10^5 \text{ K}$$

BIO APPLICATION A Fossil Both Ancient and Recent This fossil trilobite is an example of a group of marine arthropods that flourished in earth's oceans from 540 to 250 million years ago. (By comparison, the first dinosaurs did not appear until 230 million years ago.) From our perspective, this makes trilobites almost unfathomably ancient. But compared to the time that has elapsed since the Big Bang, 13.8 billion years, even trilobites are a very recent phenomenon: They first appeared when the universe was already 96% of its present age.



Repeating this calculation for the rest energies of the electron ($E = 0.511 \text{ MeV}$) and proton ($E = 938 \text{ MeV}$) gives temperatures of 10^{10} K and 10^{13} K , respectively.

EVALUATE Temperatures in excess of 10^5 K are found in the sun's interior, so most of the hydrogen there is ionized. Temperatures of 10^{10} K or 10^{13} K are not found anywhere in the solar system; as we'll see, temperatures were this high in the very early universe.

KEYCONCEPT The average kinetic energy of particles in a gas is directly proportional to the absolute temperature T . At extremely high temperatures above 10^{10} K , this average kinetic energy can be comparable to the rest energies of subatomic particles.

Uncoupling of Interactions

We've characterized the expansion of the universe by a continual increase of the scale factor R , which we can think of very roughly as characterizing the *size* of the universe, and by a corresponding decrease in average density. As the total gravitational potential energy increased during expansion, there were corresponding *decreases* in temperature and average particle energy. As this happened, the basic interactions became progressively uncoupled.

To understand the uncouplings, recall that the unification of the electromagnetic and weak interactions occurs at energies that are large enough that the differences in mass among the various spin-1 bosons that mediate the interactions become insignificant by comparison. The electromagnetic interaction is mediated by the massless photon, and the weak interaction is mediated by the weak bosons $W^\pm$ and Z^0 with masses of the order of $100 \text{ GeV}/c^2$. At energies much *less* than 100 GeV , the two interactions seem quite different. But at energies much *greater* than 100 GeV , they become part of a single interaction, because the $W^\pm$ and Z^0 weak bosons become massless like the photon. (This occurs because the average value ϕ_{av} of the Higgs field is zero at high energy, as in Fig. 44.14a.)

The grand unified theories (GUTs) provide a similar behavior for the strong interaction. It becomes unified with the electroweak interaction at energies of the order of 10^{14} GeV , but at lower energies the two appear quite distinct. One of the reasons GUTs are still very speculative is that there is no way to do controlled experiments in this energy range, which is larger by a factor of 10^{11} than energies available with any current accelerator.

Finally, at sufficiently high energies and short distances, it is assumed that gravitation becomes unified with the other three interactions. The distance at which this happens is thought to be of the order of 10^{-35} m . This distance, called the *Planck length* l_P , is determined by the speed of light c and the fundamental constants of quantum mechanics and gravitation, $\hbar$ and G , respectively:

$$l_P = \sqrt{\frac{\hbar G}{c^3}} = 1.616 \times 10^{-35} \text{ m} \quad (44.21)$$

You should verify that this combination of constants has units of length. The *Planck time* $t_P = l_P/c$ is the time required for light to travel a distance l_P :

$$t_P = \frac{l_P}{c} = \sqrt{\frac{\hbar G}{c^5}} = 0.539 \times 10^{-43} \text{ s} \quad (44.22)$$

If we mentally go backward in time toward the Big Bang at $t = 0$, we have to stop when we reach $t = 10^{-43} \text{ s}$ because we have no adequate theory that unifies all four interactions. So as yet we have no way of knowing what happened or how the universe behaved at times earlier than the Planck time or on scales smaller than the Planck length.

The Standard Model of the History of the Universe

The description that follows is called the *standard model* of the history of the universe. The title indicates that there are substantial areas of theory that rest on solid experimental foundations and are quite generally accepted. The figure on pages 1514–1515 is a graphical description of this history, with the characteristic temperature, particle energy, and scale factor at various times. Referring to this figure frequently will help you to understand the following discussion.

In this standard model, the temperature of the universe at time $t = 10^{-43} \text{ s}$ (the Planck time) was about 10^{32} K , and the average energy per particle was approximately

$$E \approx (10^{-13} \text{ GeV/K})(10^{32} \text{ K}) = 10^{19} \text{ GeV}$$


In a totally unified theory this is about the energy below which gravity begins to behave as a separate interaction. This time therefore marked the transition from any proposed TOE to the GUT period.






During the GUT period, roughly $t = 10^{-43}$ to 10^{-35} s, the strong and electroweak forces were still unified, and the universe consisted of a soup of quarks and leptons transforming into each other so freely that there was no distinction between the two families of particles. Other, much more massive particles may also have been freely created and destroyed. One important characteristic of GUTs is that at sufficiently high energies, baryon number is not conserved. (We mentioned earlier the proposed decay of the proton, which has not yet been observed.) Thus by the end of the GUT period the numbers of quarks and antiquarks may have been unequal. This point has important implications; we'll return to it at the end of the section.

By $t = 10^{-35}$ s the temperature had decreased to about 10^{27} K and the average energy to about 10^{14} GeV. At this energy the strong force separated from the electroweak force (**Fig. 44.22**), and baryon number and lepton numbers began to be separately conserved. This separation of the strong force was analogous to a phase change such as boiling a liquid, with an associated heat of vaporization. Think of it as being similar to boiling a heavy nucleus, pulling the particles apart beyond the short range of the nuclear force. As a result, the universe underwent a dramatic expansion (far more rapid than the present-day expansion rate) called *cosmic inflation*. In one model, the scale factor R increased by a factor of 10^{50} in 10^{-32} s.

At $t = 10^{-32}$ s the universe was a mixture of quarks, leptons, and the mediating bosons (gluons, photons, and the weak bosons $W^\pm$ and Z^0). It continued to expand and cool from the inflationary period to $t = 10^{-6}$ s, when the temperature was about 10^{13} K and typical energies were about 1 GeV (comparable to the rest energy of a nucleon; see Example 44.11). The quarks then began to bind together, forming nucleons and antinucleons. There were still enough photons of sufficient energy to produce nucleon–antinucleon pairs to balance the process of nucleon–antinucleon annihilation. However, by about $t = 10^{-2}$ s, most photon energies fell well below the threshold energy for such pair production. There was a slight excess of nucleons over antinucleons; as a result, virtually all of the antinucleons and most of the nucleons annihilated one another. A similar equilibrium occurred later between the production of electron–positron pairs from photons and the annihilation of such pairs. At about $t = 14$ s the average energy dropped to around 1 MeV, below the threshold for e^+e^- pair production. After pair production ceased, virtually all of the remaining positrons were annihilated, leaving the universe with many more protons and electrons than the antiparticles of each.

Figure 44.22 Schematic diagram showing the times and energies at which the various interactions are thought to have uncoupled. The energy scale is backward because the average energy decreased as the age of the universe increased.





Logarithmic scales show characteristic temperature, particle energy, and scale factor of the universe as functions of time.

A Brief History of the Universe

AGE OF IONS

Expanding, cooling
gas of ionized
H and He.

AGE OF ATOMS

Neutral atoms form;
universe becomes
transparent to most light.




AGE OF STARS AND GALAXIES

Thermonuclear fusion
begins in stars, forming
heavier nuclei.


 10^{13} s

 10^{15} s

NOW



Up until about $t = 1$ s, neutrons and neutrinos could be produced in the endoergic reaction



After this time, most electrons no longer had enough energy for this reaction. The average neutrino energy also decreased, and as the universe expanded, equilibrium reactions that involved *absorption* of neutrinos (which occurred with decreasing probability) became inoperative. At this time, in effect, the flux of neutrinos and antineutrinos throughout the universe uncoupled from the rest of the universe. Because of the extraordinarily low probability for neutrino absorption, most of this flux is still present today, although cooled greatly by expansion. The standard model of the universe predicts a present neutrino temperature of about 2 K, but no experiment has yet been able to test this prediction.

Nucleosynthesis

At about $t = 1$ s, the ratio of protons to neutrons was determined by the Boltzmann distribution factor $e^{-\Delta E/kT}$, where ΔE is the difference between the neutron and proton rest energies: $\Delta E = 1.294$ MeV. At a temperature of about 10^{10} K, this distribution factor gives about 4.5 times as many protons as neutrons. However, as we have discussed, free neutrons (with a half-life of 887 s) decay spontaneously to protons. This decay caused the proton-to-neutron ratio to increase until about $t = 225$ s. At this time, the temperature was about 10^9 K, and the average energy was well below 2 MeV.

This energy average was critical because the binding energy of the *deuteron* (a neutron and a proton combined to form a ${}^2\text{H}$ nucleus) is 2.22 MeV (see Section 43.2). A neutron bound in a deuteron does not decay spontaneously. As the average energy decreased, a proton and a neutron could combine to form a deuteron, and there were fewer and fewer photons with 2.22 MeV or more of energy to dissociate the deuterons again. Therefore the combining of protons and neutrons into deuterons halted the decay of free neutrons.

The formation of deuterons starting at about $t = 225$ s marked the beginning of the period of formation of nuclei, or *nucleosynthesis*. At this time, there were about seven protons for each neutron. The deuteron (${}^2\text{H}$) can absorb a neutron and form a triton (${}^3\text{H}$), or it can absorb a proton and form ${}^3\text{He}$. Then ${}^3\text{H}$ can absorb a proton and ${}^3\text{He}$ can absorb a neutron, each yielding ${}^4\text{He}$ (the alpha particle). A few ${}^7\text{Li}$ nuclei may also have formed by fusion of ${}^3\text{H}$ and ${}^4\text{He}$ nuclei. According to the theory, essentially all the ${}^1\text{H}$ and ${}^4\text{He}$ in the present universe formed at this time. But then the building of nuclei almost ground to a halt. The reason is that *no* nuclide with mass number $A = 5$ has a half-life greater than 10^{-21} s. Alpha particles simply do not permanently absorb neutrons or protons. The nuclide ${}^8\text{Be}$ that is formed by fusion of two ${}^4\text{He}$ nuclei is unstable, with an extremely short half-life, about 7×10^{-17} s. At this time, the average energy was still much too large for electrons to be bound to nuclei; there were not yet any atoms.

CONCEPTUAL EXAMPLE 44.11 The relative abundance of hydrogen and helium in the universe

Nearly all of the protons and neutrons in the seven-to-one ratio at $t = 225$ s either formed ${}^4\text{He}$ or remained as ${}^1\text{H}$. After this time, what was the resulting relative abundance of ${}^1\text{H}$ and ${}^4\text{He}$, by mass?

SOLUTION The ${}^4\text{He}$ nucleus contains two protons and two neutrons. For every two neutrons present at $t = 225$ s there were 14 protons. The two neutrons and two of the 14 protons make up one ${}^4\text{He}$ nucleus, leaving 12 protons (${}^1\text{H}$ nuclei). So there were eventually 12 ${}^1\text{H}$ nuclei for every ${}^4\text{He}$ nucleus. The masses of ${}^1\text{H}$ and ${}^4\text{He}$ are about 1 u and 4 u, respectively, so there were 12 u of ${}^1\text{H}$ for every 4 u of ${}^4\text{He}$. Therefore

the relative abundance, by mass, was 75% ${}^1\text{H}$ and 25% ${}^4\text{He}$. This result agrees very well with estimates of the present H–He ratio in the universe, an important confirmation of this part of the theory.

KEYCONCEPT The first atomic nuclei formed about three and a half minutes after the Big Bang. The present-day ratio of hydrogen nuclei to helium nuclei in the universe was determined by the ratio of protons to neutrons at that moment in the history of the universe.

Further nucleosynthesis did not occur until very much later, well after $t = 10^{13}$ s (about 380,000 y). At that time, the temperature was about 3000 K, and the average energy was a few tenths of an electron volt. Because the ionization energies of hydrogen and helium

atoms are 13.6 eV and 24.5 eV, respectively, almost all the hydrogen and helium was electrically neutral (not ionized). With the electrical repulsions of the nuclei canceled out, gravitational attraction could slowly pull the neutral atoms together to form clouds of gas and eventually stars. Thermonuclear reactions in stars then produced all of the more massive nuclei. In Section 43.8 we discussed one cycle of thermonuclear reactions in which ^1H becomes ^4He .

For stars whose mass is 40% of the sun's mass or greater, as the hydrogen is consumed the star's core begins to contract as the inward gravitational pressure exceeds the outward gas and radiation pressure. The gravitational potential energy decreases as the core contracts, so the kinetic energy of nuclei in the core increases. Eventually the core temperature becomes high enough to begin another process, *helium fusion*. First two ^4He nuclei fuse to form ^8Be , which is highly unstable. But because a star's core is so dense and collisions among nuclei are so frequent, there is a nonzero probability that a third ^4He nucleus will fuse with the ^8Be nucleus before it can decay. The result is the stable nuclide ^{12}C . This is called the *triple-alpha process*, since three ^4He nuclei (that is, alpha particles) fuse to form one carbon nucleus. Then successive fusions with ^4He give ^{16}O , ^{20}Ne , and ^{24}Mg . All these reactions are exoergic. They release energy to heat up the star, and ^{12}C and ^{16}O can fuse to form elements with higher and higher atomic number.

For nuclides that can be created in this manner, the binding energy per nucleon peaks at mass number $A = 56$ with the nuclide ^{56}Fe , so exoergic fusion reactions stop with Fe. But successive neutron captures followed by beta decays can continue the synthesis of more massive nuclei. If the star is massive enough, it may eventually explode as a *supernova*, sending out into space the heavy elements that were produced by the earlier processes (**Fig. 44.23**; see also Fig. 37.7). In space, the debris and other interstellar matter can gravitationally bunch together to form a new generation of stars and planets. Our sun is one such "second-generation" star. The sun's planets and everything on them (including you) contain matter that was long ago blasted into space by an exploding supernova.

Background Radiation

In 1965 Arno Penzias and Robert Wilson, working at Bell Telephone Laboratories in New Jersey on satellite communications, turned a microwave antenna skyward and found a background signal that had no apparent preferred direction. (This signal produces about 1% of the "hash" you see on a TV that's disconnected from cable and tuned to an unused channel.) Further research has shown that the radiation that is received has a frequency spectrum that fits Planck's blackbody radiation law, Eq. (39.24) (Section 39.5). The wavelength of peak intensity is 1.063 mm (in the microwave region of the spectrum), with a corresponding absolute temperature $T = 2.725\text{ K}$. Penzias and Wilson contacted physicists at Princeton University who had begun the design of an antenna to search for radiation that was a remnant from the early evolution of the universe. We mentioned above that neutral atoms began to form at about $t = 380,000$ years when the temperature was 3000 K. With far fewer charged particles present than previously, the universe became transparent at this time to electromagnetic radiation of long wavelength. The 3000 K blackbody radiation therefore survived, cooling to its present 2.725 K temperature as the universe expanded. The *cosmic background radiation* is among the most clear-cut experimental confirmations of the Big Bang theory. **Figure 44.24** shows a modern map of the cosmic background radiation.




Figure 44.23 The Veil Nebula in the constellation Cygnus is a remnant of a supernova explosion that occurred about 8000 years ago. The gas ejected from the supernova is still moving very rapidly. Collisions between this fast-moving gas and the tenuous material of interstellar space excite the gas and cause it to glow. The portion of the nebula shown here is about 40 ly (12 pc) in length.




Figure 44.24 This false-color map shows microwave radiation from the entire sky mapped onto an oval. When this radiation was emitted 380,000 years after the Big Bang, the regions shown in blue were slightly cooler and denser than average. Within these cool, dense regions formed galaxies, including the Milky Way galaxy of which our solar system, our earth, and our selves are part.

EXAMPLE 44.12 Expansion of the universe

By approximately what factor has the universe expanded since $t = 380,000$ y?

IDENTIFY and SET UP We use the idea that as the universe has expanded, all intergalactic wavelengths have expanded with it. The Wien displacement law, Eq. (39.21), relates the peak wavelength λ_m in blackbody radiation to the temperature T . Given the temperatures of the cosmic background radiation today (2.725 K) and at $t = 380,000$ y (3000 K) we can determine the factor by which wavelengths have changed and hence determine the factor by which the universe has expanded.

EXECUTE We rewrite Eq. (39.21) as

$$\lambda_m = \frac{2.90 \times 10^{-3} \text{ m} \cdot \text{K}}{T}$$

Hence the peak wavelength λ_m is inversely proportional to T . As the universe expands, all intergalactic wavelengths (including λ_m) increase

in proportion to the scale factor R . The temperature has decreased by the factor $(3000 \text{ K})/(2.725 \text{ K}) \approx 1100$, so λ_m and the scale factor must both have *increased* by this factor. Thus, between $t = 380,000$ y and the present, the universe has expanded by a factor of about 1100.

EVALUATE Our results show that since $t = 380,000$ y, any particular intergalactic *volume* has increased by a factor of about $(1100)^3 = 1.3 \times 10^9$. They also show that when the cosmic background radiation was emitted, its peak wavelength was $\frac{1}{1100}$ of the present-day value of 1.063 mm, or 967 nm. This is in the infrared region of the spectrum.

KEYCONCEPT The present-day temperature of the cosmic background radiation tells us how much the universe has expanded since the time 380,000 years after the Big Bang when the first atoms formed and the universe became transparent.

Matter and Antimatter

One of the most remarkable features of our universe is the asymmetry between matter and antimatter. You might think that the universe should have equal numbers of protons and antiprotons and of electrons and positrons, but this doesn't appear to be the case. Theories of the early universe must explain this imbalance.

We've mentioned that most GUTs include violation of conservation of baryon number at energies at which the strong and electroweak interactions have converged. If particle–antiparticle symmetry is also violated, we have a mechanism for making more quarks than antiquarks, more leptons than antileptons, and eventually more matter than antimatter. One serious problem is that any asymmetry that is created in this way during the GUT era might be wiped out by the electroweak interaction after the end of the GUT era. If so, there must be some mechanism that creates particle–antiparticle asymmetry at a much *later* time. The problem of the matter–antimatter asymmetry is still very much an open one.

There are still many unanswered questions at the intersection of particle physics and cosmology. Is the energy density of the universe precisely equal to $\rho_c c^2$, or are there small but important differences? What is dark energy? Has the density of dark energy remained constant over the history of the universe, or has the density changed? What is dark matter? What happened during the first 10^{-43} s after the Big Bang? Can we see evidence that the strong and electroweak interactions undergo a grand unification at high energies? The search for the answers to these and many other questions about our physical world continues to be one of the most exciting adventures of the human mind.

TEST YOUR UNDERSTANDING OF SECTION 44.7 Given a sufficiently powerful telescope, could we detect photons emitted earlier than $t = 380,000$ y?


ANSWER

| no Prior to $t = 380,000$ y the temperature was so high that atoms could not form, so free electrons and protons were plentiful. These charged particles are very effective at scattering photons, so light could not propagate very far and the universe was opaque. The oldest photons that we can detect date from the time $t = 380,000$ y when atoms formed and the universe became transparent.


CHAPTER 44 SUMMARY

Fundamental particles: Each particle has an antiparticle; some particles are their own antiparticles. Particles can be created and destroyed, some of them (including electrons and positrons) only in pairs or in conjunction with other particles and antiparticles.

Particles serve as mediators for the fundamental interactions. The photon is the mediator of the electromagnetic interaction. Yukawa proposed the existence of mesons to mediate the nuclear interaction. Mediating particles that can exist only because of the uncertainty principle for energy are called virtual particles.




Particle accelerators and detectors: Cyclotrons, synchrotrons, and linear accelerators are used to accelerate charged particles to high energies for experiments with particle interactions. Only part of the beam energy is available to cause reactions with targets at rest. This problem is avoided in colliding-beam experiments. (See Examples 44.1–44.3.)




Particles and interactions: Four fundamental interactions are found in nature: the strong, electromagnetic, weak, and gravitational interactions. Particles can be described in terms of their interactions and of quantities that are conserved in all or some of the interactions.

Fermions have half-integer spins; bosons have integer spins. Leptons, which are fermions, have no strong interactions. Strongly interacting particles are called hadrons. They include mesons, which are always bosons, and baryons, which are always fermions. There are conservation laws for three different lepton numbers and for baryon number. Additional quantum numbers, including strangeness and charm, are conserved in some interactions. (See Examples 44.4–44.6.)


Quarks: Hadrons are composed of quarks. There are thought to be six types of quarks. The interaction between quarks is mediated by gluons. Quarks and gluons have an additional attribute called color. (See Example 44.7.)




Symmetry and the unification of interactions: Symmetry considerations play a central role in all fundamental-particle theories. The electromagnetic and weak interactions become unified at high energies into the electroweak interaction. In grand unified theories the strong interaction is also unified with these interactions, but at much higher energies.



The expanding universe and its composition: The Hubble law shows that galaxies are receding from each other and that the universe is expanding. Observations show that the rate of expansion is accelerating due to the presence of dark energy, which makes up 69.0% of the energy in the universe. Only 4.9% of the energy in the universe is in the form of conventional matter; the remaining 26.1% is dark matter, whose nature is poorly understood. (See Examples 44.8 and 44.9.)



The history of the universe: In the standard model of the universe, a Big Bang gave rise to the first fundamental particles. They eventually formed into the lightest atoms as the universe expanded and cooled. The cosmic background radiation is a relic of the time when these atoms formed. The heavier elements were manufactured much later by fusion reactions inside stars. (See Examples 44.10–44.12.)





GUIDED PRACTICE

For assigned homework and other learning materials, go to Mastering Physics.

KEY EXAMPLE VARIATION PROBLEMS

Be sure to review **EXAMPLE 44.1** (Section 44.2) before attempting these problems.

VP44.1.1 In a cyclotron the maximum radius for the path of protons is 0.800 m. The magnetic field is 0.600 T. Find (a) the frequency of the alternating voltage applied to the dees, (b) the maximum proton kinetic energy in MeV, and (c) the maximum proton speed in terms of c .

VP44.1.2 A cyclotron at the TRIUMF laboratory in Vancouver, Canada, has a magnetic field of 0.460 T. It accelerates negative hydrogen ions with charge $-e$ and mass 1.67×10^{-27} kg. Find (a) the radius of the path of the ions when they have speed 6.00×10^6 m/s and (b) the angular frequency of the ions' motion.

VP44.1.3 The first cyclotron (in 1930) had a diameter of 11.4 cm and was used to accelerate negative hydrogen ions (charge $-e$, mass 1.67×10^{-27} kg) to kinetic energies of 80.0 keV. Find (a) the magnetic field used and (b) the frequency of the alternating voltage applied to the dees.

VP44.1.4 A cyclotron is used to accelerate alpha particles (charge $+2e$, mass 6.64×10^{-27} kg). The frequency of the alternating voltage applied to the dees is 5.80 MHz. Find (a) the magnetic field used and (b) the kinetic energy, in MeV, of the alpha particles when the radius of their circular path is 0.650 m.

Be sure to review **EXAMPLES 44.2 and 44.3** (Section 44.2) before attempting these problems.

VP44.3.1 A proton with kinetic energy K collides with a proton at rest. Both protons remain after the collision along with an eta-prime (η') particle that has rest energy 958 MeV. In order for this reaction to happen, what must be (a) the minimum available energy in the center-of-momentum system and (b) the minimum value of K ?

VP44.3.2 A proton with kinetic energy K collides with a proton at rest. The particles that remain after the collision are one proton and a delta-plus (Δ^+) that has rest energy 1232 MeV. In order for this reaction to

happen, what must be (a) the minimum available energy in the center-of-momentum system and (b) the minimum value of K ?

VP44.3.3 A negative pion (π^-) that has rest energy 140 MeV and kinetic energy K collides with a proton at rest. What remains after the collision is a delta-zero (Δ^0) that has rest energy 1232 MeV. In order for this reaction to happen, what must be (a) the minimum available energy in the center-of-momentum system and (b) the minimum value of K ?

VP44.3.4 The Large Hadron Collider gives each proton a total energy of 7 TeV $= 7.00 \times 10^3$ GeV. (a) If a proton that has this energy collides with a second proton that is stationary, what is the available energy? (b) After the collision, both protons remain as well as a neutral particle X . What is the maximum mass of X ?

Be sure to review **EXAMPLES 44.8 and 44.9** (Section 44.6) before attempting these problems.

VP44.9.1 The red spectral line of hydrogen has a wavelength of 656.3 nm when the hydrogen is at rest. This line is observed to have a wavelength of 725 nm in the spectrum of a galaxy. Find (a) the redshift z and (b) the recession speed of the galaxy in terms of c .

VP44.9.2 A prominent absorption line in the spectrum of the sun (see Fig. 39.9) and many galaxies is due to singly ionized calcium. In the rest frame, this line has an ultraviolet wavelength of 396.9 nm. If a galaxy is receding from us at $0.0711c$, find (a) the wavelength at which we'll observe this line and (b) the redshift z .

VP44.9.3 Find the distances to (a) a galaxy with a recession speed of $0.0992c$ and (b) a galaxy with a recession speed of $0.0711c$. Give your answers in pc.

VP44.9.4 The spectrum of a Type Ia supernova (an exploding star so luminous that it can be seen even in distant galaxies) has an emission line due to singly ionized silicon. If astronomers observe this emission line at 615.0 nm in the laboratory but 666 nm in the spectrum of a Type Ia supernova in a distant galaxy, find (a) the recession speed of the galaxy in m/s and (b) the distance to the galaxy in pc.

BRIDGING PROBLEM Hyperons, Pions, and the Expanding Universe

(a) A Λ^0 hyperon decays into a neutron and a neutral pion (π^0). Find the kinetic energies of the decay products and the fraction of the kinetic energy that is carried off by each particle. (b) A π^0 is at rest in the galaxy shown in Fig. 44.20. If a physicist on earth detects one of the two photons emitted in the decay of this π^0 , what is the energy of this detected photon?

SOLUTION GUIDE

IDENTIFY and SET UP

- Which quantities are conserved in the Λ^0 decay? In the π^0 decay?
- The universe expanded during the time that the photon traveled from the cluster to earth. How does this affect the wavelength and energy of the photon that the physicist detects?
- List the unknown quantities for each part of the problem and identify the target variables.
- Select the equations that will allow you to solve for the target variables.

EXECUTE

- Write the conservation equations for the decay of the Λ^0 . [Hint: It's useful to write the energy E of a particle in terms of its momentum p and mass m with $E = (p^2c^2 + m^2c^4)^{1/2}$.]
- Solve the conservation equations for the energy of one of the decay products. (Hint: Rearrange the energy conservation equation so that one of the $(p^2c^2 + m^2c^4)^{1/2}$ terms is on one side of the equation. Then square both sides.) Then use $K = E - mc^2$.
- Find the fraction of the total kinetic energy that goes into the neutron and into the pion.
- Write the conservation equations for the decay of the π^0 at rest and find the energy of each emitted photon. By what factor does the wavelength of this photon change as it travels from the galaxy to earth? By what factor does the photon energy change? (Hint: See Fig. 44.20.)

EVALUATE

- Which of the Λ^0 decay products should have the greater kinetic energy? Should the detected π^0 decay photon have more or less energy than when it was emitted?

PROBLEMS

•, ••, •••: Difficulty levels. **CP**: Cumulative problems incorporating material from earlier chapters. **CALC**: Problems requiring calculus. **DATA**: Problems involving real data, scientific evidence, experimental design, and/or statistical reasoning. **BIO**: Biosciences problems.

DISCUSSION QUESTIONS

Q44.1 Is it possible that some parts of the universe contain antimatter whose atoms have nuclei made of antiprotons and antineutrons, surrounded by positrons? How could we detect this condition without actually going there? Can we detect these antiatoms by identifying the light they emit as composed of antiphotons? Explain. What problems might arise if we actually *did* go there?

Q44.2 Given the Heisenberg uncertainty principle, is it possible to create particle–antiparticle pairs that exist for extremely short periods of time before annihilating? Does this mean that empty space is really empty?

Q44.3 When they were first discovered during the 1930s and 1940s, there was confusion as to the identities of pions and muons. What are the similarities and most significant differences?

Q44.4 The gravitational force between two electrons is weaker than the electric force by the order of 10^{-40} . Yet the gravitational interactions of matter were observed and analyzed long before electrical interactions were understood. Why?

Q44.5 When a π^0 decays to two photons, what happens to the quarks of which it was made?

Q44.6 Why can't an electron decay to two photons? To two neutrinos?

Q44.7 According to the standard model of the fundamental particles, what are the similarities between baryons and leptons? What are the most important differences?

Q44.8 According to the standard model of the fundamental particles, what are the similarities between quarks and leptons? What are the most important differences?

Q44.9 The quark content of the neutron is *udd*. (a) What is the quark content of the antineutron? Explain your reasoning. (b) Is the neutron its own antiparticle? Why or why not? (c) The quark content of the ψ is $c\bar{c}$. Is the ψ its own antiparticle? Explain your reasoning.

Q44.10 Does the universe have a center? Explain.

Q44.11 Does it make sense to ask, "If the universe is expanding, what is it expanding into?"

Q44.12 Assume that the universe has an edge. Placing yourself at that edge in a thought experiment, explain why this assumption violates the cosmological principle.

Q44.13 Explain why the cosmological principle requires that H_0 must have the same value everywhere in space, but does not require that it be constant in time.

EXERCISES

Section 44.1 Fundamental Particles—A History

44.1 • The starship *Enterprise*, of television and movie fame, is powered by combining matter and antimatter. If the entire 400-kg antimatter fuel supply of the *Enterprise* combines with matter, how much energy is released? How does this compare to the U.S. yearly energy use, which is roughly 1.0×10^{20} J?

44.2 •• **CP** Two equal-energy photons collide head-on and annihilate each other, producing a $\mu^+\mu^-$ pair. The muon mass is given in terms of the electron mass in Section 44.1. (a) Calculate the maximum wavelength of the photons for this to occur. If the photons have this wavelength, describe the motion of the μ^+ and μ^- immediately after they are produced. (b) If the wavelength of each photon is half the value calculated in part (a), what is the speed of each muon after they have moved apart? Use correct relativistic expressions for momentum and energy.

44.3 •• A positive pion at rest decays into a positive muon and a neutrino. (a) Approximately how much energy is released in the decay? (Assume the neutrino has zero rest mass. Use the muon and pion masses given in terms of the electron mass in Section 44.1.) (b) Why can't a positive muon decay into a positive pion?

44.4 • A proton and an antiproton annihilate, producing two photons. Find the energy, frequency, and wavelength of each photon (a) if the p and $\bar{p}$ are initially at rest and (b) if the p and $\bar{p}$ collide head-on, each with an initial kinetic energy of 620 MeV.

44.5 •• **CP** For the nuclear reaction given in Eq. (44.2) assume that the initial kinetic energy and momentum of the reacting particles are negligible. Calculate the speed of the α particle immediately after it leaves the reaction region.

44.6 •• Estimate the range of the force mediated by an ω^0 meson that has mass $783 \text{ MeV}/c^2$.

Section 44.2 Particle Accelerators and Detectors

44.7 • In a collision experiment, a proton at rest is struck by an antiproton. (a) What is the minimum kinetic energy of the antiproton if the available energy is 2.00 TeV? (b) If a colliding beam is used instead of a stationary target, what minimum kinetic energy for each beam is required for an available energy of 2.00 TeV?

44.8 • An electron with a total energy of 30.0 GeV collides with a stationary positron. (a) What is the available energy? (b) If the electron and positron are accelerated in a collider, what total energy corresponds to the same available energy as in part (a)?

44.9 • Deuterons in a cyclotron travel in a circle with radius 32.0 cm just before emerging from the dees. The frequency of the applied alternating voltage is 9.00 MHz. Find (a) the magnetic field and (b) the kinetic energy and speed of the deuterons upon emergence.

44.10 • Bubble-chamber photographs taken with a 0.40 T magnetic field show at one point a positron moving perpendicular to the field in a 15 cm circle. What is the magnitude of the positron's momentum at that point?

44.11 • (a) A high-energy beam of alpha particles collides with a stationary helium gas target. What must the total energy of a beam particle be if the available energy in the collision is 16.0 GeV? (b) If the alpha particles instead interact in a colliding-beam experiment, what must the energy of each beam be to produce the same available energy?

44.12 • The first cyclotron used a 1.3 T magnetic field and had an 11 cm radius. (a) Find the maximum kinetic energy of protons accelerated by this cyclotron. Is it accurate to use nonrelativistic expressions in your calculation? (b) When the cyclotron was accelerating protons, at what frequency was it operated?

44.13 •• (a) What is the speed of a proton that has total energy 1000 GeV? (b) What is the angular frequency ω of a proton with the speed calculated in part (a) in a magnetic field of 4.00 T? Use both the nonrelativistic Eq. (44.7) and the correct relativistic expression, and compare the results.

44.14 •• Calculate the minimum beam energy in a proton-proton collider to initiate the $p + p \rightarrow p + p + \eta^0$ reaction. The rest energy of the η^0 is 547.9 MeV (see Table 44.3).

44.15 •• You work for a start-up company that is planning to use antiproton annihilation to produce radioactive isotopes for medical applications. One way to produce antiprotons is by the reaction $p + p \rightarrow p + p + p + \bar{p}$ in proton-proton collisions. (a) You first consider a colliding-beam experiment in which the two proton beams have equal kinetic energies. To produce an antiproton via this reaction, what is the required minimum kinetic energy of the protons in each beam? (b) You then consider the collision of a proton beam with a stationary proton target. For this experiment, what is the required minimum kinetic energy of the protons in the beam?

Section 44.3 Particles and Interactions

44.16 • In about 8% of Ω^- decays, the decay products are a Ξ^- and a π^0 . (a) What is the energy released in this decay? (b) In the decay, what is the change in baryon number, and what is the change in strangeness? Your results should show that this decay is allowed for the weak nuclear interaction but not for the strong interaction.

44.17 • A K^+ meson at rest decays into two π mesons. (a) What are the allowed combinations of π^0 , π^+ , and π^- as decay products? (b) Find the total kinetic energy of the π mesons.

44.18 • How much energy is released when a μ^- muon at rest decays into an electron and two neutrinos? Neglect the small masses of the neutrinos.

44.19 • What is the mass (in kg) of the Z^0 ? What is the ratio of the mass of the Z^0 to the mass of the proton?

44.20 • The discovery of the Ω^- particle helped confirm Gell-Mann's eightfold way. If an Ω^- decays into a Λ^0 and a K^- , what is the total kinetic energy of the decay products?

44.21 • If a Σ^+ at rest decays into a proton and a π^0 , what is the total kinetic energy of the decay products?

44.22 • Which of the following reactions obey the conservation of baryon number? (a) $p + p \rightarrow p + e^+$; (b) $p + n \rightarrow 2e^+ + e^-$; (c) $p \rightarrow n + e^- + \bar{\nu}_e$; (d) $p + \bar{p} \rightarrow 2\gamma$.

44.23 • In which of the following decays are the three lepton numbers conserved? In each case, explain your reasoning. (a) $\mu^- \rightarrow e^- + \nu_e + \bar{\nu}_\mu$; (b) $\tau^- \rightarrow e^- + \bar{\nu}_e + \nu_\tau$; (c) $\pi^+ \rightarrow e^+ + \gamma$; (d) $n \rightarrow p + e^- + \bar{\nu}_e$.

44.24 • In which of the following reactions or decays is strangeness conserved? In each case, explain your reasoning. (a) $K^+ \rightarrow \mu^+ + \nu_\mu$; (b) $n + K^+ \rightarrow p + \pi^0$; (c) $K^+ + K^- \rightarrow \pi^0 + \pi^0$; (d) $p + K^- \rightarrow \Lambda^0 + \pi^0$.

Section 44.4 Quarks and Gluons

44.25 • Determine the electric charge, baryon number, strangeness quantum number, and charm quantum number for the following quark combinations: (a) uds ; (b) $c\bar{u}$; (c) ddd ; and (d) $d\bar{c}$. Explain your reasoning.

44.26 •• Determine the electric charge, baryon number, strangeness quantum number, and charm quantum number for the following quark combinations: (a) uus , (b) $c\bar{s}$, (c) $d\bar{d}\bar{u}$, and (d) $c\bar{b}$.

44.27 • Given that each particle contains only combinations of $u, d, s, \bar{u}, \bar{d}$, and $\bar{s}$, use the method of Example 44.7 to deduce the quark content of (a) a particle with charge $+e$, baryon number 0, and strangeness $+1$; (b) a particle with charge $+e$, baryon number -1 , and strangeness $+1$; (c) a particle with charge 0, baryon number $+1$, and strangeness -2 .

44.28 • What is the total kinetic energy of the decay products when an up quark at rest decays to $\tau^+ + \tau^-$?

Section 44.5 The Standard Model and Beyond

44.29 • Section 44.5 states that current experiments show that the mass of the Higgs boson is about $125 \text{ GeV}/c^2$. What is the ratio of the mass of the Higgs boson to the mass of a proton?

Section 44.6 The Expanding Universe

44.30 • The spectrum of the sodium atom is detected in the light from a distant galaxy. (a) If the 590.0 nm line is redshifted to 658.5 nm, at what speed is the galaxy receding from the earth? (b) Use the Hubble law to calculate the distance of the galaxy from the earth.

44.31 • A galaxy in the constellation Pisces is 5210 Mly from the earth. (a) Use the Hubble law to calculate the speed at which this galaxy is receding from earth. (b) What redshifted ratio λ_0/λ_S is expected for light from this galaxy?

44.32 • Redshift. The definition of the redshift z is given in Example 44.8. (a) Show that Eq. (44.13) can be written as $1 + z = ([1 + \beta]/[1 - \beta])^{1/2}$, where $\beta = v/c$. (b) The observed redshift for a certain galaxy is $z = 0.700$. Find the speed of the galaxy relative to the earth; assume the redshift is described by Eq. (44.14). (c) Use the Hubble law to find the distance of this galaxy from the earth.

Section 44.7 The Beginning of Time

44.33 • Calculate the reaction energy Q (in MeV) for the reaction $e^- + p \rightarrow n + \nu_e$. Is this reaction endoergic or exoergic?

44.34 • Calculate the energy (in MeV) released in the triple-alpha process $3 {}^4\text{He} \rightarrow {}^{12}\text{C}$.

44.35 • Calculate the reaction energy Q (in MeV) for the nucleosynthesis reaction



Is this reaction endoergic or exoergic?

PROBLEMS

44.36 •• The strong nuclear force can be crudely modeled as a Hooke's-law spring force, increasing linearly with the quark separation distance. The energy stored in this "spring" corresponds to the energy content of the gluon field. In this picture, as quarks are further separated, increasing energy is stored between them. At a critical separation distance, the energy converts to matter, and a new quark-antiquark pair is generated, guaranteeing that there can never be a free quark. (a) A proton has a diameter of about 1.5 fm. Estimate the repulsive Coulomb force between two up quarks separated by 0.5 fm. (b) Model the strong force as $F_s = ks$, where s is the distance between two quarks. If this force balances the electrostatic repulsion between two up quarks when $s = 0.5 \text{ fm}$, what is the effective spring constant k , in SI units? (c) Convert k into units of MeV/fm^2 . (d) How much energy is stored in the gluon field when $s = 0.5 \text{ fm}$? The mass of an up quark is thought to be about $2.3 \text{ MeV}/c^2$. (e) How much energy is needed to produce an up quark and an antiup quark? (f) How far would two up quarks need to be separated so that the gluon energy $\frac{1}{2}ks^2$ matches the rest energy of an up-antiup quark pair?

44.37 •• CP BIO Radiation Therapy with π^- Mesons. Beams of π^- mesons are used in radiation therapy for certain cancers. The energy comes from the complete decay of the π^- to stable particles. (a) Write out the complete decay of a π^- meson to stable particles. What are these particles? (b) How much energy is released from the complete decay of a single π^- meson to stable particles? (You can ignore the very small masses of the neutrinos.) (c) How many π^- mesons need to decay to give a dose of 50.0 Gy to 10.0 g of tissue? (d) What would be the equivalent dose in part (c) in Sv and in rem? Consult Table 43.3 and use the largest appropriate RBE for the particles involved in this decay.

44.38 •• A proton and an antiproton collide head-on with equal kinetic energies. Two γ rays with wavelengths of 0.720 fm are produced. Calculate the kinetic energy of the incident proton.

44.39 •• The 7 TeV proton bunches that circulate in opposite directions around the 27 km ring of the Large Hadron Collider smash together at beam crossings every 25 ns. (a) How many bunches meet every second? (b) Each bunch has 115 billion protons. Of these, typically 20 collide during each crossing. Estimate the fraction of protons that collide per crossing. (c) Estimate how many collisions take place each second. (d) The bunches are 30 cm long and are squeezed to a diameter of 20 μm . Estimate the density of protons in a bunch, in units of protons/ mm^3 . (e) Estimate the density of hadrons in ordinary matter. (*Hint:* Divide your mass, which is mostly due to hadrons, by the mass of a proton to get the number of hadrons in your body. Then divide by the estimated volume.)

44.40 •• Calculate the threshold kinetic energy for the reaction $\pi^- + p \rightarrow \Sigma^0 + K^0$ if a π^- beam is incident on a stationary proton target. The K^0 has a mass of 497.7 MeV/ c^2 .

44.41 • Each of the following reactions is missing a single particle. Calculate the baryon number, charge, strangeness, and the three lepton numbers (where appropriate) of the missing particle, and from this identify the particle. (a) $p + p \rightarrow p + \Lambda^0 + ?$; (b) $K^- + n \rightarrow \Lambda^0 + ?$; (c) $p + \bar{p} \rightarrow n + ?$; (d) $\bar{\nu}_\mu + p \rightarrow n + ?$

44.42 •• An η^0 meson at rest decays into three π mesons. (a) What are the allowed combinations of π^0 , π^+ , and π^- as decay products? (b) Find the total kinetic energy of the π mesons.

44.43 • The ϕ meson has mass 1019.4 MeV/ c^2 and a measured energy width of 4.4 MeV/ c^2 . Using the uncertainty principle, estimate the lifetime of the ϕ meson.

44.44 • Estimate the energy width (energy uncertainty) of the ψ if its mean lifetime is 7.6×10^{-21} s. What fraction is this of its rest energy?

44.45 •• CP BIO One proposed proton decay is $p^+ \rightarrow e^+ + \pi^0$, which violates both baryon and lepton number conservation, so the proton lifetime is expected to be very long. Suppose the proton half-life were 1.0×10^{18} y. (a) Calculate the energy deposited per kilogram of body tissue (in rad) due to the decay of the protons in your body in one year. Model your body as consisting entirely of water. Only the two protons in the hydrogen atoms in each H_2O molecule would decay in the manner shown; do you see why? Assume that the π^0 decays to two γ rays, that the positron annihilates with an electron, and that all the energy produced in the primary decay and these secondary decays remains in your body. (b) Calculate the equivalent dose (in rem) assuming an RBE of 1.0 for all the radiation products, and compare with the 0.1 rem due to the natural background and the 5.0 rem guideline for industrial workers. Based on your calculation, can the proton lifetime be as short as 1.0×10^{18} y?

44.46 •• A ϕ meson (see Problem 44.43) at rest decays via $\phi \rightarrow K^+ + K^-$. It has strangeness 0. (a) Find the kinetic energy of the K^+ meson. (Assume that the two decay products share kinetic energy equally, since their masses are equal.) (b) Suggest a reason the decay $\phi \rightarrow K^+ + K^- + \pi^0$ has not been observed. (c) Suggest reasons the decays $\phi \rightarrow K^+ + \pi^-$ and $\phi \rightarrow K^+ + \mu^-$ have not been observed.

44.47 ••• CP About 10,000 cosmic-ray protons, each with hundreds of MeV of energy, impinge on each square meter of our upper atmosphere each second. They collide with atmospheric nitrogen and oxygen to produce secondary showers of newly created particles, including many muons. Muons have a mass of 105.7 MeV/ c^2 and an average lifetime of 2.197 μs . Consider a secondary cosmic muon produced at an altitude of 15.00 km aimed directly downward with an energy of 6.000 GeV. With such high energy, the muon can travel a great distance into the earth without slowing down. (a) Determine the speed of this muon. (b) How far would this muon travel in one lifetime if there were no relativistic effects? (c) In the frame of the muon, what distance separates its creation position from the earth's surface? (d) If one lifetime passes in the frame of the muon, how much time passes in the frame of the earth? (e) How far does a muon travel in this time as seen from the earth? (f) Does the muon survive its trip to the surface? How far will it penetrate the earth in its lifetime?

44.48 ••• CP A Ξ^- particle at rest decays to a Λ^0 and a π^- . (a) Find the total kinetic energy of the decay products. (b) What fraction of the energy is carried off by each particle? (Use relativistic expressions for momentum and energy.)

44.49 ••• CP A Σ^- particle moving in the $+x$ -direction with kinetic energy 180 MeV decays into a π^- and a neutron. The π^- moves in the $+y$ -direction. What is the kinetic energy of the neutron, and what is the direction of its velocity? Use relativistic expressions for energy and momentum.

44.50 ••• CP The K^0 meson has rest energy 497.7 MeV. A K^0 meson moving in the $+x$ -direction with kinetic energy 225 MeV decays into a π^+ and a π^- , which move off at equal angles above and below the $+x$ -axis. Calculate the kinetic energy of the π^+ and the angle it makes with the $+x$ -axis. Use relativistic expressions for energy and momentum.

44.51 •• DATA While tuning up a medical cyclotron for use in isotope production, you obtain the data given in the table.

B (T)	0.10	0.20	0.30	0.40
K_{max} (MeV)	0.068	0.270	0.608	1.080

B is the uniform magnetic field in the cyclotron, and K_{max} is the maximum kinetic energy of the particle being accelerated, which is a proton. The radius R of the proton path at maximum kinetic energy has the same value for each magnetic-field value. (a) Compare the kinetic energy values in the table to the rest energy mc^2 of a proton. Is it necessary to use relativistic expressions in your analysis? Explain. (b) Graph your data as K_{max} versus B^2 . Use the slope of the best-fit straight line to your data to find R . (c) What is the maximum kinetic energy for a 0.25 T magnetic field? (d) What is the angular frequency ω of the proton when $B = 0.40$ T?

44.52 •• DATA The decay products from the decay of short-lived unstable particles can provide evidence that these particles have been produced in a collision experiment. As an initial step in designing an experiment to detect short-lived hadrons, you make a literature study of their decays. Table 44.3 gives experimental data for the mass and typical decay modes of the particles Σ^- , Ξ^0 , Δ^{++} , and Ω^- . (a) Which of these four particles has the largest mass? The smallest? (b) By the decay modes shown in the table, for which of these particles do the decay products have the greatest total kinetic energy? The least?


44.53 •• DATA You have entered a graduate program in particle physics and are learning about the use of symmetry. You begin by repeating the analysis that led to the prediction of the Ω^- particle. Nine of the spin- $\frac{3}{2}$ baryons are four Δ particles, each with mass 1232 MeV/ c^2 , strangeness 0, and charges $+2e$, $+e$, 0 , and $-e$; three Σ^* particles, each with mass 1385 MeV/ c^2 , strangeness -1 , and charges $+e$, 0 , and $-e$; and two Ξ^* particles, each with mass 1530 MeV/ c^2 , strangeness -2 , and charges 0 and $-e$. (a) Place these particles on a plot of S versus Q . Deduce the Q and S values of the tenth spin- $\frac{3}{2}$ baryon, the Ω^- particle, and place it on your diagram. Also label the particles with their masses. The mass of the Ω^- is 1672 MeV/ c^2 ; is this value consistent with your diagram? (b) Deduce the three-quark combinations (of u , d , and s) that make up each of these ten particles. Redraw the plot of S versus Q from part (a) with each particle labeled by its quark content. What regularities do you see?

44.54 ••• CP A positive kaon K^+ which is a bound state of an up quark and an antidown quark, can decay into an antimuon and a muon neutrino via a weak interaction process mediated by a W^+ boson. We can determine how much of the kaon's energy each of the daughter particles obtains using relativistic kinematics. (a) In the rest frame of the kaon, the speed of the antimuon is v and its Lorentz factor is $\gamma = (1 - v^2/c^2)^{-1/2}$. Using the notation E_ν for the energy of the neutrino, M_K for the mass of the kaon, and M_μ for the mass of the antimuon, write the relativistic expression for conservation of energy. Treat the neutrino as massless. (b) Write the relativistic equation for conservation of momentum. (c) Solve these two equations to determine v as a function of the mass ratio $\sigma = M_K/M_\mu$. (d) Using the masses M_K and M_μ found in Tables 44.3 and 44.2, respectively, determine the value of σ . (e) What is the energy of $E_\mu = \gamma M_\mu c^2$ of the antimuon? (f) What is the energy of the neutrino? (g) Do these energies add up to the rest energy of the kaon?

CHALLENGE PROBLEMS

44.55 ••• CALC Consider the following hypothetical universe: The Nibiruvians and the Xibalbans are minuscule “flat” societies located at an angular separation of $\theta = 60^\circ$ on the surface of a two-dimensional spherical universe with radius R , as shown in Fig. P44.55. (a) What is the distance D between the two societies in terms of R and θ ? (b) If R is increasing in time, what is the speed $V = dD/dt$ at which the civilizations are separating, as a function of R , dR/dt , and D ? (c) The separation velocity and the distance D are related by $V = BD$. Determine the “Bubble parameter” B in terms of $R(t)$. Note that B is not constant. (Similarly, the Hubble “constant” H_0 is presumed to be changing with time.) (d) The radius of this universe was $R_0 = 500.0$ m upon its creation at time $t = 0$, and it has been increasing at a constant rate of $1.00 \mu\text{m/s}$. What was the Bubble parameter exactly four years after creation? (e) What is the distance between Nibiru and Xibalba four years after creation? (f) At what speed are they separating at that time? (g) At that time, the Nibiruvians transmit isotropic ripple waves with their “light speed” of $c = 6.35 \mu\text{m/s}$ and wavelength 1.00 nm. How long does it take these waves to reach Xibalba? (*Hint:* In time dt the waves travel an angular distance $d\theta = c dt/R(t)$. Integrate to obtain an expression for the total angular distance traveled as a function of

Figure P44.55



time. Solve for the time needed to travel a given angular distance.) (h) At what wavelength will the Xibalbans observe these waves? (*Hint:* Use Eq. (44.16).)

44.56 ••• CP Consider a collision in which a stationary particle with mass M is bombarded by a particle with mass m , speed v_0 , and total energy (including rest energy) E_m . (a) Use the Lorentz transformation to write the velocities v_m and v_M of particles m and M in terms of the speed v_{cm} of the center of momentum. (b) Use the fact that the total momentum in the center-of-momentum frame is zero to obtain an expression for v_{cm} in terms of m , M , and v_0 . (c) Combine the results of parts (a) and (b) to obtain Eq. (44.9) for the total energy in the center-of-momentum frame.

MCAT-STYLE PASSAGE PROBLEMS

BIO Looking Under the Hood of PET. In the imaging method called positron emission tomography (PET), a patient is injected with molecules containing atoms that have nuclei with an excess of protons. As they decay into neutrons, these protons emit positrons. An emitted positron travels a short distance and slows to near-zero velocity; when it encounters an electron, they may annihilate each other and emit two photons in opposite directions. The patient is enclosed in a circular array of detectors, with the tissue to be imaged centered in the array. If two photons of the proper energy strike two detectors simultaneously (within 10 ns), we can conclude that the photons were produced by positron-electron annihilation somewhere along a line connecting the detectors. By observing many such simultaneous events, we can create a map of the distribution of positron-emitting atoms in the tissue. However, photons can be absorbed or scattered as they pass through tissue. The number of photons remaining after they travel a distance x through tissue is given by $N = N_0 e^{-\mu x}$, where N_0 is the initial number of photons and μ is the attenuation coefficient, which is approximately 0.1 cm^{-1} for photons of this energy. The index of refraction of biological tissue for x rays is 1.

44.57 What is the energy of each photon produced by positron-electron annihilation? (a) $\frac{1}{2}m_e v^2$, where v is the speed of the emitted positron; (b) $m_e v^2$; (c) $\frac{1}{2}m_e c^2$; (d) $m_e c^2$.

44.58 Suppose that positron-electron annihilations occur on the line 3 cm from the center of the line connecting two detectors. Will the resultant photons be counted as having arrived at these detectors simultaneously? (a) No, because the time difference between their arrivals is 100 ms; (b) no, because the time difference is 200 ms; (c) yes, because the time difference is 0.1 ns; (d) yes, because the time difference is 0.2 ns.

44.59 If the annihilation photons come from a part of the body that is separated from the detector by 20 cm of tissue, what percentage of the photons that originally traveled toward the detector remains after they have passed through the tissue? (a) 1.4%; (b) 8.6%; (c) 14%; (d) 86%.

ANSWERS

Chapter Opening Question ?

(v) Only 4.9% of the mass and energy of the universe is in the form of “normal” matter. Of the rest, 26.1% is poorly understood dark matter and 69.0% is even more mysterious dark energy.

Key Example VARIATION Problems

VP44.1.1 (a) 9.15 MHz (b) 11.0 MeV (c) $0.153c$

VP44.1.2 (a) 0.136 m (b) $4.41 \times 10^7 \text{ rad/s}$

VP44.1.3 (a) 0.717 T (b) 10.9 MHz

VP44.1.4 (a) 0.756 T (b) 11.6 MeV

VP44.3.1 (a) 2834 MeV (b) $2.41 \times 10^3 \text{ MeV} = 2.41 \text{ GeV}$

VP44.3.2 (a) 2170 MeV (b) 634 MeV

VP44.3.3 (a) 1232 MeV (b) 190 MeV

VP44.3.4 (a) 115 GeV (b) $113 \text{ GeV}/c^2$

VP44.9.1 (a) 0.105 (b) 0.0992c

VP44.9.2 (a) 426 nm (b) 0.0738

VP44.9.3 (a) $4.4 \times 10^8 \text{ pc}$ (b) $3.1 \times 10^8 \text{ pc}$

VP44.9.4 (a) $2.38 \times 10^7 \text{ m/s}$ (b) $3.5 \times 10^8 \text{ pc}$

Bridging Problem

(a) neutron: 5.78 MeV (0.140 of total);

pion: 35.62 MeV (0.860 of total)

(b) 8.1 MeV

APPENDIX A

THE INTERNATIONAL SYSTEM OF UNITS

The *Système International d'Unités*, abbreviated **SI**, is the system developed by the General Conference on Weights and Measures and adopted by nearly all the industrial nations of the world. The following material is adapted from the National Institute of Standards and Technology (<http://physics.nist.gov/cuu>).

Quantity	Name of unit	Symbol	
SI base units			
length	meter	m	
mass	kilogram	kg	
time	second	s	
electric current	ampere	A	
thermodynamic temperature	kelvin	K	
amount of substance	mole	mol	
luminous intensity	candela	cd	
SI derived units			Equivalent units
area	square meter	m ²	
volume	cubic meter	m ³	
frequency	hertz	Hz	s ⁻¹
mass density (density)	kilogram per cubic meter	kg/m ³	
speed, velocity	meter per second	m/s	
angular velocity	radian per second	rad/s	
acceleration	meter per second squared	m/s ²	
angular acceleration	radian per second squared	rad/s ²	
force	newton	N	kg • m/s ²
pressure (mechanical stress)	pascal	Pa	N/m ²
kinematic viscosity	square meter per second	m ² /s	
dynamic viscosity	newton-second per square meter	N • s/m ²	
work, energy, quantity of heat	joule	J	N • m
power	watt	W	J/s
quantity of electricity	coulomb	C	A • s
potential difference, electromotive force	volt	V	J/C, W/A
electric field strength	volt per meter	V/m	N/C
electrical resistance	ohm	Ω	V/A
capacitance	farad	F	A • s/V
magnetic flux	weber	Wb	V • s
inductance	henry	H	V • s/A
magnetic flux density	tesla	T	Wb/m ²
magnetic field strength	ampere per meter	A/m	
magnetomotive force	ampere	A	
luminous flux	lumen	lm	cd • sr
luminance	candela per square meter	cd/m ²	
illuminance	lux	lx	lm/m ²
wave number	1 per meter	m ⁻¹	
entropy	joule per kelvin	J/K	
specific heat capacity	joule per kilogram-kelvin	J/kg • K	
thermal conductivity	watt per meter-kelvin	W/m • K	

Quantity	Name of unit	Symbol	Equivalent units
radiant intensity	watt per steradian	W/sr	
activity (of a radioactive source)	becquerel	Bq	s ⁻¹
radiation dose	gray	Gy	J/kg
radiation dose equivalent	sievert	Sv	J/kg
SI supplementary units			
plane angle	radian	rad	
solid angle	steradian	sr	

Definitions of SI Units

meter (m) The *meter* is the length equal to the distance traveled by light, in vacuum, in a time of 1/299,792,458 second.

kilogram (kg) The *kilogram* is the unit of mass; it is defined by taking the value of Planck's constant h to be exactly $6.62607015 \times 10^{-34} \text{ kg} \cdot \text{m}^2/\text{s}$.

second (s) The *second* is the duration of 9,192,631,770 periods of the radiation corresponding to the transition between the two hyperfine levels of the ground state of the cesium-133 atom.

ampere (A) The *ampere* is a current of one coulomb per second, where the coulomb is defined in terms of the elementary charge e .

kelvin (K) The *kelvin*, unit of thermodynamic temperature, is defined by taking the value of the Boltzmann constant k to be exactly $1.380649 \times 10^{-23} \text{ J/K}$.

ohm (Ω) The *ohm* is the electric resistance between two points of a conductor when a constant difference of potential of 1 volt, applied between these two points, produces in this conductor a current of 1 ampere, this conductor not being the source of any electromotive force.

coulomb (C) The *coulomb* is the quantity of electricity such that the elementary charge e is exactly $1.602176634 \times 10^{-19} \text{ C}$ transported in 1 second by a current of 1 ampere.

candela (cd) The *candela* is the luminous intensity, in a given direction, of a source that emits monochromatic radiation of frequency 540×10^{12} hertz and that has a radiant intensity in that direction of 1/683 watt per steradian.

mole (mol) The *mole* is the SI unit of substance. One mole contains exactly $6.02214076 \times 10^{23}$ elementary entities.

newton (N) The *newton* is that force that gives to a mass of 1 kilogram an acceleration of 1 meter per second per second.

joule (J) The *joule* is the work done when the point of application of a constant force of 1 newton is displaced a distance of 1 meter in the direction of the force.

watt (W) The *watt* is the power that gives rise to the production of energy at the rate of 1 joule per second.

volt (V) The *volt* is the difference of electric potential between two points of a conducting wire carrying a constant current of 1 ampere, when the power dissipated between these points is equal to 1 watt.

weber (Wb) The *weber* is the magnetic flux that, linking a circuit of one turn, produces in it an electromotive force of 1 volt as it is reduced to zero at a uniform rate in 1 second.

lumen (lm) The *lumen* is the luminous flux emitted in a solid angle of 1 steradian by a uniform point source having an intensity of 1 candela.

farad (F) The *farad* is the capacitance of a capacitor between the plates of which there appears a difference of potential of 1 volt when it is charged by a quantity of electricity equal to 1 coulomb.

henry (H) The *henry* is the inductance of a closed circuit in which an electromotive force of 1 volt is produced when the electric current in the circuit varies uniformly at a rate of 1 ampere per second.

radian (rad) The *radian* is the plane angle between two radii of a circle that cut off on the circumference an arc equal in length to the radius.

steradian (sr) The *steradian* is the solid angle that, having its vertex in the center of a sphere, cuts off an area of the surface of the sphere equal to that of a square with sides of length equal to the radius of the sphere.

SI Prefixes To form the names of multiples and submultiples of SI units, apply the prefixes listed in Appendix F.

APPENDIX B

USEFUL MATHEMATICAL RELATIONS

Algebra

$$a^{-x} = \frac{1}{a^x} \quad a^{(x+y)} = a^x a^y \quad a^{(x-y)} = \frac{a^x}{a^y}$$

Logarithms: If $\log a = x$, then $a = 10^x$. $\log a + \log b = \log(ab)$ $\log a - \log b = \log(a/b)$ $\log(a^n) = n \log a$

If $\ln a = x$, then $a = e^x$. $\ln a + \ln b = \ln(ab)$ $\ln a - \ln b = \ln(a/b)$ $\ln(a^n) = n \ln a$

Quadratic formula: If $ax^2 + bx + c = 0$, $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

Binomial Theorem


$$(a + b)^n = a^n + na^{n-1}b + \frac{n(n-1)a^{n-2}b^2}{2!} + \frac{n(n-1)(n-2)a^{n-3}b^3}{3!} + \dots$$

Trigonometry

In the right triangle ABC , $x^2 + y^2 = r^2$.

Definitions of the trigonometric functions:

$$\sin \alpha = y/r \quad \cos \alpha = x/r \quad \tan \alpha = y/x$$



Identities: $\sin^2 \alpha + \cos^2 \alpha = 1$

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha$$

$$\sin \frac{1}{2} \alpha = \sqrt{\frac{1 - \cos \alpha}{2}}$$

$$\sin(-\alpha) = -\sin \alpha$$

$$\cos(-\alpha) = \cos \alpha$$

$$\sin(\alpha \pm \pi/2) = \pm \cos \alpha$$

$$\cos(\alpha \pm \pi/2) = \mp \sin \alpha$$

$$\tan \alpha = \frac{\sin \alpha}{\cos \alpha}$$

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha = 2 \cos^2 \alpha - 1 = 1 - 2 \sin^2 \alpha$$

$$\cos \frac{1}{2} \alpha = \sqrt{\frac{1 + \cos \alpha}{2}}$$

$$\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$$

$$\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \pm \sin \alpha \sin \beta$$


$$\sin \alpha + \sin \beta = 2 \sin \frac{1}{2}(\alpha + \beta) \cos \frac{1}{2}(\alpha - \beta)$$

$$\cos \alpha + \cos \beta = 2 \cos \frac{1}{2}(\alpha + \beta) \cos \frac{1}{2}(\alpha - \beta)$$

For any triangle $A'B'C'$ (not necessarily a right triangle) with sides a , b , and c and angles α , β , and γ :

Law of sines: $\frac{\sin \alpha}{a} = \frac{\sin \beta}{b} = \frac{\sin \gamma}{c}$

Law of cosines: $c^2 = a^2 + b^2 - 2ab \cos \gamma$



Geometry

Circumference of circle of radius r :

Area of circle of radius r :

Volume of sphere of radius r :

$$C = 2\pi r$$

$$A = \pi r^2$$

$$V = 4\pi r^3/3$$

Surface area of sphere of radius r :

Volume of cylinder of radius r and height h :

$$A = 4\pi r^2$$

$$V = \pi r^2 h$$

Calculus

Derivatives:

$$\frac{d}{dx}x^n = nx^{n-1}$$

$$\frac{d}{dx}\ln ax = \frac{1}{x}$$

$$\frac{d}{dx}e^{ax} = ae^{ax}$$

$$\frac{d}{dx}\sin ax = a \cos ax$$

$$\frac{d}{dx}\cos ax = -a \sin ax$$

Integrals:

$$\int x^n dx = \frac{x^{n+1}}{n+1} \quad (n \neq -1)$$

$$\int \frac{dx}{x} = \ln x$$

$$\int e^{ax} dx = \frac{1}{a}e^{ax}$$

$$\int \sin ax dx = -\frac{1}{a}\cos ax$$

$$\int \cos ax dx = \frac{1}{a}\sin ax$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin \frac{x}{a}$$

$$\int \frac{dx}{\sqrt{x^2 + a^2}} = \ln(x + \sqrt{x^2 + a^2})$$

$$\int \frac{dx}{x^2 + a^2} = \frac{1}{a}\arctan \frac{x}{a}$$

$$\int \frac{dx}{(x^2 + a^2)^{3/2}} = \frac{1}{a^2} \frac{x}{\sqrt{x^2 + a^2}}$$

$$\int \frac{xdx}{(x^2 + a^2)^{3/2}} = -\frac{1}{\sqrt{x^2 + a^2}}$$

Power series (convergent for range of x shown):

$$(1+x)^n = 1 + nx + \frac{n(n-1)x^2}{2!} + \frac{n(n-1)(n-2)}{3!}x^3$$

$$+ \cdots (|x| < 1)$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots (\text{all } x)$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots (\text{all } x)$$

$$\tan x = x + \frac{x^3}{3} + \frac{2x^5}{15} + \frac{17x^7}{315} + \cdots (|x| < \pi/2)$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots (\text{all } x)$$

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots (|x| < 1)$$

APPENDIX C

THE GREEK ALPHABET

Name	Capital	Lowercase	Name	Capital	Lowercase	Name	Capital	Lowercase
Alpha	A	α	Iota	I	ι	Rho	P	ρ
Beta	B	β	Kappa	K	κ	Sigma	Σ	σ
Gamma	Γ	γ	Lambda	Λ	λ	Tau	T	τ
Delta	Δ	δ	Mu	M	μ	Upsilon	Υ	υ
Epsilon	E	ϵ	Nu	N	ν	Phi	Φ	ϕ
Zeta	Z	ζ	Xi	Ξ	ξ	Chi	X	χ
Eta	H	η	Omicron	O	o	Psi	Ψ	ψ
Theta	Θ	θ	Pi	Π	π	Omega	Ω	ω

APPENDIX D

PERIODIC TABLE OF THE ELEMENTS

Group	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Period																		
1	1 H 1.008																	2 He 4.003
2	3 Li 6.941	4 Be 9.012											5 B 10.811	6 C 12.011	7 N 14.007	8 O 15.999	9 F 18.998	10 Ne 20.180
3	11 Na 22.990	12 Mg 24.305											13 Al 26.982	14 Si 28.086	15 P 30.974	16 S 32.065	17 Cl 35.453	18 Ar 39.948
4	19 K 39.098	20 Ca 40.078	21 Sc 44.956	22 Ti 47.867	23 V 50.942	24 Cr 51.996	25 Mn 54.938	26 Fe 55.845	27 Co 58.933	28 Ni 58.693	29 Cu 63.546	30 Zn 65.409	31 Ga 69.723	32 Ge 72.64	33 As 74.922	34 Se 78.96	35 Br 79.904	36 Kr 83.798
5	37 Rb 85.468	38 Sr 87.62	39 Y 88.906	40 Zr 91.224	41 Nb 92.906	42 Mo 95.94	43 Tc (98)	44 Ru 101.07	45 Rh 102.906	46 Pd 106.42	47 Ag 107.868	48 Cd 112.411	49 In 114.818	50 Sn 118.710	51 Sb 121.760	52 Te 127.60	53 I 126.904	54 Xe 131.293
6	55 Cs 132.905	56 Ba 137.327	71 Lu 174.967	72 Hf 178.49	73 Ta 180.948	74 W 183.84	75 Re 186.207	76 Os 190.23	77 Ir 192.217	78 Pt 195.078	79 Au 196.967	80 Hg 200.59	81 Tl 204.383	82 Pb 207.2	83 Bi 208.980	84 Po (209)	85 At (210)	86 Rn (222)
7	87 Fr (223)	88 Ra (226)	103 Lr (262)	104 Rf (267)	105 Db (270)	106 Sg (269)	107 Bh (270)	108 Hs (269)	109 Mt (278)	110 Ds (281)	111 Rg (281)	112 Cn (285)	113 Nh (286)	114 Fl (289)	115 Mc (289)	116 Lv (293)	117 Ts (293)	118 Og (294)
Lanthanoids			57 La 138.905	58 Ce 140.116	59 Pr 140.908	60 Nd 144.24	61 Pm (145)	62 Sm 150.36	63 Eu 151.964	64 Gd 157.25	65 Tb 158.925	66 Dy 162.500	67 Ho 164.930	68 Er 167.259	69 Tm 168.934	70 Yb 173.04		
Actinoids			89 Ac (227)	90 Th (232)	91 Pa (231)	92 U (238)	93 Np (237)	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)		

For each element, the average atomic mass of the mixture of isotopes occurring in nature is shown. For elements having no stable isotope, the approximate atomic mass of the longest-lived isotope is shown in parentheses. All atomic masses are expressed in atomic mass units ($1 \text{ u} = 1.660539040(20) \times 10^{-27} \text{ kg}$), equivalent to grams per mole (g/mol).

APPENDIX E

UNIT CONVERSION FACTORS

Length

1 m = 100 cm = 1000 mm = $10^6 \mu\text{m}$ = 10^9 nm
1 km = 1000 m = 0.6214 mi
1 m = 3.281 ft = 39.37 in.
1 cm = 0.3937 in.
1 in. = 2.540 cm
1 ft = 30.48 cm
1 yd = 91.44 cm
1 mi = 5280 ft = 1.609 km
1 Å = 10^{-10} m = 10^{-8} cm = 10^{-1} nm
1 nautical mile = 6080 ft
1 light-year = $9.461 \times 10^{15} \text{ m}$

Area

1 cm² = 0.155 in.²
1 m² = 10⁴ cm² = 10.76 ft²
1 in.² = 6.452 cm²
1 ft² = 144 in.² = 0.0929 m²

Volume

1 liter = 1000 cm³ = 10^{-3} m^3 = 0.03531 ft³ = 61.02 in.³
1 ft³ = 0.02832 m³ = 28.32 liters = 7.477 gallons
1 gallon = 3.788 liters

Time

1 min = 60 s
1 h = 3600 s
1 d = 86,400 s
1 y = 365.24 d = $3.156 \times 10^7 \text{ s}$

Angle

1 rad = 57.30° = $180^\circ/\pi$
1° = 0.01745 rad = $\pi/180 \text{ rad}$
1 revolution = 360° = $2\pi \text{ rad}$
1 rev/min (rpm) = 0.1047 rad/s

Speed

1 m/s = 3.281 ft/s
1 ft/s = 0.3048 m/s
1 mi/min = 60 mi/h = 88 ft/s
1 km/h = 0.2778 m/s = 0.6214 mi/h
1 mi/h = 1.466 ft/s = 0.4470 m/s = 1.609 km/h
1 furlong/fortnight = $1.662 \times 10^{-4} \text{ m/s}$

Acceleration

1 m/s² = 100 cm/s² = 3.281 ft/s²
1 cm/s² = 0.01 m/s² = 0.03281 ft/s²
1 ft/s² = 0.3048 m/s² = 30.48 cm/s²
1 mi/h · s = 1.467 ft/s²

Mass

1 kg = 10³ g = 0.0685 slug
1 g = $6.85 \times 10^{-5} \text{ slug}$
1 slug = 14.59 kg
1 u = $1.661 \times 10^{-27} \text{ kg}$
1 kg has a weight of 2.205 lb when $g = 9.80 \text{ m/s}^2$

Force

1 N = 10⁵ dyn = 0.2248 lb
1 lb = 4.448 N = $4.448 \times 10^5 \text{ dyn}$

Pressure

1 Pa = 1 N/m² = $1.450 \times 10^{-4} \text{ lb/in.}^2$ = 0.0209 lb/ft²
1 bar = 10⁵ Pa
1 lb/in.² = 6895 Pa
1 lb/ft² = 47.88 Pa
1 atm = $1.013 \times 10^5 \text{ Pa}$ = 1.013 bar
= 14.7 lb/in.^2 = 2117 lb/ft²
1 mm Hg = 1 torr = 133.3 Pa

Energy

1 J = 10⁷ ergs = 0.239 cal
1 cal = 4.186 J (based on 15° calorie)
1 ft · lb = 1.356 J
1 Btu = 1055 J = 252 cal = 778 ft · lb
1 eV = $1.602 \times 10^{-19} \text{ J}$
1 kWh = $3.600 \times 10^6 \text{ J}$

Mass–Energy Equivalence

1 kg ↔ $8.988 \times 10^{16} \text{ J}$
1 u ↔ 931.5 MeV
1 eV ↔ $1.074 \times 10^{-9} \text{ u}$

Power

1 W = 1 J/s
1 hp = 746 W = 550 ft · lb/s
1 Btu/h = 0.293 W

APPENDIX F

NUMERICAL CONSTANTS

Fundamental Physical Constants*

Name	Symbol	Value
Speed of light in vacuum	c	$2.99792458 \times 10^8 \text{ m/s}$
Magnitude of charge of electron	e	$1.602176634 \times 10^{-19} \text{ C}$
Gravitational constant	G	$6.67408(31) \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2$
Planck's constant	h	$6.62607015 \times 10^{-34} \text{ J} \cdot \text{s}$
Boltzmann constant	k	$1.380649 \times 10^{-23} \text{ J/K}$
Avogadro's number	N_A	$6.02214076 \times 10^{23} \text{ molecules/mol}$
Gas constant	$R = N_A k$	$8.314462618 \dots \text{ J/mol} \cdot \text{K}$
Mass of electron	m_e	$9.10938356(11) \times 10^{-31} \text{ kg}$
Mass of proton	m_p	$1.672621898(21) \times 10^{-27} \text{ kg}$
Mass of neutron	m_n	$1.674927471(21) \times 10^{-27} \text{ kg}$
Magnetic constant	μ_0	$1.25663706 \times 10^{-6} \text{ Wb/A} \cdot \text{m}$ (approximate) $\cong 4\pi \times 10^{-7} \text{ Wb/A} \cdot \text{m}$
Electric constant	$\epsilon_0 = 1/\mu_0 c^2$	$8.854187817 \times 10^{-12} \text{ C}^2/\text{N} \cdot \text{m}^2$ (approximate)
	$1/4\pi\epsilon_0$	$8.987551787 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$ (approximate)

Other Useful Constants*

Mechanical equivalent of heat		4.186 J/cal (15° calorie)
Standard atmospheric pressure	1 atm	$1.01325 \times 10^5 \text{ Pa}$
Absolute zero	0 K	-273.15°C
Electron volt	1 eV	$1.6021766209(98) \times 10^{-19} \text{ J}$
Atomic mass unit	1 u	$1.660539040(20) \times 10^{-27} \text{ kg}$
Electron rest energy	$m_e c^2$	0.5109989461(31) MeV
Volume of ideal gas (0°C and 1 atm)		22.413962(13) liter/mol
Acceleration due to gravity (standard)	g	9.80665 m/s^2

*Source: National Institute of Standards and Technology (<http://physics.nist.gov/cuu>). Numbers in parentheses show the uncertainty in the final digits of the main number; for example, the number 1.6454(21) means 1.6454 ± 0.0021 . Values shown without uncertainties are exact. The exact values of the magnitude of the charge of the electron, Planck's constant, the Boltzmann constant, Avogadro's number, and the gas constant are from the redefinitions adopted in 2018. As consequences of these redefinitions, the values of the magnetic constant and electric constant now have fractional uncertainties of about 2×10^{-10} . As of this writing (2018) it was expected that updated values of the magnetic and electric constants, as well as of all other constants with uncertainties, were to be announced in May 2019. These updated values will be available at <http://physics.nist.gov/cuu>.

Astronomical Data[†]

Body	Mass (kg)	Radius (m)	Orbit radius (m)	Orbital period
Sun	1.99×10^{30}	6.96×10^8	—	—
Moon	7.35×10^{22}	1.74×10^6	3.84×10^8	27.3 d
Mercury	3.30×10^{23}	2.44×10^6	5.79×10^{10}	88.0 d
Venus	4.87×10^{24}	6.05×10^6	1.08×10^{11}	224.7 d
Earth	5.97×10^{24}	6.37×10^6	1.50×10^{11}	365.3 d
Mars	6.42×10^{23}	3.39×10^6	2.28×10^{11}	687.0 d
Jupiter	1.90×10^{27}	6.99×10^7	7.78×10^{11}	11.86 y
Saturn	5.68×10^{26}	5.82×10^7	1.43×10^{12}	29.45 y
Uranus	8.68×10^{25}	2.54×10^7	2.87×10^{12}	84.02 y
Neptune	1.02×10^{26}	2.46×10^7	4.50×10^{12}	164.8 y
Pluto‡	1.30×10^{22}	1.19×10^6	5.91×10^{12}	248.0 y

[†]Source: NASA (<http://solarsystem.nasa.gov/planets/>). For each body, “radius” is its average radius and “orbit radius” is its average distance from the sun or (for the moon) from the earth.
[‡]In August 2006, the International Astronomical Union reclassified Pluto and similar small objects that orbit the sun as “dwarf planets.”

Prefixes for Powers of 10

Power of ten	Prefix	Abbreviation	Pronunciation
10^{-24}	yocto-	y	yoc-toe
10^{-21}	zepto-	z	zep-toe
10^{-18}	atto-	a	at-toe
10^{-15}	femto-	f	fem-toe
10^{-12}	pico-	p	pee-koe
10^{-9}	nano-	n	nan-oe
10^{-6}	micro-	μ	my-crow
10^{-3}	milli-	m	mil-i
10^{-2}	centi-	c	cen-ti
10^3	kilo-	k	kil-oe
10^6	mega-	M	meg-a
10^9	giga-	G	jig-a or gig-a
10^{12}	tera-	T	ter-a
10^{15}	peta-	P	pet-a
10^{18}	exa-	E	ex-a
10^{21}	zetta-	Z	zet-a
10^{24}	yotta-	Y	yot-a

Examples:

- 1 femtometer = 1 fm = 10^{-15} m
- 1 millivolt = 1 mV = 10^{-3} V
- 1 picosecond = 1 ps = 10^{-12} s
- 1 kilopascal = 1 kPa = 10^3 Pa
- 1 nanocoulomb = 1 nC = 10^{-9} C
- 1 megawatt = 1 MW = 10^6 W
- 1 microkelvin = 1 μ K = 10^{-6} K
- 1 gigahertz = 1 GHz = 10^9 Hz

ANSWERS TO ODD-NUMBERED PROBLEMS

Chapter 1

- 1.1 (a) 1.61 km (b) 3.28×10^3 ft
 1.3 1.02 ns
 1.5 31.7 y
 1.7 (a) 23.4 km/L (b) 1.4 tanks
 1.9 9.0 cm
 1.11 (a) 72 mm² (b) 0.50 (c) 36 mm (d) 6 mm (e) 2.0
 1.13 0.45%
 1.15 (a) no (b) no (c) no (d) no (e) no
 1.17 \$300 million
 1.19 2×10^5
 1.21 7.8 km, 38° north of east
 1.23 144 m, 41° south of west.
 1.25 $A_x = 0$, $A_y = -8.00$ m, $B_x = 7.50$ m, $B_y = 13.0$ cm, $C_x = -10.9$ cm, $C_y = -5.07$ m, $D_x = -7.99$ m, $D_y = 6.02$ m
 1.27 (a) -6.00 m (b) 11.3 m
 1.29 (a) 9.01 m, 33.7° (b) 9.01 m, 33.7° (c) 22.3 m, 250° (d) 22.3 m, 70.3°
 1.31 2.81 km, 38.5° north of west
 1.33 (a) 2.48 cm, 18.4° (b) 4.09 cm, 83.7° (c) 4.09 cm, 264°
 1.35 $\vec{A} = -(8.00 \text{ m})\hat{j}$, $\vec{B} = (7.50 \text{ m})\hat{i} + (13.0 \text{ m})\hat{j}$, $\vec{C} = (-10.9 \text{ m})\hat{i} + (-5.07 \text{ m})\hat{j}$, $\vec{D} = (-7.99 \text{ m})\hat{i} + (6.02 \text{ m})\hat{j}$
 1.37 (a) $\vec{A} = (1.23 \text{ m})\hat{i} + (3.38 \text{ m})\hat{j}$, $\vec{B} = (-2.08 \text{ m})\hat{i} + (-1.20 \text{ m})\hat{j}$ (b) $\vec{C} = (12.0 \text{ m})\hat{i} + (14.9 \text{ m})\hat{j}$ (c) 19.2 m, 51.2°
 1.39 (a) $A = 5.38$, $B = 4.36$ (b) $-5.00\hat{i} + 2.00\hat{j} + 7.00\hat{k}$ (c) 8.83, yes
 1.41 (a) -104 m² (b) -148 m² (c) 40.6 m²
 1.43 (a) 165° (b) 28° (c) 90°
 1.45 (a) $(-63.9 \text{ m}^2)\hat{k}$ (b) $(63.9 \text{ m}^2)\hat{k}$
 1.47 -2.0 m, -x-direction
 1.49 (a) 5.49 g/cm³ (b) 1.1×10^6 g/cm³ (c) 4.7×10^{14} g/cm³
 1.51 (a) 1.64×10^4 km (b) $2.57\tau_E$
 1.53 $\approx 6 \times 10^{27}$
 1.55 (a) $0.60\hat{i} - 0.80\hat{k}$ (b) $-0.60\hat{i} + 0.80\hat{k}$ (c) $\vec{B}_+ = 0.80\hat{i} + 0.60\hat{k}$ and $\vec{B}_- = -0.80\hat{i} - 0.60\hat{k}$
 1.57 144 m, 41° south of west
 1.59 361 N/C, 184.73° from the +x-axis
 1.61 7.55 N
 1.63 60.9 km, 33.0° south of west
 1.65 28.8 m, 11.4° north of east
 1.67 71.9 m, 64.1° north of west
 1.69 (a) 818 m, 15.8° west of south
 1.71 18.6° east of south, 29.6 m
 1.73 28.2 m
 1.75 -2.00 J
 1.77 124°
 1.79 (a) $-bd$, $-bc\hat{i} - ad\hat{j} - ac\hat{k}$ (b) $-bd$, ad in the -y-direction
 1.81 156 m²
 1.83 28.0 m
 1.85 $C_x = -8.0$, $C_y = -6.1$
 1.87 D, F, B, C, A, E
 1.89 (b) (i) 0.9857 AU (ii) 1.3820 AU (iii) 1.695 AU (c) 54.6°
 1.91 (a) 76.2 ly (b) 129°
 1.93 Choice (a)

Chapter 2

- 2.1 25.0 m
 2.3 55 min
 2.5 (a) 0.313 m/s (b) 1.56 m/s
 2.7 (a) 12.0 m/s (b) (i) 0 (ii) 15.0 m/s (iii) 12.0 m/s (c) 13.3 m/s
 2.9 (a) 2.33 m/s, 2.33 m/s (b) 2.33 m/s, 0.33 m/s
 2.11 6.7 m/s, 6.7 m/s, 0, -40.0 m/s, -40.0 m/s, 0
 2.13 (a) 2.00 cm/s, 50.0 cm, -0.125 cm/s² (b) 16.0 s (c) 32.0 s (d) 6.20 s, 1.23 cm/s; 25.8 s, -1.23 cm/s; 36.4 s, -2.55 cm/s
 2.15 (a) 0.500 m/s² (b) 0, 1.00 m/s²
 2.17 (a) 2.17 m, 9.60 m/s²; 15.0 m, -38.4 m/s²
 2.19 (a) 8.33 m/s (b) 1.11 m/s²
 2.21 (a) 675 m/s² (b) 0.0667 s
 2.23 1.70 m
 2.25 0.38 m
 2.27 (a) 3.1×10^6 m/s² = 3.2×10^5 g (b) 1.6 ms (c) no
 2.29 (a) 0, 6.3 m/s², -11.2 m/s² (b) 100 m, 230 m, 320 m
 2.31 (a) 2.94 m/s (b) 0.600 s
 2.33 1.67 s
 2.35 (a) 33.5 m (b) 15.8 m/s
 2.37 (a) upward, downward, decreasing (b) downward, downward, increasing
 2.39 (a) $t = \sqrt{2d/g}$ (b) 0.190 s
 2.41 (a) 646 m (b) 16.4 s, 112 m/s
 2.43 15.9 m
 2.45 0.0868 m/s²
 2.47 37.6 m/s
 2.49 (a) 467 m (b) 110 m/s
 2.51 (a) $x = (0.25 \text{ m/s}^3)t^3 - (0.010 \text{ m/s}^4)t^4$, $v_x = (0.75 \text{ m/s}^3)t^2 - (0.040 \text{ m/s}^4)t^3$ (b) 39.1 m/s
 2.53 (a) 10.0 m (b) (i) 8.33 m/s (ii) 9.09 m/s (iii) 9.52 m/s
 2.55 250 km
 2.57 (a) 197 m/s (b) 169 m/s
 2.59 (a) 4.24 m/s (b) 7.54 s
 2.61 (a) 92.0 m (b) 92.0 m
 2.63 67 m
 2.65 (a) 7.56 s (b) 37.2 m (c) 25.7 m/s (car), 15.9 m/s (truck)
 2.67 3.2 m/s²
 2.69 (a) 2.0 s (b) 16 m/s in the -x-direction, -40 m/s² in the -x-direction
 2.71 (a) -4.00 m/s (b) 12.0 m/s
 2.73 38.2 m
 2.75 (a) 6.69 m/s (b) 4.49 m (c) 1.42 s
 2.77 (a) 3.3 s (b) 9H
 2.79 (a) 380 m (b) 184 m
 2.81 (a) 0.625 m/s³ (b) 107 m
 2.83 (a) car A (b) 2.27 s, 5.73 s (c) 1.00 s, 4.33 s (d) 2.67 s
 2.85 (a) 0.0510 s²/m (b) lower than (c) no
 2.87 4.8
 2.89 (a) 8.3 m/s (b) (i) 0.411 m (ii) 1.15 km (c) 9.8 m/s (d) 4.9 m/s
 2.91 Choice (b)

Chapter 3

- 3.1 (a) 1.4 m/s, -1.3 m/s (b) 1.9 m/s, 317°
 3.3 (a) 7.1 cm/s, 45° (b) 5.0 cm/s, 90°; 7.1 cm/s, 45°; 11 cm/s, 27°

- 3.5 (b) -8.67 m/s², -2.33 m/s² (c) 8.98 m/s², 195°
 3.7 (b) $\vec{v} = \alpha\hat{i} - 2\beta t\hat{j}$, $\vec{a} = -2\beta\hat{j}$ (c) 5.4 m/s, 297°; 2.4 m/s², 270° (d) speeding up and turning right
 3.9 (a) 1.13 m (b) 0.528 m (c) $v_x = 1.10$ m/s, $v_y = -4.70$ m/s, 4.83 m/s, 76.8° below the horizontal
 3.11 2.57 m
 3.13 (a) 24.1 m/s (b) 31.0 m/s
 3.15 1.28 m/s²
 3.17 (a) 0.683 s, 2.99 s (b) 24.0 m/s, 11.3 m/s; 24.0 m/s, -11.3 m/s (c) 30.0 m/s, 36.9° below the horizontal
 3.19 (a) 1.5 m (b) -0.89 m/s
 3.21 (a) 13.6 m (b) 34.6 m/s (c) 103 m
 3.25 (a) $0.034 \text{ m/s}^2 = 0.0034g$ (b) 1.4 h
 3.27 52.8 m/s²
 3.29 120 m/s, 270 mph
 3.31 (a) 2.57 m/s² upward (b) 2.57 m/s² downward (c) 14.7 s
 3.33 (a) 32.9 m/s (b) 27.7 m/s² (c) 35.5 rpm
 3.35 (a) 14 s (b) 70 s
 3.37 0.36 m/s, 52.5° south of west
 3.39 (a) 24° west of south (b) 5.5 h
 3.41 (a) 4.7 m/s, 25° south of east (b) 120 s (c) 240 m
 3.43 (a) 14°, north of west (b) 310 km/h
 3.45 $2b/3c$
 3.47 (a) 128 m (b) 315 m
 3.49 274 m
 3.51 (a) 55.5 m south
 3.53 33.7 m
 3.55 (a) 16.6 m/s (b) 10.9 m/s, 40.5° below the horizontal
 3.57 (a) 1.50 m/s (b) 4.66 m
 3.59 (a) $v = \sqrt{v_0^2 + 2gH}$ (b) $v = \sqrt{v_0^2 + 2gH}$ (c) $\sqrt{v_0^2 + 2gH}$ (d) stays the same
 3.61 (a) 6.91 m (c) no
 3.63 (a) 17.8 m/s (b) in the river, 28.4 m horizontally from his launch point
 3.65 (a) 49.5 m/s (b) 50 m
 3.67 6.25 s
 3.69 (a) 13.3 m/s (b) 3.8 m
 3.71 (a) 44.7 km/h, 26.6° west of south (b) 10.5° north of west
 3.73 7.39 m/s, 12.4° north of east
 3.75 3.01 m/s, 33.7° north of east
 3.77 (a) graph R^2 versus h (b) 16.4 m/s (c) 23.8 m
 3.79 70.5°
 3.81 5.15 s
 3.83 Choice (b)
 3.85 Choice (c)

Chapter 4

- 4.1 494 N, 31.8°
 4.3 3.15 N
 4.5 (a) -8.10 N, 3.00 N (b) 8.64 N
 4.7 46.7 N, opposite to the motion of the skater
 4.9 21.8 kg
 4.11 (a) 3.12 m, 3.12 m/s (b) 21.9 m, 6.24 m/s
 4.13 (a) 45.0 N, between 2.0 s and 4.0 s (b) between 2.0 s and 4.0 s (c) 0 s, 6.0 s
 4.15 (a) $A = 100$ N, $B = 12.5$ N/s² (b) (i) 21.6 N, 2.70 m/s² (ii) 134 N, 16.8 m/s² (c) 26.6 m/s²

- 4.17 2940 N
 4.19 (a) 4.49 kg (b) 4.49 kg, 8.13 N
 4.21 825 N, blocks
 4.23 50 N
 4.25 7.4×10^{-23} m/s²
 4.27 (b) yes
 4.29 (a) yes (b) no
 4.33 2.58 s
 4.35 (a) 17 N, 90° clockwise from the +x-axis
 (b) 840 N
 4.37 (a) 2.50 m/s² (b) 10.0 N (c) to the right,
 $F > T$ (d) 25.0 N
 4.39 (a) 4.4 m (b) 300 m/s (c) (i) 2.7×10^4 N
 (ii) 9.0×10^3 N
 4.41 (a) 0.603 m/s², upward
 (b) 1.26 m/s², downward
 4.43 1630 N away from him
 4.45 (a) 4.34 kg (b) 5.30 kg
 4.47 309 N
 4.49 7.78 m
 4.51 (a) largest: Ferrari; smallest: Alpha Romeo and
 Honda Civic (b) largest: Ferrari; smallest:
 Volvo (c) 7.5 kN, smaller (d) zero
 4.53 (b) 26 kg, 8.3 m/s²
 4.55 (a) 0.593 m, 1.33 s, 4.00 N (b) 2.00 s, 2.00 m/s
 4.57 Choice (d)
 4.59 Choice (a)

Chapter 5

- 5.1 (a) 25.0 N (b) 50.0 N
 5.3 (a) 990 N, 735 N (b) 926 N
 5.5 48°
 5.7 (a) $T_A = 0.732w$, $T_B = 0.897w$, $T_C = w$
 (b) $T_A = 2.73w$, $T_B = 3.35w$, $T_C = w$
 5.9 (a) 574 N (b) 607 N
 5.11 1.10×10^8 N (b) 5w (c) 8.4 s
 5.13 (a) 4610 m/s² = 470g
 (b) 9.70×10^5 N = 471w (c) 0.0187 s
 5.15 (b) 2.96 m/s² (c) 191 N; greater than; less than
 5.17 (b) 3.75 m/s² (c) 2.48 kg (d) $T <$ weight of the
 hanging block
 5.19 (a) 3.4 m/s (c) 2.2w
 5.21 (a) 14.0 m (b) 18.0 m/s
 5.23 50°
 5.25 (a) 33 N (b) 3.1 m
 5.27 (a) μ_k : 0.710; μ_k : 0.472 (b) 258 N (c) (i) 51.8 N
 (ii) 4.97 m/s²
 5.31 (a) 18.3 m/s² (b) 2.29 m/s²
 5.33 (a) 57.1 N (b) 146 N up the ramp
 5.35 (a) 52.5 m (b) 16.0 m/s
 5.37 (a) $\mu_k(m_A + m_B)g$ (b) $\mu_k m_A g$
 5.39 (a) 0.218 m/s (b) 11.7 N
 5.41 (a) $\frac{\mu_k mg}{\cos\theta - \mu_k \sin\theta}$ (b) $1/\tan\theta$
 5.43 (b) 8.2 m/s
 5.45 (a) 61.8 N (b) 30.4 N
 5.47 3.66 s
 5.49 (a) 21.0°, no (b) 11,800 N (car), 23,600 N (truck)
 5.51 6200 N (horizontal cable), 1410 (upper cable)
 5.53 (a) 1.5 rev/min (b) 0.92 rev/min
 5.55 (a) $\sqrt{Lg}$ (b) $5mg$
 5.57 2.42 m/s
 5.59 (a) rope making 60° angle (b) 6400 N
 5.61 $T_B = 4960$ N, $T_C = 1200$ N
 5.63 3.18 m/s
 5.65 (a) 101 N (b) 33.8 N (c) 62.3 N
 5.67 762 N
 5.69 (a) (i) -3.80 m/s (ii) 24.6 m/s
 (b) 4.36 m (c) 2.45 s

- 5.71 245 N, equal to it
 5.73 12.3 m/s
 5.75 1.78 m/s
 5.77 (a) $m_1(\sin\alpha + \mu_k \cos\alpha)$ (b) $m_1(\sin\alpha - \mu_k \cos\alpha)$
 (c) $m_1(\sin\alpha - \mu_s \cos\alpha) \leq m_2 \leq m_1(\sin\alpha + \mu_s \cos\alpha)$
 5.79 (a) 1.44 N (b) 1.80 N
 5.81 920 N
 5.83 1.30 kg, 2.19 kg
 5.85 (a) 294 N (18.0-cm wire), 152 N, 152 N
 (b) 40.0 N
 5.87 3.0 N
 5.89 (a) 12.9 kg (b) $T_{AB} = 47.2$ N, $T_{BC} = 101$ N
 5.91 $a_1 = \frac{2m_2g}{4m_1 + m_2}$, $a_2 = \frac{m_2g}{4m_1 + m_2}$
 5.93 (a) 7.75 N (b) 0.242
 5.95 g/μ_s
 5.97 (b) 0.452
 5.99 (b) 0.28 (c) no
 5.101 (a) 1.42×10^4 N (b) 817 m/s² (c) 27.1 m/s²
 5.103 (b) 8.8 N (c) 31.0 N (d) 1.54 m/s²
 5.105 $v = (2mg/k)\left[\frac{1}{2} + e^{-(k/m)t}\right]$
 5.107 (a) 81.1° (b) no (c) The bead rides at the bottom
 of the hoop.
 5.109 (a) 0.371 (b) 0.290 (c) yes, same slope,
 less-negative intercept
 5.111 (a) 5/8 in (b) 23.9 kN (c) 3.57 kN, smaller
 (d) larger; accurate
 5.113 $F = (M + m)g \tan\alpha$
 5.115 $\cos^2\beta$
 5.117 Choice (b)

Chapter 6

- 6.1 (a) 3.60 J (b) -0.900 J (c) 0 (d) 0
 (e) 2.70 J
 6.3 (a) 74 N (b) 333 J (c) -330 J (d) 0, 0
 (e) 0
 6.5 (a) -1750 J (b) no
 6.7 (a) (i) 9.00 J (ii) -9.00 J (b) (i) 0
 (ii) 9.00 J (iii) -9.00 J (iv) 0
 (c) zero for each block
 6.9 -6.7 J
 6.11 (a) 0° (b) 4
 6.13 -572 J
 6.15 (a) 120 J (b) -108 J (c) 24.3 J
 6.17 (a) 36,000 J (b) 4
 6.19 (a) 1.0×10^{16} J (b) 2.4 times
 6.21 $4W_1$
 6.23 $\sqrt{2gh}(1 + \mu_k/\tan\alpha)$
 6.25 48.0 N
 6.27 (a) 4.48 m/s (b) 3.61 m/s
 6.29 (a) twice as fast (b) both decrease by a
 factor of $1/\sqrt{3}$
 6.31 (a) $v_0^2/2\mu_k g$ (b) (i) $1/2$ (ii) 4 (iii) 2
 6.33 (b) 13.1 cm (bottom), 14.1 cm (middle),
 15.2 cm (top)
 6.35 (a) 2.83 m/s (b) 3.46 m/s
 6.37 8.5 cm
 6.39 (a) 1.76 (b) 0.666 m/s
 6.41 (a) 4.0 J (b) 0 (c) -1.0 J (d) 3.0 J (e) -1.0 J
 6.43 (a) 2.83 m/s (b) 2.40 m/s
 6.45 8.17 m/s
 6.47 360,000 J; 100 m/s
 6.53 (a) 84.6/min (b) 22.7/min
 6.55 29.6 kW
 6.57 0.20 W
 6.59 (a) 608 J (b) -395 J (c) 0 (d) -189 J
 (e) 24 J (f) 1.5 m/s
 6.61 (a) -35.5 J (b) decreases (c) -51.6 J

- 6.63 (a) 5.62 J (20.0-N block), 3.38 J (12.0-N block)
 (b) 2.58 J (20.0-N block), 1.54 J (12.0-N block)
 6.65 (a) 1.8 m/s = 4.0 mi/h
 (b) $180 \text{ m/s}^2 \approx 18g$, 900 N
 6.67 (a) 5.11 m (b) 0.304 (c) 10.3 m
 6.69 (a) $V/\sqrt{2}$, greater than (b) $W_B/4$, less than
 6.71 (a) 0.074 N (b) 4.7 N (c) 0.22 J
 6.73 6.3×10^4 N/m
 6.75 1.1 m
 6.77 (a) 2.39 m/s (b) 9.42 m/s, away from the wall
 6.79 (a) 0.600 m (b) 1.50 m/s
 6.81 0.786
 6.83 1.3 m
 6.85 (a) 1.10×10^5 J (b) 1.30×10^5 J (c) 3.99 kW
 6.87 (a) 47.0 J, -29.6 J, 2.41 m/s
 (b) 23.2 J, 29.6 J, -6.4 J
 (c) 40.6 J, 47.0 J, -6.4 J
 6.89 (a) 1.26×10^5 J (b) 1.46 W
 6.91 (b) $v^2 = -\frac{k}{m}d^2 + 2d\left[\frac{k}{m}(0.400 \text{ m}) - \mu_k g\right]$
 (c) 1.29 m/s, 0.204 m (d) 12.0 N/m, 0.800
 6.93 (a) $Mv^2/6$ (b) 6.1 m/s (c) 3.9 m/s
 (d) 0.40 J, 0.60 J
 6.95 Choice (a)
 6.97 Choice (d)

Chapter 7

- 7.1 (a) 6.6×10^5 J (b) -7.7×10^5 J
 7.3 (a) 610 N (b) (i) 0 (ii) 550 J
 7.5 (a) 24.0 m/s (b) 24.0 m/s (c) part (b)
 7.7 (a) 2.0 m/s (b) 9.8×10^{-7} J, 2.0 J/kg
 (c) 200 m, 63 m/s (d) 5.9 J/kg
 (e) in its tensed legs
 7.9 (a) (i) 0 (ii) 0.98 J (b) 2.8 m/s
 (c) Only gravity is constant. (d) 5.1 N
 -5400 J
 7.11 -5400 J
 7.13 (a) 11.8 J, -15.7 J (b) (0.200 m) T , -(0.200 m) T
 (c) 0, -3.9 J, 0.75 m/s
 7.15 (a) 52.0 J (b) 3.25 J
 7.17 (a) (i) $4U_0$ (ii) $U_0/4$ (b) (i) $x_0\sqrt{2}$
 (ii) $x_0/\sqrt{2}$
 7.19 (a) 5.48 cm (b) 3.92 cm
 7.21 (a) 6.32 cm (b) 12 cm
 7.23 (a) 3.03 m/s, as it leaves the spring
 (b) 95.9 m/s², when the spring has its maximum
 compression
 7.25 (a) 4.46×10^5 N/m (b) 0.128 m
 7.27 (a) -5.4 J (b) -5.4 J (c) -10.8 J
 (d) nonconservative
 7.29 (a) 8.16 m/s (b) 766 J
 7.31 1.29 N, +x-direction
 7.33 130 m/s^2 , 132° counterclockwise from the
 +x-axis
 7.35 (a) $F(r) = (12a/r^{13}) - (6b/r^7)$
 (b) $(2a/b)^{1/6}$, yes (c) $b^2/4a$
 (d) $a = 6.67 \times 10^{-138} \text{ J} \cdot \text{m}^{12}$,
 $b = 6.41 \times 10^{-78} \text{ J} \cdot \text{m}^6$
 7.37 (a) zero (gravel), 637 N (box) (b) 2.99 m/s
 7.39 0.41
 7.41 (a) 16.0 m/s (b) 11,500 N
 7.43 (a) 20.0 m along the rough bottom (b) -78.4 J
 7.45 (a) 22.2 m/s (b) 16.4 m (c) no
 7.47 10.8 J
 7.49 15.5 m/s
 7.51 4.4 m/s
 7.53 (a) 7.00 m/s (b) 8.82 N
 7.55 48.2°
 7.57 (a) 0.392 (b) -0.83 J
 7.59 (a) $U(x) = \frac{1}{2}\alpha x^2 + \frac{1}{3}\beta x^3$ (b) 7.85 m/s
 7.61 (a) $\alpha/(x + x_0)$ (b) 3.27 m/s

- 7.63 7.01 m/s
 7.65 (a) 0.747 m/s (b) 0.931 m/s
 7.67 (a) 0.480 m/s (b) 0.566 m/s
 7.69 (a) 3.87 m/s (b) 0.10 m
 7.71 0.456 N
 7.73 119 J
 7.75 (a) -50.6 J (b) -67.5 J (c) nonconservative
 7.77 (a) 57.0 m (b) 16.5 m (c) negative work done by air resistance
 7.79 (a) yes (b) 0.14 J
 (d) -1.0 m, 0, 1.0 m
 (e) positive: $-1.5 \text{ m} < x < -1.0 \text{ m}$ and $0 < x < 1.0 \text{ m}$
 negative: $-1.0 \text{ m} < x < 0$ and $1.0 \text{ m} < x < 1.5 \text{ m}$ (f) -0.55 m, 0.12 J
 7.81 Choice (c)
 7.83 Choice (b)

Chapter 8

- 8.1 (a) $1.20 \times 10^5 \text{ kg} \cdot \text{m/s}$ (b) (i) 60.0 m/s
 (ii) 26.8 m/s
 8.3 (a) $-30 \text{ kg} \cdot \text{m/s}$, $-55 \text{ kg} \cdot \text{m/s}$
 (b) 0, $52 \text{ kg} \cdot \text{m/s}$ (c) 0, $-3.0 \text{ kg} \cdot \text{m/s}$
 8.5 (a) $22.5 \text{ kg} \cdot \text{m/s}$, to the left (b) 838 J
 8.7 562 N, not significant
 8.9 (a) 10.8 m/s, to the right (b) 0.750 m/s, to the left
 8.11 (a) 500 N/s^2 (b) $5810 \text{ N} \cdot \text{s}$ (c) 2.70 m/s
 8.13 (a) $2.50 \text{ N} \cdot \text{s}$, in the direction of the force
 (b) (i) 6.25 m/s, to the right (ii) 3.75 m/s, to the right
 8.15 (a) $8.00 \times 10^2 \text{ kg} \cdot \text{m/s}$ (b) $8.00 \times 10^3 \text{ J}$
 (c) 160 N opposite to her velocity
 8.17 $0.87 \text{ kg} \cdot \text{m/s}$, in the same direction as the bullet is traveling
 8.19 (a) 6.79 m/s (b) 55.2 J
 8.21 (a) 0.790 m/s (b) -0.0023 J
 8.23 1.97 m/s
 8.25 (a) 0.0559 m/s (b) 0.0313 m/s
 8.27 (a) 7.20 m/s, 38.0° from Rebecca's original direction (b) -680 J
 8.29 (a) 4.3 m/s (c) 4.3 m/s
 8.31 0.31 m/s toward the south
 8.33 (a) 0.846 m/s (b) 2.10 J
 8.35 13.5 J
 8.37 5.9 m/s, 58° north of east
 8.39 5.46 m/s, 36.0° south of east
 8.41 19.5 m/s (car), 21.9 m/s (truck)
 8.43 (a) 2.93 cm (b) 866 J (c) 1.73 J
 8.45 13.6 N
 8.47 (a) 3.00 J; 0.500 m/s for both
 (b) A: -1.00 m/s ; B: 1.00 m/s
 8.49 (a) $v_i/3$ (b) $K_i/9$ (c) 10
 8.51 (a) B (b) 3 times greater (c) 3.0 m/s to the left
 8.53 2520 km
 8.55 0.488 m
 8.57 0.73 m/s
 8.59 $F_x = -(1.50 \text{ N/s})t$, $F_y = 0.25 \text{ N}$, $F_z = 0$
 8.61 (a) 0.053 kg (b) 5.19 N
 8.65 (a) $-1.14 \text{ N} \cdot \text{s}$, $0.330 \text{ N} \cdot \text{s}$
 (b) 0.04 m/s, 1.8 m/s
 8.67 (a) 5.21 J, -0.0833 m/s
 (b) -2.17 m/s (A), 0.333 m/s (B)
 8.69 (a) 1.75 m/s, 0.260 m/s (b) -0.092 J
 8.71 0.946 m
 8.73 1.8 m
 8.75 $v_A/2$ for each one
 8.77 12 m/s (SUV), 21 m/s (sedan)
 8.79 (a) 2.60 m/s (b) 325 m/s
 8.81 (a) 5.3 m/s (b) 5.7 m
 8.83 (a) 1 (b) $3 - 2\sqrt{2}$, $3 + 2\sqrt{2}$
 8.85 (a) 0.0781 (b) 248 J (c) 0.441 J
 8.87 (a) 9.35 m/s (b) 3.29 m/s
 8.89 4.04 m/s
 8.91 1.33 m
 8.93 $H/9$
 8.95 (a) 0.150 J (b) 15.4 J (c) 0.560 m
 8.97 (a) 71.6 m/s (0.28-kg piece), 14.3 m/s (1.40-kg piece) (b) 347 m
 8.99 (a) yes (b) no, decreases by 4800 J
 8.101 (a) maximum: C, minimum: B (b) 69 N/m
 (c) 0.12 m
 8.103 (a) $g/3$ (b) 14.7 m (c) 29.4 g
 8.105 $0, 4a/3\pi$
 8.107 Choice (b)
 8.109 Choice (b)

Chapter 9

- 9.1 (a) 0.600 rad, 34.4° (b) 6.27 cm (c) 1.05 m
 9.3 (a) rad/s, rad/s^3 (b) (i) 0 (ii) 15.0 rad/s^2
 (c) 9.50 rad
 9.5 (a) $\omega_z = \gamma + 3\beta t^2$ (b) 0.400 rad/s
 (c) 1.30 rad/s, 0.700 rad/s
 9.7 (a) $\pi/4 \text{ rad}$, 2.00 rad/s , -0.139 rad/s^3
 (b) 0 (c) 19.5 rad, 9.36 rad/s
 9.9 (a) 2.00 rad/s (b) 4.38 rad
 9.11 (a) 24.0 s (b) 68.8 rev
 9.13 3.00 rad/s
 9.15 (a) 300 rpm (b) 75.0 s, 312 rev
 9.17 9.00 rev
 9.21 (a) 15.1 m/s^2 (b) 15.1 m/s^2
 9.25 0.107 m, no
 9.27 5.97 rad/s^2
 9.29 0.180 m/s^2 , 0.432 m/s^2 , 0.468 m/s^2
 9.31 (a) (i) $0.469 \text{ kg} \cdot \text{m}^2$ (ii) $0.117 \text{ kg} \cdot \text{m}^2$ (iii) 0
 (b) (i) $0.0433 \text{ kg} \cdot \text{m}^2$ (ii) $0.0722 \text{ kg} \cdot \text{m}^2$
 (c) (i) $0.0288 \text{ kg} \cdot \text{m}^2$ (ii) $0.0144 \text{ kg} \cdot \text{m}^2$
 9.33 (a) $1.93 \text{ kg} \cdot \text{m}^2$ (b) $6.53 \text{ kg} \cdot \text{m}^2$
 (c) 0 (d) $1.15 \text{ kg} \cdot \text{m}^2$
 9.35 $0.193 \text{ kg} \cdot \text{m}^2$
 9.37 0.208 s
 9.39 $\omega_1 \sqrt{3/5}$
 9.41 6.49 m/s
 9.43 (a) wheel B (b) $\frac{3}{16}$
 9.45 $7.35 \times 10^4 \text{ J}$
 9.47 (a) 0.673 m (b) 45.5%
 9.49 46.5 kg
 9.51 an axis that is parallel to a diameter and is 0.516R from the center
 9.53 $M(a^2 + b^2)/3$
 9.55 $\frac{1}{2}MR^2$
 9.57 (a) $\gamma L^2/2$ (b) $ML^2/2$ (c) $ML^2/6$
 9.59 7.68 m
 9.61 (a) 0.600 m/s^3 (b) $\alpha = (2.40 \text{ rad/s}^3)t$ (c) 3.54 s
 (d) 17.7 rad
 9.63 1.22 kg
 9.65 8.44 kg
 9.67 2.99 cm
 9.69 68.4 J
 9.71 $4.65 \text{ kg} \cdot \text{m}^2$
 9.73 (a) -0.882 J (b) 5.42 rad/s (c) 5.42 m/s
 (d) 5.42 m/s compared to 4.43 m/s
 9.75 1.46 m/s
 9.77 $\sqrt{\frac{2gd(m_B - \mu_k m_A)}{m_A + m_B + l/R^2}}$
 9.79 (a) $2.25 \times 10^{-3} \text{ kg} \cdot \text{m}^2$ (b) 3.40 m/s
 (c) 4.95 m/s
 9.81 13.9 m
 9.85 (a) 55.3 kg (b) $0.804 \text{ kg} \cdot \text{m}^2$
 9.87 (a) 4.0 rev, no (b) 15 rad/s (c) 9.5 rad/s
 9.89 (a) yes (b) 3.15 m/s (c) $0.348 \text{ kg} \cdot \text{m}^2$
 (d) 36.4 N

- 9.91 (a) $s(\theta) = r_0\theta + \frac{\beta}{2}\theta^2$
 (b) $\theta(t) = \frac{1}{\beta}(\sqrt{r_0^2 + 2\beta vt} - r_0)$
 (c) $\omega_z(t) = \frac{v}{\sqrt{r_0^2 + 2\beta vt}}$,
 $\alpha_z(t) = -\frac{\beta v t}{(r_0^2 + 2\beta vt)^{3/2}}$, no
 (d) 25.0 mm, $0.247 \mu\text{m/rad}$, $2.13 \times 10^4 \text{ rev}$
 9.93 Choice (d)
 9.95 Choice (d)

Chapter 10

- 10.1 (a) $40.0 \text{ N} \cdot \text{m}$, out of the page (b) $34.6 \text{ N} \cdot \text{m}$, out of the page (c) $20.0 \text{ N} \cdot \text{m}$, out of the page
 (d) $17.3 \text{ N} \cdot \text{m}$, into the page (e) 0 (f) 0
 10.3 $2.50 \text{ N} \cdot \text{m}$, out of the page
 10.5 (b) $-\hat{k}$ (c) $(-1.05 \text{ N} \cdot \text{m})\hat{k}$
 10.7 (a) $2.56 \text{ N} \cdot \text{m}$ (b) $4.25 \text{ N} \cdot \text{m}$, perpendicular to handle
 10.9 $8.38 \text{ N} \cdot \text{m}$
 10.11 (a) 14.8 rad/s^2 (b) 1.52 s
 10.13 (a) 7.5 N (at book on table), 18.2 N (at hanging book) (b) $0.16 \text{ kg} \cdot \text{m}^2$
 10.15 $0.255 \text{ kg} \cdot \text{m}^2$
 10.17 $0.0131 \text{ kg} \cdot \text{m}^2$
 10.19 (a) 1.56 m/s (b) 5.35 J
 (c) (i) 3.12 m/s to the right (ii) 0 (iii) 2.21 m/s at 45° below the horizontal
 (d) (i) 1.56 m/s to the right (ii) 1.56 m/s to the left (iii) 1.56 m/s downward
 10.21 (a) $\frac{1}{3}$ (b) $\frac{2}{7}$ (c) $\frac{2}{5}$ (d) $\frac{5}{13}$
 10.23 (a) 0.613 (b) no (c) no slipping
 10.25 14.0 m
 10.29 (a) 3.76 m (b) 8.58 m/s
 10.31 (a) 67.9 rad/s (b) 8.35 J
 10.33 (a) 0.309 rad/s (b) 100 J (c) 6.67 W
 10.35 (a) $0.704 \text{ N} \cdot \text{m}$ (b) 157 rad (c) 111 J (d) 111 J
 10.37 (a) $115 \text{ kg} \cdot \text{m}^2/\text{s}$ into the page (b) $125 \text{ kg} \cdot \text{m}^2/\text{s}^2$ out of the page
 10.39 $4.71 \times 10^{-6} \text{ kg} \cdot \text{m}^2/\text{s}$
 10.41 (a) A: rad/s^2 ; B: rad/s^4 (b) (i) $59.0 \text{ kg} \cdot \text{m}^2/\text{s}$ (ii) $56.1 \text{ N} \cdot \text{m}$
 10.43 4600 rad/s
 10.45 1.14 rev/s
 10.47 (a) 1.38 rad/s (b) 1080 J, 495 J
 10.49 (a) 0.120 rad/s (b) $3.20 \times 10^{-4} \text{ J}$ (c) work done by the bug
 10.51 4.82 m/s^2
 10.53 $\frac{m_{\text{rock}} v}{\left(m + \frac{M}{2}\right)R}$
 10.55 $2.4 \times 10^{-12} \text{ N} \cdot \text{m}$
 10.59 0.483
 10.61 (a) 16.3 rad/s^2 (b) no, decrease (c) 5.70 rad/s
 10.63 0.921 m/s^2 , 7.68 rad/s^2 , 35.5 N (at A), 21.4 N (at B)
 10.65 $\frac{7v_{\text{cm}}^2}{10g}$, $\frac{5v_{\text{cm}}^2}{6g}$, hollow
 10.67 68.0 N
 10.69 270 N
 10.71 $a = \frac{2g}{2 + (R/b)^2}$, $\alpha = \frac{2g}{2b + R^2/b}$,
 $T = \frac{2mg}{2(b/R)^2 + 1}$
 10.73 (a) $3H_0/5$
 10.75 29.0 m/s
 10.77 (a) 26.0 m/s (b) no change
 10.79 $g/3$

A-12 Answers to Odd-Numbered Problems

- 10.81 (a) $\frac{6}{19}v/L$ (b) $\frac{3}{19}$
 10.83 3200 J
 10.85 (a) 2.00 rad/s (b) 6.58 rad/s
 10.87 0.776 rad/s
 10.89 (a) A: solid sphere, B: solid cylinder, C: hollow sphere, D: hollow cylinder (b) same (c) D (d) 0.350
 10.91 (a) $mv_1^2 r_1^2 / r^3$ (b) $\frac{mv_1^2}{2} r_1^2 \left(\frac{1}{r_2^2} - \frac{1}{r_1^2} \right)$ (c) same
 10.93 (a) 39.2 N upward, 39.2 N upward (b) 60.0 N upward, 18.4 N upward (c) 165 N upward, 86.2 N downward (d) 0.0940 rev/s
 10.95 Choice (c)
 10.97 Choice (a)

Chapter 11

- 11.1 29.8 cm
 11.3 1.35 m
 11.5 9.70 cm
 11.7 6.6 kN
 11.9 (a) 1000 N, 0.800 m from end where 600-N force is applied (b) 800 N, 0.75 m from end where 600-N force is applied
 11.11 (a) 550 N (b) 0.614 m from A
 11.13 (a) 1920 N (b) 1140 N
 11.15 (a) $T = 2.60w$; 3.28w, 37.6° (b) $T = 4.10w$; 5.39w, 48.8°
 11.17 (a) 3410 N (b) 3410 N, 7600 N
 11.19 (b) 530 N (c) 600 N, 270 N; downward
 11.21 220 N (left), 255 N (right), 42°
 11.23 (a) 0.800 m (b) clockwise (c) 0.800 m, clockwise
 11.25 6.12 kg
 11.27 1.9 mm
 11.29 2.0×10^{11} Pa
 11.31 2.0×10^3 atm
 11.33 (a) 150 atm (b) 1.5 km, no
 11.35 4.8×10^9 Pa, 2.1×10^{-10} Pa⁻¹
 11.37 (a) 2.4×10^{-2} (b) 2.4×10^{-3} m
 11.39 $1.4T_{\text{steel}}$
 11.41 $0.19\Delta p_1$
 11.43 3.41×10^7 Pa
 11.47 20.0 kg
 11.49 (a) 525 N (b) 222 N, 328 N (c) 1.48
 11.51 (a) 140 N (b) 6 cm to the right
 11.53 (a) 409 N (b) 161 N
 11.55 49.9 cm
 11.57 (a) $\arctan(mg/2T)$ (b) $\sqrt{\frac{3g \sin \theta}{L}}$
 11.59 (a) $\arctan\left[\left(\frac{1}{\alpha} - 1\right) \tan \theta\right]$ (b) 1/2 (c) 0
 11.61 5500 N
 11.63 (b) 2000 N = 2.72mg (c) 4.4 mm
 11.65 (a) 1.39 m (b) 1.8×10^{-3} m
 11.67 (a) 175 N at each hand, 200 N at each foot (b) 91 N at each hand and at each foot
 11.69 (a) 1150 N (b) 1940 N (c) 918 N (d) 0.473
 11.71 590 N (person above), 1370 N (person below); person above
 11.73 (a) $\frac{T_{\text{max}} h D}{L \sqrt{h^2 + D^2}}$
 (b) $\frac{T_{\text{max}} h}{L \sqrt{h^2 + D^2}} \left(1 - \frac{D^2}{h^2 + D^2} \right)$, positive
 11.75 (a) 375 N (b) 325 N (c) 512 N
 11.77 (a) 0.424 N (A), 1.47 N (B), 0.424 N (C) (b) 0.848 N
 11.79 (a) 27° to tip, 31° to slip, tips first (b) 27° to tip, 22° to slip, slips first
 11.81 (a) 80 N (A), 870 N (B) (b) 1.92 m
 11.83 (a) 0.70 m from A (b) 0.60 m from A

- 11.85 (a) 4.2×10^4 N (b) 65 m
 11.87 (b) $x = 1.50 \text{ m} + \frac{(1.30 \text{ m})m_1 - (0.38 \text{ m})M}{m_2}$
 (c) 1.59 kg (d) 1.50 m
 11.89 (a) 391 N (4.00-m ladder), 449 N (3.00-m ladder) (b) 322 N (c) 334 N (d) 937 N
 11.91 (a) 0.66 mm (b) 0.022 J (c) 8.35×10^{-3} J (d) -3.04×10^{-2} J (e) 3.04×10^{-2} J
 11.93 Choice (a)
 11.95 Choice (d)

Chapter 12

- 12.1 no (41.8 N)
 12.3 7020 kg/m³; yes
 12.5 1.6
 12.7 (a) 1.86×10^6 Pa (b) 184 m
 12.9 0.581 m
 12.11 828 kg/m³
 12.13 2.8 m
 12.15 6.0×10^4 Pa
 12.17 2.27×10^5 N
 12.19 (a) 636 Pa (b) (i) 1170 Pa (ii) 1170 Pa
 12.21 10.9
 12.23 10.3 m
 12.25 0.0898 m³, 1020 kg/m³
 12.27 0.122 m
 12.29 6.43×10^{-4} m³, 2.78×10^3 kg/m³
 12.31 10.5 N
 12.33 (a) 116 Pa (b) 921 Pa (c) 0.822 kg, 822 kg/m³
 12.35 1640 kg/m³
 12.37 501 N
 12.39 9.6 m/s
 12.41 (a) 17.0 m/s (b) 0.317 m
 12.43 (a) 0.536 mm (b) 30.8 s, yes
 12.45 28.4 m/s
 12.47 1.47×10^5 Pa
 12.49 2.03×10^4 Pa
 12.51 2.25×10^5 Pa
 12.53 1.19D
 12.55 (a) 5.9×10^5 N (b) 1.8×10^5 N
 12.57 2.61×10^4 N · m
 12.59 0.800 kg
 12.61 0.964 cm, rise
 12.63 (a) 1470 Pa (b) 13.9 cm
 12.65 (a) 0.0500 m³ (b) 10.0 kg
 12.67 9.8×10^6 kg, yes
 12.71 (a) 16.5 cm (b) 1.75 m
 12.73 (a) 5.07 m/s, 1.28 (b) 32.4 min, 2.08
 12.75 (a) $1 - \frac{\rho_B}{\rho_L}$ (b) $\left(\frac{\rho_L - \rho_B}{\rho_L - \rho_w} \right) L$ (c) 4.60 cm
 12.77 (a) $2\sqrt{h(H-h)}$ (b) h
 12.79 5.47 m
 12.81 (a) 0.200 m³/s (b) 6.97×10^4 Pa
 12.83 $3h_1$
 12.85 (b) no
 12.87 (a) 2.5×10^{-4} m²/Pa (slope), 16 m² (intercept) (b) 8.2 m, 800 kg/m³
 12.89 Choice (b)
 12.91 Choice (a)

Chapter 13

- 13.1 2.18
 13.3 (a) 1.2×10^{-11} m/s² (b) 15 days (c) no, increase
 13.5 2.1×10^{-9} m/s², downward
 13.7 (a) 2.4×10^{-3} N (b) $F_{\text{moon}}/F_{\text{earth}} = 3.5 \times 10^{-6}$
 13.9 (a) 0.634 m from 3m (b) (i) unstable (ii) stable
 13.11 1.38×10^7 m
 13.13 (a) 0.37 m/s² (b) 1700 kg/m³
 13.15 (a) 12.3 m/s² (b) 80.0% the density of earth

- 13.17 (a) 5020 m/s (b) 60,600 m/s
 13.19 9.03 m/s²
 13.21 2.40×10^{12} N
 13.23 -2.00×10^6 J, -4.00×10^6 J
 13.25 (a) 7410 m/s (b) 1.71 h
 13.27 7330 m/s
 13.29 (a) 4.1 m/s = 9.1 mph, yes (b) 2.6 h
 13.31 (a) 82,700 m/s (b) 14.5 days
 13.33 (a) 7.84×10^9 s = 248 y (b) 4.44×10^{12} m, 7.38×10^{12} m
 13.35 (a) 0 (b) 1.07×10^{-12} N (c) 1.25×10^{-12} N
 13.37 (a) (i) 5.31×10^{-9} N (ii) 2.67×10^{-9} N
 13.39 (a) $-\frac{GmM}{\sqrt{x^2 + a^2}}$ (b) $-GmM/x$
 (c) $\frac{GmMx}{(x^2 + a^2)^{3/2}}$, toward the ring (d) GmM/x^2
 (e) $U = -GmM/a$, $F_x = 0$
 13.41 (a) 33.7 N (b) 32.8 N
 13.43 (a) 4.3×10^{37} kg = $(2.1 \times 10^7)M_{\text{S}}$ (b) no
 (c) 6.32×10^{10} m, yes
 13.45 (a) 9.67×10^{-12} N, at 45° above +x-axis (b) 3.02×10^{-5} m/s
 13.47 (b) (i) 1.49×10^{-5} m/s (50.0-kg sphere), 7.46×10^{-6} m/s (100.0-kg sphere) (ii) 2.24×10^{-5} m/s (c) 26.6 m
 13.49 (a) 3.59×10^7 m
 13.51 177 m/s
 13.53 (a) 7.36 h (b) 2.47 h
 13.55 1.83×10^{27} kg
 13.57 22.8 m
 13.59 $4\pi GM$
 13.61 $v = \sqrt{\frac{2Gm_E h}{R_E(R_E + h)}}$
 13.63 (a) $GM^2/4R^2$ (b) $v = \sqrt{GM/4R}$, $T = 4\pi\sqrt{R^3/GM}$ (c) $GM^2/4R$
 13.65 6.8×10^4 m/s
 13.67 (a) 7910 s (b) 1.53 (c) 8430 m/s (perigee), 5510 m/s (apogee) (d) 2410 m/s; 3250 m/s; perigee
 13.69 5.36×10^9 J
 13.71 9.36 m/s²
 13.73 $GmMx/(a^2 + x^2)^{3/2}$
 13.75 (a) $U(r) = \frac{Gm_E m}{2R_E^3} r^2$ (b) 7.90×10^3 m/s
 13.77 (a) It is considerable and shows no apparent pattern. (b) Earth (5500 kg/m³), Mercury (5400 kg/m³), Venus (5300 kg/m³), Mars (3900 kg/m³), Neptune (1600 kg/m³), Uranus (1200 kg/m³), Jupiter (1200 kg/m³), Saturn (530 kg/m³) (c) no effect (d) 90 m/s²
 13.79 (a) opposite; opposite (b) 259 days (c) 44.1°
 13.81 $\frac{2GmM}{a^2} \left(1 - \frac{x}{\sqrt{a^2 + x^2}} \right)$
 13.83 Choice (c)

Chapter 14

- 14.1 (a) 2.15 ms, 2930 rad/s (b) 2.00×10^4 Hz, 1.26×10^5 rad/s (c) 1.3×10^{-15} s $\leq T \leq 2.3 \times 10^{-15}$ s, 4.3×10^{14} Hz $\leq f \leq 7.5 \times 10^{14}$ Hz (d) 2.0×10^{-7} s, 3.1×10^7 rad/s
 14.3 5530 rad/s, 1.14 ms
 14.5 0.0625 s
 14.7 (a) 0.80 s (b) 1.25 Hz (c) 7.85 rad/s (d) 3.0 cm (e) 148 N/m
 14.9 (a) 0.167 s (b) 37.7 rad/s (c) 0.0844 kg
 14.11 (a) 0.150 s (b) 0.0750 s
 14.13 (a) 0.98 m (b) $\pi/2$ rad (c) $x = (-0.98 \text{ m}) \sin[(12.2 \text{ rad/s})t]$
 14.15 (a) T/6 (b) T/12 (c) no
 14.17 120 kg

- 14.19 (a) 0.253 kg (b) 1.21 cm (c) 3.03 N
 14.21 (a) 1.51 s (b) 26.0 N/m (c) 30.8 cm/s
 (d) 1.92 N (e) -0.0125 m , 30.4 cm/s , 0.216 m/s^2
 (f) 0.324 N
 14.23 (a) $\omega^2 A \sin \omega t$ (b) 0 (c) 0
 (d) $\pm A$ (e) 1.3 s
 14.25 92.2 m/s^2
 14.27 (a) 0.0336 J (b) 0.0150 m (c) 0.669 m/s
 14.29 (a) 1.20 m/s (b) 1.11 m/s (c) 36 m/s^2
 (d) 13.5 m/s^2 (e) 0.36 J
 14.31 $\frac{\sqrt{3}}{2}A$, larger
 14.33 0.240 m
 14.35 (a) 0.376 m (b) 59.3 m/s^2 (c) 119 N
 14.37 (a) 4.06 cm (b) 1.21 m/s (c) 29.8 rad/s
 14.39 (a) 0, 0, 3.92 J, 3.92 J (b) 3.92 J, 0, 0, 3.92 J
 (c) 0.98 J, 0.98 J, 1.96 J, 3.92 J
 14.41 (a) $2.7 \times 10^{-8} \text{ kg} \cdot \text{m}^2$ (b) $4.3 \times 10^{-6} \text{ N} \cdot \text{m/rad}$
 14.43 (a) 0.25 s (b) 0.25 s
 14.45 0.407 swings per second
 14.47 10.7 m/s^2
 14.49 (a) $g \sin \theta$ (b) $g \sqrt{\cos^2(\theta/2) + 4[\cos(\theta/2) - \cos \theta]^2}$
 (c) $2g(1 - \cos \theta)$, $\sqrt{2gL(1 - \cos \theta)}$
 14.51 $A: 2\pi\sqrt{\frac{L}{g}}$, $B: \frac{2\sqrt{2}}{3}\left(2\pi\sqrt{\frac{L}{g}}\right)$; pendulum A
 14.53 $A: 2\pi\sqrt{\frac{L}{g}}$, $B: \sqrt{\frac{11}{10}}\left(2\pi\sqrt{\frac{L}{g}}\right)$, pendulum B
 14.55 $\frac{\sqrt{3}}{2}\omega$
 14.57 (a) 0.393 Hz (b) 1.73 kg/s
 14.59 (a) $A_1/3$ (b) $2A_1$
 14.61 (a) 12 (b) 72
 14.63 0.353 m
 14.65 (a) $16k_1$ (b) double it
 14.67 599 N/m
 14.69 (a) 24.4 cm (b) 0.221 s
 (c) 1.19 m/s
 14.71 2.00 m
 14.73 $0.921\left(\frac{1}{2\pi}\sqrt{\frac{g}{L}}\right)$
 14.75 (a) 0.784 s (b) $-1.12 \times 10^{-4} \text{ s per s}$; shorter
 (c) 0.419 s
 14.77 (a) 0.150 m/s (b) 0.112 m/s^2 downward
 (c) 0.700 s (d) 4.38 m
 14.79 (a) 2.6 m/s (b) 0.21 m (c) 0.49 s
 14.81 1.17 s
 14.83 0.421 s
 14.85 0.705 Hz, 14.5°
 14.87 $2\pi\sqrt{\frac{M}{3k}}$
 14.89 (a) 1.60 s (b) 0.625 HZ (c) 3.93 rad/s
 (d) 5.1 cm; 0.4 s, 1.2 s, 1.8 s (e) 79 cm/s^2 ; 0.4 s,
 1.2 s, 1.8 s (f) 4.9 kg
 14.91 (b) The angular amplitude increases as L
 decreases. (c) about 53°
 14.93 (a) $Mv^2/6$ (c) $\omega = \sqrt{3k/M}$, $M' = M/3$
 14.95 Choice (a)

Chapter 15

- 15.1 (a) 0.439 m, 1.28 ms (b) 0.219 m
 15.3 $220 \text{ m/s} = 800 \text{ km/h}$
 15.5 (a) 1.7 cm to 17 m (b) $4.3 \times 10^{14} \text{ Hz}$ to
 $7.5 \times 10^{14} \text{ Hz}$ (c) 1.5 cm (d) 6.4 cm
 15.7 (a) 25.0 Hz, 0.0400 s, 19.6 rad/m
 (b) $y(x, t) = (0.0700 \text{ m}) \cos[(19.6 \text{ m}^{-1})x +$
 $(157 \text{ rad/s})t]$ (c) 4.95 cm (d) 0.0050 s
 15.9 (a) yes (b) yes (c) no (d) $v_y =$
 $\omega A \cos(kx + \omega t)$, $a_y = -\omega^2 A \sin(kx + \omega t)$
 15.11 (a) 4 mm (b) 0.040 s (c) 0.14 m, 3.6 m/s
 (d) 0.24 m, 6.0 m/s (e) no

- 15.13 (b) $+x$ -direction
 15.15 (a) 17.5 m/s (b) 0.146 m (c) both would increase
 by a factor of $\sqrt{2}$
 15.17 0.337 kg
 15.19 (a) 9.53 N (b) 20.8 m/s
 15.23 4.10 mm
 15.25 (a) 95 km (b) $0.25 \mu\text{W/m}^2$ (c) 110 kW
 15.27 (a) 0.050 W/m^2 (b) 22 kJ
 15.35 (a) $(1.33 \text{ m})n$, $n = 0, 1, 2, \dots$
 (b) $(1.33 \text{ m})(n + \frac{1}{2})$, $n = 0, 1, 2, \dots$
 15.37 (a) 96.0 m/s (b) 461 N (c) 1.13 m/s, 4.26 m/s^2
 15.39 (b) 2.80 cm (c) 277 cm (d) 185 cm, 7.96 Hz,
 0.126 s, 1470 cm/s (e) 280 cm/s
 $y(x, t) = (5.60 \text{ cm}) \times$
 $\sin[(0.0906 \text{ rad/cm})x] \sin[(133 \text{ rad/s})t]$
 15.41 (a) 0.240 m (b) 0.0600 m
 15.43 (a) 45.0 cm (b) no
 15.45 $1.80 \times 10^3 \text{ m/s}$, $1.06 \times 10^{-4} \text{ m}$
 15.47 (a) 20.0 Hz, 126 rad/s, 3.49 rad/m
 (b) $y(x, t) = (2.50 \times 10^{-3} \text{ m}) \cos[(3.49 \text{ rad/m})x$
 $-(126 \text{ rad/s})t]$
 (c) $y(0, t) = (2.50 \times 10^{-3} \text{ m}) \cos[(126 \text{ rad/s})t]$
 (d) $y(1.35 \text{ m}, t) = (2.50 \times 10^{-3} \text{ m}) \times$
 $\cos[(126 \text{ rad/s})t - 3\pi/2 \text{ rad}]$
 (e) 0.315 m/s (f) $-2.50 \times 10^{-3} \text{ m}$, 0
 15.49 $6.00 \times 10^{-2} \text{ kg}$
 15.51 (a) 62.1 m
 15.53 13.7 Hz, 25.0 m
 15.55 1.83 m
 15.57 360 Hz (copper), 488 Hz (aluminum)
 15.59 (a) 18.8 cm (b) 0.0169 kg
 15.61 (a) 7.07 cm (b) 0.400 kW
 15.63 (a) 0.0673 kg/m (b) 14.3 N (c) 14.6 m/s
 (d) 19.5 Hz (e) 1, 26, 51, 76, 101
 (f) 0.0615 m (g) $(n-1)(15.0 \text{ mm})$
 (h) 3.64 m/s
 15.65 (a) 2.22 g (b) $2.24 \times 10^4 \text{ m/s}^2$
 15.67 233 N
 15.69 1780 kg/m^3
 15.71 (a) 148 N (b) 26%
 15.73 (c) 47.5 Hz (d) 138 g
 15.75 (a) because $P_{\text{av}} = 2\pi^2\sqrt{\mu FA^2 f^2}$
 (b) 280 m/s (c) 1100 rad/s
 15.77 (a) 392 N (b) 392 N + $(7.70 \text{ N/m})x$
 (c) 3.89 s
 15.79 Choice (b)

Chapter 16

- 16.1 (a) 0.344 m (b) $1.2 \times 10^{-5} \text{ m}$
 (c) 6.9 m, 50 Hz
 16.3 (a) 7.78 Pa (b) 77.8 Pa (c) 778 Pa
 16.5 0.156 s
 16.7 90.8 m
 16.9 81.4°C
 16.11 (a) $5.5 \times 10^{-15} \text{ J}$ (b) 0.074 mm/s
 16.13 15.0 cm
 16.15 (a) 4.14 Pa (b) 0.0208 W/m² (c) 103 dB
 16.17 (a) $4.4 \times 10^{-12} \text{ W/m}^2$ (b) 6.4 dB
 (c) $5.8 \times 10^{-11} \text{ m}$
 16.19 14.0 dB
 16.21 (a) $2.0 \times 10^{-7} \text{ W/m}^2$ (b) 6.0 m (c) 290 m
 (d) yes, no
 16.23 (a) fundamental: 0.60 m; 0, 1.20 m; first overtone:
 0.30 m, 0.90 m; 0, 0.60 m, 1.20 m; second overtone:
 0.20 m, 0.60 m, 1.00 m; 0, 0.40 m, 0.80 m, 1.20 m
 (b) fundamental: 0; 1.20 m; first overtone: 0, 0.80 m;
 0.40 m, 1.20 m; second overtone: 0, 0.48 m, 0.96 m;
 0.24 m, 0.72 m, 1.20 m
 16.25 506 Hz, 1517 Hz, 2529 Hz
 16.27 (a) 35.2 Hz (b) 17 Hz
 16.29 (a) closed (b) 5th harmonic (c) 0.453 m
 (d) 190 Hz

- 16.31 (a) 614 Hz (b) 1230 Hz
 16.33 (a) 172 Hz (b) 86 Hz
 16.35 0.125 m
 16.37 (a) $(820 \text{ Hz})n$, $n = 1, 2, 3, \dots$
 (b) $(410 \text{ Hz})(2n + 1)$, $n = 0, 1, 2, \dots$
 16.39 (a) 433 Hz (b) loosen
 16.41 780 m/s
 16.43 (a) 375 Hz (b) 371 Hz (c) 4 Hz
 16.45 (a) 0.25 m/s (b) 0.91 m
 16.47 19.8 m/s
 16.49 (a) 355 m/s (b) 0.710 m (c) 500 Hz
 16.51 (a) 7.02 m/s, toward (b) 1404 Hz
 16.53 (a) 36.0° (b) 2.94 s
 16.55 (b) 344 m/s (c) no
 16.57 (a) 0.879 s (b) 27.3 Hz
 16.59 (a) stopped (b) 7th and 9th (c) 0.439 m
 16.61 (a) 0.026 m, 0.53 m, 1.27 m, 2.71 m, 9.01 m
 (b) 0.26 m, 0.86 m, 1.84 m, 4.34 m (c) 86 Hz
 16.63 (a) 0.0823 m (b) 120 Hz
 16.67 (b) 2.0 m/s
 16.69 (a) 38 Hz (b) no
 16.71 (a) 375 m/s (b) 1.39 (c) 0.8 cm
 16.73 (a) 441 Hz (b) 0.430 m
 16.75 (d) 9.69 cm/s, 667 m/s²
 16.77 Choice (b)
 16.79 Choice (a)
 16.81 Choice (b)

Chapter 17

- 17.1 (a) -81.0°F (b) 134.1°F (c) 88.0°F
 17.3 (a) 27.2°C (b) -55.6°C
 17.5 (a) -18.0°F (b) -10.0°C
 17.7 0.964 atm
 17.9 (a) -282°C (b) no, 47,600 Pa
 17.11 0.39 m
 17.13 1.9014 cm; 1.8964 cm
 17.15 49.4°C
 17.17 $1.7 \times 10^{-5} (\text{C}^\circ)^{-1}$
 17.19 (a) 1.431 cm^2 (b) 1.436 cm^2
 17.21 (a) 6.0 mm (b) $-1.0 \times 10^8 \text{ Pa}$
 17.23 $\frac{L_{\text{Invar}}}{L_{\text{Al}}} = \frac{1}{0.225} = 4$
 17.25 554 kJ
 17.27 23 min
 17.29 240 J/kg $\cdot$ K
 17.31 0.526°C
 17.33 0.0613°C
 17.35 (a) 215 J/kg $\cdot$ K (b) water (c) too small
 17.37 0.114 kg
 17.39 27.5°C
 17.41 150°C
 17.43 54.5 kJ, 13.0 kcal, 51.7 Btu
 17.45 357 m/s
 17.47 3.45 L
 17.49 $5.05 \times 10^{15} \text{ kg}$
 17.51 0.0674 kg
 17.53 2370 J/kg $\cdot$ K
 17.55 157 W/m $\cdot$ K
 17.57 (a) -0.86°C (b) 24 W/m^2
 17.61 105.5°C
 17.63 (a) 21 kW (b) 6.4 kW
 17.65 2.1 cm^2
 17.67 (a) $1.61 \times 10^{11} \text{ m}$ (b) $5.43 \times 10^6 \text{ m}$
 17.69 35.0°C
 17.71 $c = \frac{Q\beta}{\rho\Delta V}$
 17.73 (a) 35.1°M (b) 39.6°C
 17.75 69.4°C
 17.77 23.0 cm (first rod), 7.0 cm (second rod)
 17.79 (b) $1.9 \times 10^8 \text{ Pa}$
 17.81 (a) 87°C (b) -80°C
 17.83 $2.00 \times 10^3 \text{ J/kg} \cdot \text{K}$

A-14 Answers to Odd-Numbered Problems

- 17.85 (a) 83.6 J (b) 1.86 J/mol · K (c) 5.60 J/mol · K
 17.87 (a) 4.20×10^7 J (b) 10.7 C° (c) 30.0 C°
 17.89 (a) 0.60 kg (b) 0.80 bottles/h
 17.91 3.4×10^5 J/kg
 17.93 (a) no (b) 0.0°C, 0.156 kg
 17.95 (a) 86.1°C (b) no ice, 0.130 kg liquid water, no steam
 17.97 (a) 100°C (b) 0.0214 kg steam, 0.219 kg liquid water
 17.99 (a) 93.9 W (b) 1.35
 17.101 2.9
 17.103 (a) 1.04 kW (b) 87.1 W (c) 1.13 kW (d) 28 g (e) 1.1 bottles
 17.105 (c) 170 h (d) 1.5×10^{10} s $\approx$ 500 y, no
 17.107 5.82 g
 17.109 (a) 3.00×10^4 J/kg (b) 1.00×10^3 J/kg · K (liquid), 1.33×10^3 J/kg · K (solid)
 17.111 A: 216 W/m · K, B: 130 W/m · K
 17.113 $H = \frac{k\pi R_1 R_2}{L}(T_H - T_C)$
 17.115 (a) $H = \frac{(T_2 - T_1)2\pi kL}{\ln(b/a)}$
 (b) $T = T_2 - \frac{(T_2 - T_1)\ln(r/a)}{\ln(b/a)}$ (d) 73°C
 (e) 49 W
 17.117 Choice (a)
 17.119 Choice (a)

Chapter 18

- 18.1 (a) 0.122 mol (b) 14,700 Pa, 0.145 atm
 18.3 0.100 atm
 18.5 (a) 5.7×10^6 Pa (b) 3.79×10^6 Pa, -25.2%
 18.7 503°C
 18.9 19.7 kPa
 18.11 0.159 L
 18.13 0.0508 V
 18.15 0.421 mol
 18.17 (a) 6.95×10^{-16} kg (b) 2.32×10^{-13} kg/m³
 18.19 55.6 mol, 3.35×10^{25} molecules
 18.21 (a) 2.20×10^6 molecules (b) 2.44×10^{19} molecules
 18.23 (a) 3.45×10^{-9} m (b) about 10 times the diameter of a typical molecule (c) 10 times the spacing of atoms in solids
 18.25 (a) 5.83×10^7 J (b) 242 m/s
 18.27 6.46×10^{23}
 18.29 (a) 1.93×10^6 m/s, no (b) 7.3×10^{10} K
 18.31 (a) 6.21×10^{-21} J (b) 2.34×10^5 m²/s² (c) 484 m/s (d) 2.57×10^{-23} kg · m/s (e) 1.24×10^{-19} N (f) 1.24×10^{-17} Pa (g) 8.17×10^{21} molecules (h) 2.45×10^{22} molecules
 18.33 3800°C
 18.35 1100 m/s
 18.37 (a) 1870 J (b) 1120 J
 18.39 (a) 741 J/kg · K, $c_w = 5.65c_{N_2}$ (b) 5.65 kg; 4850 L
 18.41 (a) 337 m/s (b) 380 m/s (c) 412 m/s
 18.43 (a) 1.34 (b) 1.23
 18.45 (a) 610 Pa (b) 22.12 MPa
 18.49 (a) 11.8 kPa (b) 0.566 L
 18.51 272°C
 18.53 0.195 kg
 18.55 (a) $a = \left(\frac{T_{in}}{T_{out}} - 1\right)g$ (b) $T_{in} = T_{out}\left(\frac{a}{g} + 1\right)$
 18.57 (a) 0.0959 m³ (b) 17.8 mol/s (c) 42.6 m
 18.59 (a) 30.7 cylinders (b) 8420 N (c) 7800 N
 18.61 (a) 26.2 m/s (b) 16.1 m/s, 5.44 m/s (c) 1.74 m
 18.63 $\approx 5 \times 10^{27}$ atoms
 18.65 (a) 4.65×10^{-26} kg (b) 6.11×10^{-21} J (c) 2.04×10^{24} molecules (d) 12.5 kJ

- 18.67 (b) r_2 (c) $r_1 = \frac{R_0}{2^{1/6}}$, $r_2 = R_0$, $2^{-1/6}$ (d) U_0
 18.69 (a) $2R = 16.6$ J/mol · K (b) less than
 18.71 (b) 1.40×10^5 K (N₂), 1.01×10^4 K (H₂) (c) 6370 K (N₂), 459 K (H₂)
 18.73 $3kT/m$, same
 18.75 (b) 0.0421 N (c) $(2.94 \times 10^{-21})N$ (d) 0.0297 N, $(2.08 \times 10^{-21})N$ (e) 0.0595 N, $(4.15 \times 10^{-21})N$
 18.77 (a) $p_0 + \frac{mg}{\pi r^2}$ (b) $-\left(\frac{y}{h}\right)(p_0\pi r^2 + mg)$
 (c) $\frac{1}{2\pi}\sqrt{\frac{g}{h}\left(1 + \frac{p_0\pi r^2}{mg}\right)}$, no
 18.79 (a) 42.6% (b) 3 km (c) 1 km
 18.81 (a) 4.1×10^{-46} kg · m² (c) $v_f = 0$, $\omega = v_i/r$ (d) 5.5×10^{12} rad/s
 18.83 (a) 4.5×10^{11} m (b) 703 m/s, 6.4×10^8 s (≈ 20 y) (c) 1.4×10^{-14} Pa (d) 650 m/s, $v_H > v_{esc}$, evaporate (f) 2×10^5 K, $> 3T_{sun}$, no
 18.85 Choice (a)
 18.87 Choice (c)

Chapter 19

- 19.1 (b) 1330 J
 19.3 (b) -6180 J
 19.5 (a) 1.04 atm
 19.7 (a) $(p_1 - p_2)(V_2 - V_1)$ (b) negative of work done in reverse direction
 19.9 (a) 34.7 kJ (b) 80.4 kJ (c) no
 19.11 (a) 278 K, at a (b) 0; 162 J (c) 53 J
 19.13 (a) $T_a = 535$ K, $T_b = 9350$ K, $T_c = 15,000$ K (b) 21 kJ done by gas (c) 36 kJ
 19.15 (a) $Q = +100$ J, $W = -300$ J (b) $Q = -100$ J, $W = -300$ J (c) $Q = +100$ J, $W = +300$ J (d) -100 J, $W = +300$ J
 19.17 (b) 208 J (c) on the piston (d) 712 J (e) 920 J (f) 208 J
 19.19 (a) 948 K (b) 900 K
 19.21 $\frac{2}{5}$
 19.23 (a) 747 J (b) 1.30
 19.25 (a) -605 J (b) 0 (c) yes, 605 J, liberate
 19.27 (a) 476 kPa (b) -10.6 kJ (c) 1.59, heated
 19.29 (b) 314 J (c) -314 J
 19.31 11.6°C
 19.33 $4\sqrt{2}$
 19.35 (a) increase (b) 4.8 kJ
 19.37 (a) 0.681 mol (b) 0.0333 m³ (c) 2.23 KJ (d) 0
 19.39 (a) 45.0 J (b) liberate, 65.0 J (c) 23.0 J, 22.0 J
 19.41 (a) the same (b) 4.0 kJ, absorb (c) 8.0 kJ
 19.43 (a) 0.80 L (b) 305 K, 1220 K, 1220 K (c) ab : 76 J, into the gas
 ca : -107 J, out of the gas
 bc : 56 J, into the gas
 (d) ab : 76 J, increased
 bc : 0, no change
 ca : -76 J, decreased
 19.45 (b) -300 J, out of gas
 19.47 (b) 6.00 L, 2.5×10^4 Pa, 75.0 K (c) 95 J (d) heat it at constant volume
 19.49 (b) 11.9 C°
 19.51 (a) 0.169 m (b) 198°C (c) 71.3 kJ
 19.53 (a) $Q = 450$ J, $\Delta U = 0$ (b) $Q = 0$, $\Delta U = -450$ J (c) $Q = 1125$ J, $\Delta U = 675$ J
 19.55 (a) $W = 738$ J, $Q = 2590$ J, $\Delta U = 1850$ J (b) $W = 0$, $Q = -1850$ J, $\Delta U = -1850$ J (c) $\Delta U = 0$
 19.57 (a) $W = -187$ J, $Q = -654$ J, $\Delta U = -467$ J (b) $W = 113$ J, $Q = 0$, $\Delta U = -113$ J (c) $W = 0$, $Q = 580$ J, $\Delta U = 580$ J

- 19.59 (a) a : adiabatic, b : isochoric, c : isobaric (b) 28.0°C (c) a : -30.0 J, a : 0, a : 20.0 J (d) a (e) a : decrease, b : stay the same, c : increase
 19.61 (a) 53.5 J + $v^{0.40}[(21.4 \text{ J})\ln v - 53.5 \text{ J}]$ (b) 7 (c) 7 (d) 7 (e) 63.1 J
 19.63 Choice (c)
 19.65 Choice (d)

Chapter 20

- 20.1 (a) 6500 J (b) 34%
 20.3 (a) 23% (b) 12,400 J (c) 0.350 g (d) 222 kW = 298 hp
 20.5 (a) 12.3 atm (b) 5470 J, ca (c) 3723 J, bc (d) 1747 J (e) 31.9%
 20.7 (a) 12 (b) 7.4 kJ, 20 kJ
 20.9 (a) 58% (b) 1.4%
 20.11 1.2 h
 20.13 576 K, 504 K
 20.15 (a) 215 J (b) 378 K (c) 39.0%
 20.17 (a) 38 kJ (b) 590°C
 20.19 (a) 492 J (b) 212 W (c) 5.4
 20.21 161 K
 20.23 (a) 429 J/K (b) -393 J/K (c) 36 J/K
 20.25 -6.31 J/K
 20.27 (a) 6.05×10^3 J/K (b) about five times greater for vaporization
 20.29 10.0 J/K
 20.31 (a) 121 J (b) 3800 cycles
 20.33 (a) 90.2 J (b) 320 J (c) 45°C (d) 0 (e) 263 g
 20.35 -5.8 J/K, decrease
 20.37 (b) absorbed: bc ; rejected: ab and ca (c) $T_a = T_b = 241$ K, $T_c = 481$ K (d) 610 J, 610 J (e) 8.7%
 20.39 (a) 21.0 kJ (enters), 16.6 kJ (leaves) (b) 4.4 kJ, 21% (c) $e = 0.31e_{max}$
 20.41 (a) 7.0% (b) 3.0 MW; 2.8 MW (c) 6×10^5 kg/h = 6×10^5 L/h
 20.43 (a) 1: 2.00 atm, 4.00 L; 2: 2.00 atm, 6.00 L; 3: 1.11 atm, 6.00 L; 4: 1.67 atm, 4.00 L (b) $1 \rightarrow 2$: 1422 J, 405 J; $2 \rightarrow 3$: -1355 J, 0; $3 \rightarrow 4$: -274 J, -274 J; $4 \rightarrow 1$: 339 J, 0 (c) 131 J (d) 7.44%; $e = 0.168e_C$
 20.45 $1 - T_C/T_H$, same
 20.47 (a) 122 J, -78 J (b) 5.10×10^{-4} m³ (c) b : 2.32 MPa, 4.81×10^{-5} m³, 771 K (c): 4.01 MPa, 4.81×10^{-5} m³, 1332 K (d): 0.147 MPa, 5.10×10^{-4} m³, 518 K (d) 61.1%, 77.5%
 20.49 6.23
 20.53 (a) A: 28.9%, B: 38.3%, C: 53.8%, D: 24.4% (b) C (c) B > D > A
 20.55 (a) 4.83% (b) 4.83% (c) 6.25% (d) $e = \frac{0.80T_d - 200}{12T_d - 2700}$, 6.67%
 20.57 (a) 151 K (b) 305 Pa (c) 5.39 kJ (d) 4.39 kJ
 20.59 Choice (b)
 20.61 Choice (d)

Chapter 21

- 21.1 (a) 2.00×10^{10} (b) 8.59×10^{-13}
 21.3 3.4×10^{18} m/s² (proton), 6.3×10^{21} m/s² (electron)
 21.5 1.3 nC
 21.7 3.7 km
 21.9 (a) 0.742 μ C (b) 0.371 μ C, 1.48 μ C
 21.11 (a) 2.21×10^4 m/s²
 21.13 +0.750 nC
 21.15 1.8×10^{-4} N, in the + x -direction

- 21.17 $2.58 \mu\text{N}$, in the $-y$ -direction
 21.19 Attractive
 21.21 (a) $4.40 \times 10^{-16} \text{ N}$ (b) $2.63 \times 10^{11} \text{ m/s}^2$
 (c) $2.63 \times 10^5 \text{ m/s}$
 21.23 (a) $3.30 \times 10^6 \text{ N/C}$, to the left (b) 14.2 ns
 (c) $1.80 \times 10^3 \text{ N/C}$, to the right
 21.25 (a) $-21.9 \mu\text{C}$ (b) $1.02 \times 10^{-7} \text{ N/C}$
 21.27 (a) 364 N/C
 (b) no; $2.73 \mu\text{m}$, downward
 21.29 (a) $-\hat{j}$ (b) $\frac{\sqrt{2}}{2}\hat{i} + \frac{\sqrt{2}}{2}\hat{j}$ (c) $-0.39\hat{i} + 0.92\hat{j}$
 21.31 $3.09 \times 10^{-3} \text{ N}$ in the $+y$ -direction
 21.33 20.0 cm
 21.35 (a) (i) 574 N/C , $+x$ -direction
 (ii) 268 N/C , $-x$ -direction
 (iii) 404 N/C , $-x$ -direction
 (b) (i) $9.20 \times 10^{-17} \text{ N}$, $-x$ -direction
 (ii) $4.30 \times 10^{-17} \text{ N}$, $+x$ -direction
 (iii) $6.48 \times 10^{-17} \text{ N}$, $+x$ -direction
 21.37 $1.04 \times 10^7 \text{ N/C}$, toward the $-2.00\text{-}\mu\text{C}$ charge
 21.39 (a) 8740 N/C , to the right (b) 6540 N/C , to the right
 (c) $1.40 \times 10^{-15} \text{ N}$, to the right
 21.41 $1.73 \times 10^{-8} \text{ N}$, toward the point midway between the electrons
 21.43 (a) $E_x = E_y = E = 0$
 (b) $E_x = 2660 \text{ N/C}$, $E_y = 0$, $E = 2660 \text{ N/C}$, $+x$ -direction
 (c) $E_x = 129 \text{ N/C}$, $E_y = -510 \text{ N/C}$, $E = 526 \text{ N/C}$, 284° counterclockwise from the $+x$ -axis
 (d) $E_x = 0$, $E_y = 1380 \text{ N/C}$, $E = 1380 \text{ N/C}$, $+y$ -direction
 21.45 113 N/C , $-x$ direction
 21.47 (a) 1340 N/C , $+x$ -direction
 (b) 1340 N/C , $-x$ -direction
 (c) 8260 N/C , $+x$ -direction
 (d) 8260 N/C , $-x$ -direction
 21.49 (a) $(7.0 \text{ N/C})\hat{i}$ (b) $(1.75 \times 10^{-5} \text{ N})\hat{i}$
 21.51 (a) $1.4 \times 10^{-11} \text{ C} \cdot \text{m}$, from q_1 toward q_2
 (b) 860 N/C
 21.53 (a) $\vec{p}$ aligned in the same or the opposite direction as $\vec{E}$
 (b) stable: $\vec{p}$ aligned in the same direction as $\vec{E}$; unstable: $\vec{p}$ aligned in the opposite direction
 21.55 (a) 1680 N , from the $+5.00\text{-}\mu\text{C}$ charge toward the $-5.00\text{-}\mu\text{C}$ charge
 (b) $22.3 \text{ N} \cdot \text{m}$, clockwise
 21.57 (b) $\frac{Q^2}{8\pi\epsilon_0 L^2}(1 + 2\sqrt{2})$, away from the center of the square
 21.59 (a) $8.63 \times 10^{-5} \text{ N}$, $-5.52 \times 10^{-5} \text{ N}$
 (b) $1.02 \times 10^{-4} \text{ N}$, 32.6° below the $+x$ -axis
 21.61 (b) $2.80 \mu\text{C}$ (c) 39.5°
 21.63 $3.41 \times 10^4 \text{ N/C}$, to the left
 21.65 (a) 1 nC (b) 6×10^{10}
 21.67 between the charges, 0.24 m from the 0.500-nC charge
 21.69 at $x = d/3$, $q = -4Q/9$
 21.71 (a) $\frac{6q^2}{4\pi\epsilon_0 L^2}$, away from the vacant corner
 (b) $\frac{3q^2}{4\pi\epsilon_0 L^2}\left(\sqrt{2} + \frac{1}{2}\right)$, toward the center of the square
 21.73 (a) 6.0×10^{23}
 (b) $4.1 \times 10^{-31} \text{ N}$ (gravitational), 510 kN (electric)
 (c) yes (electric), no (gravitational)
 21.75 2190 km/s
 21.77 (a) $\frac{mv_0^2 \sin^2 \alpha}{2eE}$ (b) $\frac{mv_0^2 \sin^2 \alpha}{eE}$
 (d) h_{max} : 0.418 m , d : 2.89 m

- 21.79 (a) $E_x = \frac{Q}{4\pi\epsilon_0 a}\left(\frac{1}{x-a} - \frac{1}{x}\right)$, $E_y = 0$
 (b) $\frac{qQ}{4\pi\epsilon_0 a}\left(\frac{1}{r} - \frac{1}{r+a}\right)\hat{i}$
 21.81 (a) 1.56 N/C , $+x$ -direction (c) smaller
 (d) 4.7%
 21.83 $E_x = E_y = \frac{Q}{2\pi^2\epsilon_0 a^2}$
 21.85 (a) $6.25 \times 10^4 \text{ N/C}$, 225° counterclockwise from an axis pointing to the right at point P
 (b) $1.00 \times 10^{-14} \text{ N}$, opposite the electric field direction
 21.87 (a) $\pi(R_2^2 - R_1^2)\sigma$
 (b) $\frac{\sigma}{2\epsilon_0}\left(\frac{1}{\sqrt{(R_1/x)^2 + 1}} - \frac{1}{\sqrt{(R_2/x)^2 + 1}}\right)\frac{|x|}{x}\hat{i}$
 (c) $\frac{\sigma}{2\epsilon_0}\left(\frac{1}{R_1} - \frac{1}{R_2}\right)x\hat{i}$; $x \ll R_1$
 (d) $\frac{1}{2\pi}\sqrt{\frac{q\sigma}{2\epsilon_0 m}\left(\frac{1}{R_1} - \frac{1}{R_2}\right)}$
 21.89 (a) $q_1 = 8.00 \mu\text{C}$, $q_2 = 3.00 \mu\text{C}$
 (b) 7.50 N , in the $-x$ -direction (c) $x = 0.248 \text{ m}$
 21.91 (a) $\frac{\sigma}{2\epsilon_0}\left[\frac{1}{\sqrt{1 + (R/\Delta)^2}}\right]$ downward (b) $q\sigma/2g\epsilon_0$
 (c) $-\frac{q\sigma}{2\epsilon_0}(\sqrt{\Delta^2 + R^2} - R)$ (d) $mg\Delta$
 (e) $\frac{2R\alpha}{1 - \alpha^2}$ (f) $\sqrt{2g\left(\Delta - \frac{\sqrt{\Delta^2 + R^2} - R}{\alpha}\right)}$
 (g) 57.7 g , 10.7 cm
 21.93 (b) $-\frac{\lambda}{4\pi\epsilon_0 a}\left[\frac{y_2 - L}{\sqrt{(y_2 - L)^2 + a^2}} - \frac{y_2}{\sqrt{y_2^2 + a^2}}\right]\lambda dy_2$
 (c) $\frac{Q^2}{2\pi\epsilon_0 L^2}(\sqrt{(L/a)^2 + 1} - 1)\hat{i}$
 (e) $\frac{Q^2}{2\pi\epsilon_0 L^2}\left[L \ln\left(\frac{L + \sqrt{a^2 + L^2}}{a}\right) + a - \sqrt{a^2 + L^2}\right]$
 (f) 21.5 m/s
 21.95 (b) $q_1 < 0$, $q_2 > 0$ (c) $0.843 \mu\text{C}$ (d) 56.2 N
 21.97 (a) $\frac{Q}{2\pi\epsilon_0 L}\left(\frac{1}{2x+a} - \frac{1}{2L+2x+a}\right)$
 21.99 Choice (c)
 21.101 Choice (b)

Chapter 22

- 22.1 (a) $1.8 \text{ N} \cdot \text{m}^2/\text{C}$ (b) no (c) (i) 0° (ii) 90°
 22.3 (a) $3.53 \times 10^5 \text{ N} \cdot \text{m}^2/\text{C}$ (b) $3.13 \mu\text{C}$
 22.5 $7.70 \times 10^3 \text{ N/C}$
 22.7 14.8 cm^2
 22.9 (a) 0 (b) $1.22 \times 10^8 \text{ N/C}$, radially inward
 (c) $3.64 \times 10^7 \text{ N/C}$, radially inward
 22.11 $0.977 \text{ N} \cdot \text{m}^2/\text{C}$, inward
 22.13 0.0810 N
 22.15 (a) 350 N/C (b) 700 N/C (c) 1400 N/C
 22.17 (a) $6.47 \times 10^5 \text{ N/C}$, $+y$ -direction
 (b) $7.2 \times 10^4 \text{ N/C}$, $-y$ -direction
 22.19 (a) $5.73 \mu\text{C}/\text{m}^2$ (b) $6.47 \times 10^5 \text{ N/C}$
 (c) $-5.65 \times 10^4 \text{ N} \cdot \text{m}^2/\text{C}$
 22.21 (a) $0.260 \mu\text{C}/\text{m}^3$ (b) 1960 N/C
 22.23 (a) $6.56 \times 10^{-21} \text{ J}$ (b) $1.20 \times 10^5 \text{ m/s}$
 22.25 (a) 6.00 nC (b) -1.00 nC
 22.27 (a) $2\pi R\sigma$ (b) $\frac{\sigma R}{\epsilon_0 r}$ (c) $\frac{\lambda}{2\pi\epsilon_0 r}$
 22.29 1.16 km/s
 22.31 10.2°
 22.33 (a) $750 \text{ N} \cdot \text{m}^2/\text{C}$
 (b) 0 (c) 577 N/C , $+x$ -direction
 (d) within and outside
 22.35 (a) $\frac{Q}{4\pi\epsilon_0 r^2}$, toward the center of the shell (b) 0

- 22.37 (a) $\frac{\lambda}{2\pi\epsilon_0 r}$, radially outward
 (b) $\frac{\lambda}{2\pi\epsilon_0 r}$, radially outward
 (d) $-\lambda$ (inner), $+\lambda$ (outer)
 22.39 (a) $\frac{\rho r}{2\epsilon_0}$ (b) $\frac{\lambda}{2\pi\epsilon_0 r}$ (c) They are equal.
 22.41 (a) $r < R$: 0; $R < r < 2R$: $\frac{1}{4\pi\epsilon_0 r^2} \frac{Q}{r^2}$, radially outward; $r > 2R$: $\frac{1}{4\pi\epsilon_0 r^2} \frac{2Q}{r^2}$, radially outward
 22.43 (a) (i) 0 (ii) 0 (iii) $\frac{q}{2\pi\epsilon_0 r^2}$, radially outward
 (iv) 0 (v) $\frac{3q}{2\pi\epsilon_0 r^2}$, radially outward (b) (i) 0
 (ii) $+2q$ (iii) $-2q$ (iv) $+6q$
 22.45 (a) $\frac{\alpha}{2\epsilon_0}\left(1 - \frac{a^2}{r^2}\right)$ (b) $q = +2\pi\alpha a^2$, $E = \frac{\alpha}{2\epsilon_0}$
 22.47 (c) $\frac{Qr}{4\pi\epsilon_0 R^3}\left(4 - \frac{3r}{R}\right)$ (e) $2R/3$, $\frac{Q}{3\pi\epsilon_0 R^2}$
 22.49 $\frac{\alpha r}{\epsilon_0}\left(\frac{1}{2} - \frac{r}{3R}\right)$, $\frac{\alpha R^2}{6\epsilon_0 r}$, yes
 22.51 $|x| > d$ (outside the slab): $\frac{\rho_0 d}{3\epsilon_0} \frac{x}{|x|}\hat{x}$;
 $|x| < d$ (inside the slab): $\frac{\rho_0 x^3}{3\epsilon_0 d^2}\hat{x}$
 22.53 (b) $\frac{\rho \vec{b}}{3\epsilon_0}$
 22.55 (a) uniform line of charge: A; uniformly charged sphere: B
 (b) $\lambda = 1.50 \times 10^{-7} \text{ C/m}$, $\rho = 2.81 \times 10^{-3} \text{ C/m}^3$
 22.57 (i) 377 N/C (ii) 653 N/C (iii) 274 N/C
 (iv) 0
 22.59 (a) $\frac{\pi\rho_0 R^2}{3}$ (b) $T = 2\pi\left(\frac{R_{\text{orbit}}}{R_{\text{cylinder}}}\right)\sqrt{\frac{6\epsilon_0 M}{Q\rho_0}}$
 22.61 (a) $\frac{QL}{2\epsilon_0\sqrt{(L/2)^2 + R^2}}$
 (b) $\frac{QL}{4\epsilon_0}\left(\frac{2}{L} - \frac{1}{\sqrt{R^2 + (L/2)^2}}\right)$
 (c) $\frac{QL}{4\epsilon_0}\left(\frac{2}{L} - \frac{1}{\sqrt{R^2 + (L/2)^2}}\right)$ (d) Q/ϵ_0
 22.63 Choice (a)
 22.65 Choice (b)

Chapter 23

- 23.1 -0.356 J
 23.3 $3.46 \times 10^{-13} \text{ J} = 2.16 \text{ MeV}$
 23.5 (a) 12.5 m/s (b) 0.323 m
 23.7 $1.94 \times 10^{-5} \text{ N}$
 23.9 (a) 13.6 km/s ; very long after release
 (b) $2.45 \times 10^{17} \text{ m/s}^2$; just after release
 23.11 89.3 V , point a
 23.13 7.42 m/s , faster
 23.15 (a) 0 (b) 0.750 mJ (c) -2.06 mJ
 23.17 (a) 0 (b) -175 kV (c) -0.875 J
 23.19 (a) -737 V (b) -704 V (c) $8.2 \times 10^{-8} \text{ J}$
 23.21 (a) negative (b) 171 V/m (c) -17.1 V
 23.23 (a) b (b) 800 V/m (c) $-48.0 \mu\text{J}$
 23.25 (a) 0.0539 J (b) 3.00 m/s
 23.27 (a) (i) 180 V (ii) -270 V (iii) -450 V
 (b) 719 V , inner shell
 23.29 (a) oscillatory (b) $1.67 \times 10^7 \text{ m/s}$
 23.31 150 m/s
 23.33 (a) 78.2 kV (b) 0
 23.35 0.474 J
 23.37 (a) 8.00 kV/m (b) $19.2 \mu\text{N}$ (c) $0.864 \mu\text{J}$
 (d) $-0.864 \mu\text{J}$
 23.39 -760 V

- 23.41 (a) (i) $V = \frac{q}{4\pi\epsilon_0} \left(\frac{1}{r_a} - \frac{1}{r_b} \right)$
(ii) $V = \frac{q}{4\pi\epsilon_0} \left(\frac{1}{r} - \frac{1}{r_b} \right)$ (iii) $V = 0$
(d) 0 (e) $E = \frac{q - Q}{4\pi\epsilon_0 r^2}$
- 23.43 (a) $E_x = -Ay + 2Bx$, $E_y = -Ax - C$, $E_z = 0$
(b) $x = -C/A$, $y = -2BC/A^2$, any value of z
- 23.45 (a) 0.762 nC
- 23.47 (a) $-0.360 \mu\text{J}$ (b) $x = 0.074 \text{ m}$
- 23.49 $+1.90 \times 10^5 \text{ V}$, point A
- 23.51 $4.2 \times 10^6 \text{ V}$
- 23.53 (a) $-21.5 \mu\text{J}$ (b) -2.83 kV (c) 35.4 kV/m
- 23.55 (a) $7.85 \times 10^4 \text{ V/m}^4/3$
(b) $E_x(x) = -(1.05 \times 10^5 \text{ V/m}^4/3)x^{1/3}$
(c) $3.13 \times 10^{-15} \text{ N}$, toward the anode
- 23.57 (a) $-\frac{1.46q^2}{\pi\epsilon_0 d}$
- 23.59 47.8 V
- 23.61 (a) (i) $V = (\lambda/2\pi\epsilon_0)\ln(b/a)$
(ii) $V = (\lambda/2\pi\epsilon_0)\ln(b/r)$
(iii) $V = 0$
(d) $(\lambda/2\pi\epsilon_0)\ln(b/a)$
- 23.63 (a) $1.76 \times 10^{-16} \text{ N}$, downward
(b) $1.93 \times 10^{14} \text{ m/s}^2$, downward
(c) 0.822 cm
(d) 15.3° (e) 3.29 cm
- 23.65 (a) 97.1 kV/m (b) 30.3 pC
- 23.67 $\frac{3}{5} \left(\frac{Q^2}{4\pi\epsilon_0 R} \right)$
- 23.69 360 kV
- 23.71 (a) $\frac{Q}{4\pi\epsilon_0 a} \ln \left(\frac{x+a}{x} \right)$
(b) $\frac{Q}{4\pi\epsilon_0 a} \ln \left(\frac{a + \sqrt{a^2 + y^2}}{y} \right)$
(c) (a): $\frac{Q}{4\pi\epsilon_0 x}$, (b): $\frac{Q}{4\pi\epsilon_0 y}$
- 23.73 (a) $U = \frac{1}{2}kd^2 + \frac{1}{4\pi\epsilon_0} \frac{e^2}{d}$
(b) $d_0 = \left(\frac{e^2}{4\pi\epsilon_0 k} \right)^{1/3}$ (c) $2.30 \times 10^{17} \text{ N/m}$
(d) 2.16 MeV (e) $1.44 \times 10^7 \text{ m/s}$
- 23.75 $2.48 \times 10^{-14} \text{ m}$
- 23.77 (a) $A = -6.0 \text{ V/m}^2$, $B = -4.0 \text{ V/m}^3$,
 $C = -2.0 \text{ V/m}^6$, $D = 10 \text{ V}$, $l = 2.0$, $m = 3.0$,
 $n = 6.0$
(b) (0, 0, 0): 10.0 V, 0;
(0.50 m, 0.50 m, 0.50 m):
8.0 V, 6.7 V/m;
(1.00 m, 1.00 m, 1.00 m):
-2.0 V, 21 V/m 23.79
(c) $4.79 \times 10^{-19} \text{ C}$ (drop 1),
 $1.59 \times 10^{-19} \text{ C}$ (drop 2),
 $8.09 \times 10^{-19} \text{ C}$ (drop 3),
 $3.23 \times 10^{-19} \text{ C}$ (drop 4)
(d) 3 (drop 1), 5 (drop 3), 2 (drop 4)
(e) $1.60 \times 10^{-19} \text{ C}$ (drop 1),
 $1.59 \times 10^{-19} \text{ C}$ (drop 2),
 $1.62 \times 10^{-19} \text{ C}$ (drops 3 and 4);
 $1.61 \times 10^{-19} \text{ C}$ 23.81
23.81 (a) $4.75 \times 10^{-15} \text{ C}$ (b) 94.9 kV/m
(c) $5.81 \times 10^{-15} \text{ C}$ (d) 36,300
23.83 $1.01 \times 10^{-12} \text{ m}$, $1.11 \times 10^{-13} \text{ m}$, $2.54 \times 10^{-14} \text{ m}$
23.85 Choice (b)

Chapter 24

- 24.1 (a) 10.0 kV (b) 22.6 cm^2 (c) 8.00 pF
24.3 (a) 604 V (b) 90.8 cm^2 (c) 1840 kV/m
(d) $16.3 \mu\text{C/m}^2$

- 24.5 (a) $120 \mu\text{C}$ (b) $60 \mu\text{C}$ (c) $480 \mu\text{C}$
24.7 (a) 1.05 mm (b) 84.0 V
24.9 (a) 4.35 pF (b) 2.30 V
24.11 (a) 15.0 pF (b) 3.09 cm (c) 31.2 kN/C
24.13 5.57 μF
24.15 (a) series (b) 5000
24.17 (a) $Q_1 = Q_2 = 22.4 \mu\text{C}$, $Q_3 = 44.8 \mu\text{C}$,
 $Q_4 = 67.2 \mu\text{C}$
(b) $V_1 = V_2 = 5.6 \text{ V}$, $V_3 = 11.2 \text{ V}$, $V_4 = 16.8 \text{ V}$
(c) 11.2 V
24.19 (a) $Q_1 = 156 \mu\text{C}$, $Q_2 = 260 \mu\text{C}$
(b) $V_1 = V_2 = 52.0 \text{ V}$
24.21 (a) 19.3 nF (b) 482 nC (c) 162 nC (d) 25 V
24.23 0.0283 J/m^3
24.25 (a) 90.0 pF (b) 0.0152 m³ (c) 4.5 kV
(d) 1.80 μJ
24.27 (a) $U_p = 4U_s$ (b) $Q_p = 2Q_s$ (c) $E_p = 2E_s$
24.29 (a) 24.2 μC (b) $Q_{35} = 7.7 \mu\text{C}$, $Q_{75} = 16.5 \mu\text{C}$
(c) 2.66 mJ (d) $U_{35} = 0.85 \text{ mJ}$, $U_{75} = 1.81 \text{ mJ}$
(e) 220 V
24.31 (a) 3.60 mJ (before), 13.5 mJ (after)
(b) 9.9 mJ, increase
24.33 (a) $0.620 \mu\text{C/m}^2$ (b) 1.28
24.35 $\frac{2KQ_0}{1 + K}$ increases
24.37 0.0135 m^2
24.39 (a) 6.3 μC (b) 6.3 μC (c) none
24.41 (a) 10.1 V (b) 2.25
24.43 (a) $\frac{Q}{\epsilon_0 AK}$ (b) $\frac{Qd}{\epsilon_0 AK}$ (c) $K \frac{\epsilon_0 A}{d} = KC_0$
24.45 (a) 421 J (b) 0.054 F
24.47 (a) 0.531 pF (b) 0.224 mm
24.49 (a) 0.0160 C (b) 533 V (c) 4.26 J (d) 2.14 J
24.51 (a) 158 μJ (b) 72.1 μJ
24.53 (a) 2.5 μF
(b) $Q_1 = 550 \mu\text{C}$, $Q_2 = 370 \mu\text{C}$,
 $Q_3 = Q_4 = 180 \mu\text{C}$, $Q_5 = 550 \mu\text{C}$;
 $V_1 = 65 \text{ V}$, $V_2 = 87 \text{ V}$,
 $V_3 = V_4 = 43 \text{ V}$, $V_5 = 65 \text{ V}$
24.55 $C_2 = 6.00 \mu\text{F}$, $C_3 = 4.50 \mu\text{F}$
24.57 (a) 76 μC (b) 1.4 mJ (c) 11 V (d) 1.3 mJ
24.59 (a) 2.3 μF (b) $Q_1 = 970 \mu\text{C}$, $Q_2 = 640 \mu\text{C}$
(c) 47 V
24.61 1.67 μF
24.63 (a) $\frac{Q^2}{2A\epsilon_0 K}$ (b) $\frac{d_0}{2A^2\epsilon_0 YK}$ (c) $7.53 \times 10^7 \text{ mJ/C}^2$
(d) 95.7 kV (e) 133 kV
24.65 (a) $\frac{\pi\epsilon_0 r^2 V^2}{2z^2}$ (b) $\frac{\pi\epsilon_0 r^2 V^2}{4d}$ (c) $\frac{\pi\epsilon_0 r^2 V^2}{2d}$
(d) $\frac{\pi\epsilon_0 r^2 V^2}{4d}$ (e) no
24.67 (a) $C_1 = 6.00 \mu\text{F}$, $C_2 = 3.00 \mu\text{F}$
(b) same charge; C_2 stores more energy
(c) C_1 stores more charge and energy
24.69 (a) first (connected) (b) 144 cm^2
(c) disconnected
24.71 (a) $\frac{\epsilon_0 A(K - 1)}{d \ln K}$
24.73 Choice (c)
24.75 Choice (a)

Chapter 25

- 25.1 1.0 C
25.3 (a) 3.12×10^{19} (b) $1.51 \times 10^6 \text{ A/m}^2$
(c) 0.111 mm/s (d) both (b) and
(c) would increase
25.5 (a) 0.26 mm/s (b) 0.0120 N/C
25.7 (a) 330 C (b) 41 A
25.9 9.0 μA
25.11 $4.02 \times 10^{-8} \Omega \cdot \text{m}$

- 25.13 (a) $1.06 \times 10^{-5} \Omega \cdot \text{m}$ (b) $0.00105 (\text{C}^\circ)^{-1}$
25.15 (a) 0.206 mV (b) 0.176 mV
25.17 0.457
25.19 0.125 Ω
25.21 (a) 11 A (b) 3.1 V (c) 0.28 Ω
25.23 (a) 99.54 Ω (b) 0.0158 Ω
25.25 (a) 27.4 V (b) 12.3 MJ
25.27 (a) 0 (b) 5.0 V (c) 5.0 V
25.29 3.08 V, 0.067 Ω , 1.80 Ω
25.31 (a) 1.41 A, clockwise (b) 13.7 V (c) -1.0 V
25.33 21 W, 21 W
25.35 (a) 144 Ω (b) 240 Ω
(c) 100 W: 0.833 A; 60 W: 0.500 A
25.37 (a) 29.8 W (b) 0.248 A
25.39 (a) 3.1 W (b) 7.2 W (c) 4.1 W
25.41 (a) 300 W (b) 0.90 J
25.43 (a) 2.6 MJ (b) 0.063 L (c) 1.6 h
25.45 12.3%
25.47 (a) 24.0 W (b) 4.0 W (c) 20.0 W
25.49 (a) $1.55 \times 10^{-12} \text{ s}$
25.51 (a) $3.65 \times 10^{-8} \Omega \cdot \text{m}$ (b) 172 A
(c) 2.58 mm/s
25.53 0.060 Ω
25.55 (a) 2.5 mA (b) 21.4 $\mu\text{V/m}$ (c) 85.5 $\mu\text{V/m}$
(d) 0.180 mV
25.57 (a) 80 C $^\circ$ (b) no
25.59 (a) $\frac{\rho h}{\pi r_1 r_2}$
25.61 (a) 0.20 Ω (b) 8.7 V
25.63 (a) 1.0 k Ω (b) 100 V (c) 10 W
25.65 (a) \$78.90 (b) \$140.27
25.67 (a) 171 $\mu\Omega$ (b) 176 $\mu\text{V/m}$ (c) left: 54.7 $\mu\Omega$;
right: 116 $\mu\Omega$
25.69 (a) 4.25 ppt (b) 510 pC
(c) 1.33 V (d) 1.60 ppt
25.71 6.67 V
25.73 (b) no (c) yes (d) 9.40 W (e) 4.12 W
25.75 (a) inward (b) $\frac{V}{\ln(r_{\text{outer}}/r_{\text{inner}})}$
(c) $\frac{\rho}{2\pi L} \ln(r_{\text{outer}}/r_{\text{inner}})$
(d) 616 $\Omega \cdot \text{m}$
25.77 (a) $R = \frac{\rho_0 L}{A} \left(1 - \frac{1}{e} \right)$, $I = \frac{V_0 A}{\rho_0 L \left(1 - \frac{1}{e} \right)}$
(b) $E(x) = \frac{V_0 e^{-x/L}}{L \left(1 - \frac{1}{e} \right)}$
 $V_0 \left(e^{-x/L} - \frac{1}{e} \right)$
(c) $V(x) = \frac{V_0 \left(e^{-x/L} - \frac{1}{e} \right)}{1 - \frac{1}{e}}$
25.79 Choice (c)
25.81 Choice (d)

Chapter 26

- 26.1 $3R/4$
26.3 22.5 W
26.5 (a) 3.50 A (b) 4.50 A (c) 3.15 A (d) 3.25 A
26.7 0.769 A
26.9 (a) $I = \frac{\mathcal{E}}{R}$ (b) $I = \frac{\mathcal{E}}{6R}$ (c) parallel
26.11 (a) $I_1 = 8.00 \text{ A}$, $I_3 = 12.0 \text{ A}$
(b) 84.0 V
26.13 5.00 Ω ; $I_{3,00} = 8.00 \text{ A}$, $I_{4,00} = 9.00 \text{ A}$,
 $I_{6,00} = 4.00 \text{ A}$, $I_{12,0} = 3.00 \text{ A}$
26.15 (a) $I_1 = 1.50 \text{ A}$, $I_2 = I_3 = I_4 = 0.500 \text{ A}$
(b) $P_1 = 10.1 \text{ W}$, $P_2 = P_3 = P_4 = 1.12 \text{ W}$;
bulb R_1
(c) $I_1 = 1.33 \text{ A}$, $I_2 = I_3 = 0.667 \text{ A}$
(d) $P_1 = 8.00 \text{ W}$, $P_2 = P_3 = 2.00 \text{ W}$
(e) brighter: R_2 and R_3 ; less bright: R_1

- 26.17 18.0 V, 3.00 A
 26.19 1010 s
 26.21 (a) 0.100 A (b) $P_{400} = 4.0$ W, $P_{800} = 8.0$ W
 (c) 12.0 W (d) $I_{400} = 0.300$ A, $I_{800} = 0.150$ A
 (e) $P_{400} = 36.0$ W, $P_{800} = 18.0$ W (f) 54.0 W
 (g) series: 800- Ω bulb; parallel: 400- Ω bulb
 (h) parallel
 26.23 (a) 2.00 A (b) 5.00 Ω (c) 42.0 V (d) 3.50 A
 26.25 (a) 8.00 A (b) $\mathcal{E}_1 = 36.0$ V, $\mathcal{E}_2 = 54.0$ V
 (c) 9.00 Ω
 26.27 (a) 1.60 A (top), 1.40 A (middle),
 0.20 A (bottom) (b) 10.4 V
 26.29 (a) 36.4 V (b) 0.500 A
 26.31 (a) 2.14 V, a (b) 0.050 A, 0; down
 26.33 (a) 0.641 Ω (b) 975 Ω
 26.35 (a) 17.9 V (b) 22.7 V (c) 21.4%
 26.37 (a) 0.849 μF (b) 2.89 s
 26.39 (a) 0 (b) 245 V (c) 0 (d) 32.7 mA
 (e) (a): 245 V; (b): 0; (c): 1.13 mC; (d): 0
 26.41 (a) 4.21 ms (b) 0.125 A
 26.43 192 μC
 26.45 13.6 A
 26.47 (a) 0.937 A (b) 0.606 A
 26.49 (a) 165 μC (b) 463 Ω (c) 12.6 ms
 26.51 900 W
 26.53 (a) $U_C = \frac{1}{2}C\mathcal{E}^2$ (b) $U_C = C\mathcal{E}^2$
 (c) $U_R = \frac{1}{2}C\mathcal{E}^2$ (d) 1/2, 1/2
 26.55 (a) 2.29 A (b) 3.71 A (c) 1.43 A
 26.57 (a) +0.22 V (b) 0.464 A
 26.59 $I_1 = 0.848$ A, $I_2 = 2.14$ A, $I_3 = 0.171$ A
 26.63 (a) 109 V; no (b) 13.5 s
 26.67 (a) -12.0 V (b) 1.71 A (c) 4.21 Ω
 26.69 (a) $\epsilon_0\rho K$ (b) 1.28 nC (c) 165 s (d) 2.68 pA
 26.71 (a) 11.8 pC (b) 0.917 A (c) 4.92 pJ (d) 703 pJ
 (e) 1300 pJ (f) 595 pJ
 26.73 (a) 114 V (b) 263 V (c) 266 V
 26.75 (a) 18.0 V (b) a (c) 6.00 V
 (d) both decrease by 36.0 μC
 26.77 1.7 M Ω , 3.1 μF
 26.79 (a) -1.23 ms (slope), 79.5 μC (y-intercept)
 (b) 247 Ω , 15.9 V (c) 1.22 ms (d) 11.9 V
 26.81 (a) $V_{\text{out}} = 21\text{V} - 10V_{\text{in}}$ (b) 1.35 V (c) -10
 26.85 (b) 4 (c) 3.2 M Ω , 4.0×10^{-3} (d) 3.4×10^{-4}
 (e) 0.88
 26.87 Choice (d)

Chapter 27

- 27.1 (a) $(-6.68 \times 10^{-4} \text{ N})\hat{k}$
 (b) $(6.68 \times 10^{-4} \text{ N})\hat{i} + (7.27 \times 10^{-4} \text{ N})\hat{j}$
 27.3 (a) positive (b) 0.0505 N
 27.5 172 m/s, -x-direction
 27.7 (a) 1.46 T, in the xz-plane at 40° from the +x-axis
 toward the -z-axis
 (b) 7.47×10^{-16} N, in the xz-plane at 50° from the
 -x-axis toward the -z-axis
 27.9 (a) 3.05 mWb (b) 1.83 mWb (c) 0
 27.11 -0.78 mWb
 27.13 (a) 0 (b) -0.0115 Wb (c) +0.0115 Wb (d) 0
 27.15 (a) 0.160 mT, into the page (b) 0.111 μs
 27.17 7.93×10^{-10} N, toward the south
 27.19 (a) 2.84×10^6 m/s, negative (b) yes (c) same
 27.21 0.838 mT
 27.23 (a) 7900 N/C, $\hat{i}$ (b) 7900 N/C, $\hat{i}$
 27.25 0.0445 T, out of the page
 27.27 (a) 4.92 km/s (b) 9.96×10^{-26} kg
 27.29 0.724 N, 63.4° below the current direction in the
 upper wire segment
 27.31 (a) 817 V (b) 113 m/s²
 27.33 (a) a (b) 3.21 kg
 27.35 (a) 55 turns (b) counterclockwise
 (c) 8.73 N $\cdot$ m, upward

- 27.37 (b) $F_{cd} = 1.20$ N (c) 0.420 N $\cdot$ m
 27.39 (a) A_2 (b) 290 rad/s²
 27.41 (a) $-NIAB\hat{i}$, 0 (b) 0, $-NIAB$ (c) $+NIAB\hat{i}$, 0
 (d) 0, $+NIAB$
 27.43 (a) 1.13 A (b) 3.69 A (c) 98.2 V (d) 362 W
 27.45 (a) 4.7 mm/s (b) $+4.5 \times 10^{-3}$ V/m,
 +z-direction (c) 53 μV
 27.47 (a) $x = 0.565$ m, $y = 0.825$ m, $z = 0.0900$ m
 (b) 31.3 m/s
 27.49 (a) 8.3×10^6 m/s (b) 0.14 T
 27.51 4.81×10^7 C/kg
 27.53 1.6 mm
 27.55 $\frac{Mg \tan \theta}{LB}$, right to left
 27.57 (a) 8.46 mT (b) 27.2 cm (c) 2.2 cm; yes
 27.59 (a) ILB , to the right (b) $\frac{v^2 m}{2ILB}$ (c) 1960 km
 27.61 (a) 1.13 N, toward the origin
 (b) 0.800 N, +z-direction
 27.63 (a) 48,000 A (b) not feasible (c) 19 G
 (d) 0.83 N
 27.65 0.024 T, +y-direction
 27.67 (a) $F_{PQ} = 0$; $F_{RP} = 12.0$ N, into the page;
 $F_{QR} = 12.0$ N, out of the page (b) 0
 (c) $\tau_{PQ} = \tau_{RP} = 0$; $\tau_{QR} = 3.60$ N $\cdot$ m
 (d) 3.60 N $\cdot$ m; yes (e) out
 27.69 $-(0.444 \text{ N})\hat{j}$
 27.71 (b) left: $(B_0 LI/2)\hat{i}$; top: $-IB_0 L\hat{j}$;
 right: $-(B_0 LI/2)\hat{i}$; bottom: 0 (c) $-IB_0 L\hat{j}$
 27.73 (a) $\frac{Mg}{2WNB}$ (b) $R(1 + \pi/2)$
 (c) $Mg(\sigma - 1)\sqrt{h(2R - h)}$
 (d) $\left[\sigma \cos\left(\frac{h}{R} - 1\right) - 1 \right] MgR$
 (e) $(1 - \sigma)Mgh$
 (f) $\left\{ \frac{h}{R} - \sigma \left[\sin\left(\frac{h}{R} - 1\right) + 1 \right] \right\} (MgR)$
 (g) $R[1 + \arccos(1/\sigma)]$ (h) $\sigma < \frac{1}{2}\left(1 + \frac{\pi}{2}\right)$
 27.75 (a) $-IA\hat{k}$ (b) $B_x = \frac{3D}{IA}$, $B_y = \frac{4D}{IA}$, $B_z = -\frac{12D}{IA}$
 27.77 (b) 1.85×10^{-28} kg (c) 1.20 kV
 (d) 8.32×10^5 m/s
 27.79 (a) $r = R \sin \theta$ (b) $dI = \sigma \omega R^2 \sin \theta d\theta$
 (c) $d\mu = \pi R^4 \sigma \omega \sin^3 \theta d\theta$ (d) $\frac{Q\omega R^2}{3} \hat{k}$
 (e) $\frac{Q\omega BR^2}{3} (-\sin \alpha \hat{j} - \cos \alpha \hat{i}) B$
 27.81 (a) 5.14 m (b) 1.72 μs (c) 6.08 mm
 (d) 3.05 cm
 27.83 Choice (c)
 27.85 Choice (a)

Chapter 28

- 28.1 (a) $-(19.2 \mu\text{T})\hat{k}$ (b) 0 (c) $(19.2 \mu\text{T})\hat{i}$
 (d) $(6.79 \mu\text{T})\hat{i}$
 28.3 (a) 60.0 nT, out of the page at A and B
 (b) 0.120 μT , out of the page (c) 0
 28.5 (a) 0 (b) $-(1.31 \mu\text{T})\hat{k}$ (c) $-(0.462 \mu\text{T})\hat{k}$
 (d) $(1.31 \mu\text{T})\hat{j}$
 28.7 (a) left (b) 0 (c) 1.21 pT, out of the page
 (d) 1.21 pT, into the page
 28.9 (a) 0.440 μT , out of the page (b) 16.7 nT, out of
 the page (c) 0
 28.11 (a) $(50.0 \text{ pT})\hat{j}$ (b) $-(50.0 \text{ pT})\hat{i}$
 (c) $-(17.7 \text{ pT})(\hat{i} - \hat{j})$ (d) 0
 28.13 17.6 μT , into the page
 28.15 (a) 0.8 mT (b) 40 μT (20 times larger)
 28.17 25 μA
 28.19 (a) $-(0.10 \mu\text{T})\hat{i}$ (b) 2.19 μT , at 46.8° from the
 +x-axis to the +z-axis
 (c) $(7.9 \mu\text{T})\hat{i}$

- 28.21 (a) 0 (b) 6.67 μT , toward the top of the page
 (c) 7.54 μT , to the left
 28.23 (a) 0 (b) 0 (c) 0.40 mT, to the left
 28.25 (a) P : 41 μT , into the page; Q : 25 μT , out of the
 page (b) P : 9.0 μT , out of the page; Q : 9.0 μT ,
 into the page
 28.27 $\frac{\mu_0 I^2}{2\pi g \lambda}$
 28.29 (a) 6.00 μN ; repulsive (b) 24.0 μN
 28.31 0.38 μA
 28.33 $\frac{\mu_0 |I_1 - I_2|}{4R}$; 0
 28.35 18.0 A, counterclockwise
 28.37 (a) 305 A (b) -3.83×10^{-4} T $\cdot$ m
 28.39 (a) $\mu_0 I / 2\pi r$ (b) 0
 28.41 (a) $\frac{\mu_0 I_1}{2\pi r}$ (b) $\frac{\mu_0 (I_1 + I_2)}{2\pi r}$
 28.43 (a) 1790 turns per meter (b) 63.0 m
 28.45 (a) 3.72 MA (b) 124 kA (c) 237 A
 28.47 (a) (i) 1.13 mT (ii) 4.68 MA/m (iii) 5.88 T
 28.49 (a) 1.00 μT , into the page (b) $(74.9 \text{ nN})\hat{j}$
 28.51 0.200 m
 28.53 (a) $\mu_0 I^2 R / d$ (b) $\frac{4B^2}{\mu_0 d} \left(\frac{A}{\pi} \right)^{3/2}$
 (c) $\frac{1}{2} \sqrt{\mu_0 F d} \left(\frac{\pi}{A} \right)^{3/4}$
 28.55 (a) 5.7×10^{12} m/s², away from the wire
 (b) 32.5 N/C, away from the wire (c) no
 28.59 (a) 2.00 A, out of the page (b) 2.13 μT , to the
 right (c) 2.06 μT
 28.61 23.2 A
 28.63 (a) $I_0 = 2\pi b \delta (1 - e^{-a/\delta})$, 81.5 A (b) $\frac{\mu_0 I_0}{2\pi r}$
 (c) $\left(\frac{e^{r/\delta} - 1}{e^{a/\delta} - 1} \right) I_0$ (d) $\frac{\mu_0 I}{2\pi r} \left(\frac{e^{r/\delta} - 1}{e^{a/\delta} - 1} \right)$
 (e) $r = \delta$: 175 μT ;
 $r = a$: 326 μT ;
 $r = 2a$: 163 μT
 28.65 (a) $\frac{3I}{2\pi R^3}$ (b) (i) $B = \frac{\mu_0 I r^2}{2\pi R^3}$ (ii) $B = \frac{\mu_0 I}{2\pi r}$
 28.67 (b) $B = \frac{\mu_0 I_0}{2\pi r}$ (c) $\frac{I_0 r^2}{a^2} \left(2 - \frac{r^2}{a^2} \right)$
 (d) $B = \frac{\mu_0 I_0 r}{2\pi a^2} \left(2 - \frac{r^2}{a^2} \right)$
 28.69 (a) $B = \mu_0 I n / 2$, +x-direction
 (b) $B = \mu_0 I n / 2$, -x-direction
 28.71 (a) $\frac{Q_1 \omega_1}{2\pi}$ (b) $\frac{\mu_0 \omega_1 Q_1}{2\pi H} \hat{k}$
 (c) $\frac{\mu_0 Q_1 Q_2 \omega_1 \omega_2 R^2}{8\pi H} \sin \theta$
 (d) $\frac{1}{2} M R^2 \omega_2$ (e) $\frac{\mu_0}{4\pi H} \frac{Q_1 Q_2 \omega_1 \sin \theta}{M}$
 28.73 (a) no (c) 64.9 A, 1.19 cm
 28.75 (a) nq (b) $n^{2/3} q$ (c) $\frac{1}{2} \mu_0 n q v R \hat{\phi}$
 (d) $n^{2/3} q v R d \hat{\phi}$ (e) $\frac{\mu_0}{2} n^{5/3} q^2 v^2 R^2 L d \hat{\phi}$
 (f) $\pi \mu_0 n^{5/3} q^2 v^2 R^2 L$, $\frac{\mu_0}{2} n^{5/3} q^2 v^2 R^2$
 (g) 1.9×10^{-5} Pa
 28.77 (a) $\frac{Q\omega}{4\pi} \sin \theta d\theta$ (b) $\frac{1}{4} Q \omega R^2 \sin^3 \theta d\theta$, +z-direction
 (c) $\frac{Q\omega R^2}{3}$, +z-direction
 (d) $\frac{2}{3} M R^2 \omega$ (e) $\frac{Q}{2M}$ (f) 1
 28.79 (b) $\frac{1}{2g} \left(\frac{\mu_0 Q_0^2}{4\pi \lambda R C d} \right)^2$
 28.81 Choice (b)
 28.83 Choice (c)

Chapter 29

- 29.1 (a) 17.1 mV (b) 28.5 mA
 29.3 (a) Wb/s, Wb/s³ (b) $\alpha = (0.750 \text{ s}^2)\beta$
 (c) -1.20 V
 29.5 (a) 34 V (b) counterclockwise
 29.7 (a) $\mu_0 i / 2\pi r$, into the page (b) $\frac{\mu_0 i}{2\pi r} L dr$
 (c) $\frac{\mu_0 i L}{2\pi} \ln(b/a)$ (d) $\frac{\mu_0 L}{2\pi} \ln(b/a) \frac{di}{dt}$
 (e) 0.506 μV
 29.9 (a) 5.44 mV (b) clockwise
 29.11 (a) 12.6 mA (b) 626 μA
 29.13 10.4 rad/s
 29.15 (a) counterclockwise (b) clockwise
 (c) no induced current
 29.17 (a) C: counterclockwise; A: clockwise
 (b) toward the wire
 29.19 (a) a to b (b) b to a (c) b to a
 29.21 (a) clockwise (b) no induced current
 (c) counterclockwise
 29.23 (a) 0.72 V (b) 0
 29.25 (a) 0.675 V (b) b (c) 2.25 V/m, b to a (d) b
 (e) (i) 0 (ii) 0
 29.27 (a) 3.00 V (b) b to a (c) 0.800 N, to the right
 (d) 6.00 W for each
 29.29 (a) counterclockwise (b) 42.4 mW
 29.31 35.0 m/s, to the right
 29.33 (a) 0.225 A, clockwise (b) 0
 (c) 0.225 A, counterclockwise
 29.35 9.21 A/s
 29.37 0.950 mV
 29.39 (a) 0.599 nC (b) 6.00 mA (c) 6.00 mA
 29.41 (a) 2.2 nA (b) 0.75 s
 29.43 (a) $2.5 \times 10^5 \text{ C}$ (b) 65 A (c) 11 T
 29.45 (a) 3.7 A (b) 1.33 mA (c) counterclockwise
 29.47 16.2 μV
 29.49 (a) $\frac{\mu_0 I a b v}{2\pi r(r+a)}$ (b) clockwise
 29.51 (a) 17.9 mV (b) a to b
 29.53 $\mu_0 I W / 4\pi$
 29.55 (a) $\frac{\mu_0 I v}{2\pi} \ln(1 + L/d)$ (b) a (c) 0
 29.57 (a) 0.165 V (b) 0.165 V (c) 0; 0.0412 V
 29.59 (a) $B^2 L^2 v / R$
 29.61 $a: \frac{qr dB}{2 dt}$, to the left; $b: \frac{qr dB}{2 dt}$, toward the top of the
 page; $c: 0$
 29.63 5.0 s
 29.65 (a) 0.3071 s⁻¹ (b) 3.69 T (c) a (d) 2.26 s
 29.67 (a) $\vec{v} - (1.26 \text{ m/s}) \sin \theta \hat{i}$
 (b) $-1.26 \sin \theta \text{ T} \cdot \text{m/s} \hat{j}$
 (c) $(10.0 \text{ cm})(\cos \theta \hat{j} - \sin \theta \hat{k}) d\theta$
 (d) 4.73 A (e) upward
 29.69 (a) a to b (b) $\frac{R m g \tan \phi}{L^2 B^2 \cos \phi}$ (c) $\frac{m g \tan \phi}{LB}$
 (d) $\frac{R m^2 g^2 \tan^2 \phi}{L^2 B^2}$ (e) $\frac{R m^2 g^2 \tan^2 \phi}{L^2 B^2}$; same
 29.71 Choice (c)
 29.73 Choice (a)

Chapter 30

- 30.1 (a) 0.270 V; yes (b) 0.270 V
 30.3 (a) 1.96 H (b) 7.11 mWb
 30.5 (a) 1940 (b) 800 A/s
 30.7 (a) 0.250 H (b) 0.450 mWb
 30.9 (a) 4.68 mV (b) a
 30.11 (b) 0.111 μH
 30.13 0.0300 T

- 30.15 2850
 30.17 (a) 0.161 T (b) 10.3 kJ/m³ (c) 0.129 J
 (d) 40.2 μH
 30.19 91.7 J
 30.21 (a) 2.40 A/s (b) 0.800 A/s (c) 0.413 A
 (d) 0.750 A
 30.23 (a) 17.3 μs (b) 30.7 μs
 30.25 15.4 V
 30.27 (a) 0.250 A (b) 0.137 A (c) 32.9 V; c
 (d) 0.462 ms
 30.29 15.3 V
 30.31 (a) 0 (b) 0 (c) $\mathcal{E}^2 / 4R$
 30.33 (a) 443 nC (b) 358 nC
 30.35 (a) 105 rad/s, 59.6 ms (b) 0.720 mC (c) 4.32 mJ
 (d) -0.542 mC (e) -0.050 A, counterclockwise
 (f) $U_C = 2.45 \text{ mJ}$, $U_L = 1.87 \text{ mJ}$
 30.37 (a) 7.50 μC (b) 15.9 kHz (c) 21.2 mJ
 30.39 (a) 298 rad/s (b) 83.8 Ω
 30.43 (a) 59 NH (b) 26.8 μV
 30.45 20 km/s; about 30 times smaller
 30.47 (a) 5.00 H (b) 31.7 m; no
 30.49 222 μF , 9.31 μH
 30.51 (a) 111 N (b) 37.1 ms (c) $4.00 \times 10^3 \text{ rad/s}^2$
 30.53 (a) 15.0 mC (b) 93.8 μs (c) 2.67 kHz
 (d) 11.3 J (e) 1.30 ms
 30.55 (a) 24.0 mV (b) 1.55 mA (c) 72.1 nJ
 (d) 5.20 μC , 18.0 nJ
 30.57 (a) 0, 20.0 V (b) 0.267 A, 0 (c) 0.147 A, 9.0 V
 30.59 (b) 5.0 Ω , 8.5 H (c) 1.7 kJ; 2.0 kW
 30.61 (a) 60.0 V (b) a (c) 60.0 V (d) c
 (e) -96.0 V (f) b (g) 156 V (h) d
 30.63 (a) 0; $v_{ac} = 0$, $v_{cb} = 36.0 \text{ V}$
 (b) 0.180 A, $v_{ac} = 9.0 \text{ V}$, $v_{cb} = 27.0 \text{ V}$
 (c) $i_0 = (0.180 \text{ A})(1 - e^{-t/(0.020 \text{ s})})$,
 $v_{ac} = (9.0 \text{ V})(1 - e^{-t/(0.020 \text{ s})})$,
 $v_{cb} = (9.0 \text{ V})(3.00 + e^{-t/(0.020 \text{ s})})$
 30.65 (a) $A_1 = A_4 = 0.455 \text{ A}$, $A_2 = A_3 = 0$
 (b) $A_1 = 0.585 \text{ A}$, $A_2 = 0.320 \text{ A}$, $A_3 = 0.160 \text{ A}$,
 $A_4 = 0.107 \text{ A}$
 30.67 (a) $v_L = \frac{\mathcal{E}}{R + R_L}(R_L + R e^{-(R+R_L)t/L})$
 (b) 50.0 V (c) 30.0 V; 3.00 A (d) 6.67 Ω
 (e) 40.0 mH
 30.69 (a) $\frac{\mu_0 I (N_1 - N_2) \pi a^2}{\lambda}$ (b) $\frac{\mu_0 I \pi (N_1 b^2 - N_2 a^2)}{\lambda}$
 (c) $\frac{\mu_0 \pi}{\lambda} (N_1^2 b^2 - N_2^2 a^2)$
 (d) $\frac{\mu_0 \pi}{\lambda} (N_1^2 b^2 + 2N_1 N_2 a^2 + N_2^2 a^2)$
 (e) 10.3 mH (f) 16.0 mH
 30.71 (a) $i_1 = \frac{\mathcal{E}}{R_1}(1 - e^{-R_1 t/L})$, $i_2 = \frac{\mathcal{E}}{R_2} e^{-t/R_2 C}$,
 $q_2 = \mathcal{E} C (1 - e^{-t/R_2 C})$
 (b) $i_1 = 0$, $i_2 = 9.60 \text{ mA}$
 (c) $i_1 = 1.92 \text{ A}$, $i_2 = 0$; $t \gg L/R_1$
 and $t \gg R_2 C$ (d) 1.6 ms (e) 9.4 mA
 (f) 0.22 s
 30.73 Choice (b)
 30.75 Choice (c)

Chapter 31

- 31.1 1.06 A
 31.3 (a) 90°; lead (b) 193 Hz
 31.5 (a) 1510 Ω (b) 0.239 H (c) 497 Ω
 (d) 16.6 μF
 31.7 (a) 1700 Ω (b) 106 Ω (c) 150 Hz
 31.9 (a) $(12.5 \text{ V}) \cos[(480 \text{ rad/s})t]$ (b) 7.17 V
 31.11 (a) $i = (0.0253 \text{ A}) \cos[(720 \text{ rad/s})t]$
 (b) 180 Ω
 (c) $v_L = -(4.56 \text{ V}) \sin[(720 \text{ rad/s})t]$

- 31.13 (a) 601 Ω (b) 49.9 mA (c) -70.6°; lag
 (d) $V_R = 9.98 \text{ V}$, $V_L = 4.99 \text{ V}$, $V_C = 33.3 \text{ V}$
 31.15 50.0 V
 31.17 32.8°
 31.19 (a) 40.0 W (b) 0.167 A (c) 720 Ω
 31.21 (b) 32.4 Ω
 31.23 (a) 45.8°, 0.697 (b) 344 Ω (c) 155 V
 (d) 48.6 W (e) 48.6 W (f) 0 (g) 0
 31.25 (a) 0.302 (b) 0.370 W
 (c) 0.370 W (resistor), 0, 0
 31.27 (a) 113 Hz; 15.0 mA (b) 7.61 mA; lag
 31.29 (a) 150 V (b) $V_R = 150 \text{ V}$,
 $V_L = V_C = 1290 \text{ V}$ (c) 37.5 W
 31.31 (a) 1.00 (b) 75.0 W (c) 75.0 W
 31.33 (a) 945 rad/s (b) 70.6 Ω
 (c) $V_L = V_C = 450 \text{ V}$, $V_R = 120 \text{ V}$
 31.35 (a) 10 (b) 2.40 A (c) 28.8 W (d) 500 Ω
 31.37 0.124 H
 31.39 230 Ω
 31.41 $3.59 \times 10^7 \text{ rad/s}$
 31.43 (a) inductor (b) 0.133 H
 31.45 (a) 0.831 (b) 161 W
 31.47 $\frac{V_{\text{out}}}{V_s} = \sqrt{\frac{R^2 + \omega^2 L^2}{R^2 + \left(\omega L - \frac{1}{\omega C}\right)^2}}$
 31.51 (a) 102 Ω (b) 0.882 A (c) 270 V
 31.53 (a) $C = 5\epsilon_0 a^2 \theta / g$ (b) 131° (c) 7.41 V
 (d) 1230 kHz (e) 7.74 V
 31.55 (b) 5770 rad/s (c) 2.40 A (d) 2.40 A
 (e) 0.139 A (f) 0.139 A
 31.57 (a) $\omega = 28,800 \text{ rad/s}$ so $\phi = 60^\circ$
 (b) $P_R = 0.375 \text{ W}$, $P_L = P_C = 0$; 0.100 A
 31.59 (a) 0.750 A (b) 160 Ω (c) 341 Ω , 619 Ω
 (d) 341 Ω
 31.61 (a) $\frac{V}{R}$ (b) $\frac{V}{R} \sqrt{\frac{L}{C}}$ (c) $\frac{V}{R} \sqrt{\frac{L}{C}}$ (d) $\frac{1}{2} L \frac{V^2}{R^2}$
 (e) $\frac{1}{2} L \frac{V^2}{R^2}$
 31.63 20.0 Ω , 0.180 H
 31.65 (a) $\frac{V_{\text{in}}}{\sqrt{R^2 + (\omega L)^2}}$ (b) $\arctan\left(\frac{\omega L}{R}\right)$
 (c) $\frac{1}{\sqrt{1 + (R/\omega L)^2}}$ (d) $\frac{R}{2.00\pi L}$ (e) 1.6 mH
 31.67 (a) $\frac{1}{2} V_R I$ (b) 0 (c) 0
 31.69 Choice (b)
 31.71 Choice (d)

Chapter 32

- 32.1 (a) 1.28 s (b) $8.15 \times 10^{13} \text{ km}$
 32.3 13.3 nT, + y-direction
 32.5 $3.0 \times 10^{18} \text{ Hz}$, $3.3 \times 10^{-19} \text{ s}$, $6.3 \times 10^{10} \text{ rad/s}$
 32.7 (a) (i) 60 kHz (ii) $6.0 \times 10^{13} \text{ Hz}$
 (iii) $6.0 \times 10^{16} \text{ Hz}$
 (b) (i) $4.62 \times 10^{-14} \text{ m} = 4.62 \times 10^{-5} \text{ nm}$
 (ii) 508 m = $5.08 \times 10^{11} \text{ nm}$
 32.9 (a) +y-direction (b) 0.149 mm
 (c) $\vec{B} = (1.03 \text{ mT}) \cos[(4.22 \times 10^4 \text{ rad/m})y - (1.265 \times 10^{13} \text{ rad/s})t] \hat{i}$
 32.11 (a) 361 m (b) 0.0174 rad/m
 (c) $5.22 \times 10^6 \text{ rad/s}$ (d) 0.0144 V/m
 32.13 (a) 0.381 μm (b) 0.526 μm (c) 1.38
 (d) 1.90
 32.15 (a) 330 W/m² (b) 500 V/m, 1.7 μT
 32.17 $2.5 \times 10^{25} \text{ W}$
 32.19 93.8 W
 32.21 12.0 V/m, 40.0 nT
 32.23 (a) 0.18 mW (b) 274 V/m, 0.913 μT
 (c) 0.18 mJ/s (d) 0.010 W/cm²

- 32.25 (a) 637 W/m^2 (b) 693 V/m , $2.31 \mu\text{T}$
 (c) $2.12 \mu\text{J/m}^3$
- 32.27 (a) 30.5 cm (b) 2.46 GHz (c) 2.11 GHz
- 32.29 7.50 cm
- 32.31 (a) 0.375 mJ (b) 4.08 mPa
 (c) 604 nm , $3.70 \times 10^{14} \text{ Hz}$
 (d) 30.3 kV/m , $101 \mu\text{T}$
- 32.33 (a) $6.02 \times 10^{-3} \text{ W/m}^2$ (b) 2.13 V/m , 7.10 nT
 (c) 1.20 pN ; no
- 32.35 (a) at $r = R$: 64 MW/m^2 , 0.21 Pa ; at $r = R/2$:
 260 MW/m^2 , 0.85 Pa (b) no
- 32.37 $3.89 \times 10^{-13} \text{ rad/s}^2$
- 32.39 (a) $\rho I/\pi a^2$, in the direction of the current
 (b) $\mu_0 I/2\pi a$, counterclockwise if the current is out
 of the page
 (c) $\frac{\rho I^2}{2\pi^2 a^3}$, radially inward
 (d) $\frac{\rho I^2}{\pi a^2} = I^2 R$
- 32.41 (a) 1.364 m (b) 10.91 m
- 32.43 (a) $9.75 \times 10^{-15} \text{ W/m}^2$
 (b) $2.71 \mu\text{V/m}$, $9.03 \times 10^{-15} \text{ T}$, 67.3 ms
 (c) $3.25 \times 10^{-23} \text{ Pa}$ (d) 0.190 m
- 32.45 (a) $\frac{4\rho G\pi MR^3}{3r^2}$ (b) $\frac{LR^2}{4cr^2}$ (c) $0.19 \mu\text{m}$; no
- 32.47 (b) $3.00 \times 10^8 \text{ m/s}$
- 32.49 (a) $\frac{I_0}{c}(2 - e)$ (b) $3.3 \times 10^{-16} \text{ N}$ (c) 0.55
- 32.51 (b) 1.39×10^{-11} (c) 2.54×10^{-8}
- 32.53 (c) $66.0 \mu\text{m}$
- 32.55 Choice (d)

Chapter 33

- 33.1 39.4°
- 33.3 (a) $2.04 \times 10^8 \text{ m/s}$ (b) 442 nm
- 33.5 $2.11 \times 10^8 \text{ m/s}$, 417 nm , $5.07 \times 10^{14} \text{ Hz}$
- 33.7 (a) 47.5° (b) 66.0°
- 33.9 $2.51 \times 10^8 \text{ m/s}$
- 33.11 (a) 2.34 (b) 82°
- 33.13 38.5°
- 33.15 (a) 51.3° (b) 33.6°
- 33.17 (a) 58.1° (b) 22.8°
- 33.19 1.77
- 33.21 (a) 48.9° (b) 28.7°
- 33.23 0.6°
- 33.25 (a) red: 1.36 , violet: 1.40 (b) red: $2.21 \times 10^8 \text{ m/s}$,
 violet: $2.14 \times 10^8 \text{ m/s}$
- 33.27 $0.375I_0$
- 33.29 (a) $A: I_0/2$, $B: 0.125I_0$, $C: 0.0938I_0$ (b) 0
- 33.31 (a) 1.40 (b) 35.5°
- 33.33 6.38 W/m^2
- 33.35 (a) $0.374I$ (b) $2.35I$
- 33.37 (a) 46.7° (b) 13.4°
- 33.39 72.1°
- 33.41 1.28
- 33.43 3.52×10^4
- 33.45 1.84
- 33.47 (a) 48.6° (b) 48.6°
- 33.49 39.1°
- 33.51 (b) 0.23° ; about the same
- 33.53 23.3°
- 33.55 (a) A : 1.46, carbon tetrachloride;
 B : 1.33, water;
 C : 1.63, carbon disulfide;
 D : 1.50, benzene
 (b) A : 2.13, B : 1.77, C : 2.66, D : 2.25
 (c) all: $5.09 \times 10^{14} \text{ Hz}$
- 33.57 (a) 35° (b) $I_0 = 10.1 \text{ W/m}^2$, $I_p = 19.9 \text{ W/m}^2$

- 33.59 (a) $\theta = \arcsin\left(\frac{2R - d}{2R + d}\right)$ (b) $R = \frac{d(n_1 + n_2)}{2(n_1 - n_2)}$
 (c) 2.07 cm (d) $\approx 50 \text{ cm}$ (e) $1.49 \mu\text{s}$
- 33.61 (a) $\Delta = 2\theta_a^A - 6 \arcsin\left(\frac{\sin \theta_a^A}{n}\right) + 2\pi$
 (b) $\theta_2 = \arccos\sqrt{\frac{n^2 - 1}{8}}$
 (c) violet: $\theta_2 = 71.55^\circ$, $\Delta = 233.2^\circ$
 red: $\theta_2 = 71.94^\circ$, $\Delta = 230.1^\circ$; violet
- 33.63 Choice (d)

Chapter 34

- 34.1 39.2 cm to the right of the mirror, 4.85 cm
- 34.3 9.0 cm ; tip of the lead
- 34.5 (b) 33.0 cm to the left of the vertex, 1.20 cm ,
 inverted, real
- 34.7 0.213 mm
- 34.9 18.0 cm from the vertex; 0.50 cm , erect, virtual
- 34.11 (a) $+4.00$ (b) 48.0 cm to the right of the mirror;
 virtual
- 34.13 (a) concave (b) $f = 2.50 \text{ cm}$, $R = 5.00 \text{ cm}$
- 34.15 (a) 10.0 cm to the left of the shell vertex, 2.20 mm
 (b) 4.29 cm to the right of the shell vertex,
 0.944 mm
- 34.19 2.67 cm
- 34.21 3.30 m
- 34.23 (a) at the center of the bowl, 1.33 (b) no
- 34.25 39.5 cm
- 34.27 (a) 30 cm , 9.60 mm , inverted (b) 60.0 cm ,
 9.60 mm , upright
- 34.29 (a) 107 cm to the right of the lens, 17.8 mm ; real;
 inverted (b) the same
- 34.31 71.2 cm to the right of the lens; -2.97
- 34.33 3.69 cm ; 2.82 cm to the left of the lens
- 34.35 1.67
- 34.37 12.0 mm
- 34.39 -12 cm
- 34.41 26.3 cm from the lens, 12.4 mm ; erect; same side
- 34.43 (a) 200 cm to the right of the first lens, 4.80 cm
 (b) 150 cm to the right of the second lens,
 7.20 cm
- 34.45 (a) 53.0 cm (b) real (c) 2.50 mm ; inverted
- 34.47 (a) $s' = -2f$, virtual, left (b) $3h$, upright (c) yes
- 34.49 8.69 cm ; no
- 34.51 (a) 12 cm to the right of the converging lens
 (b) -8 cm (c) 24 cm to the right of the diverging
 lens (d) 12 cm to the right of the converging lens,
 -4 cm , 6 cm to the right of the diverging lens
- 34.53 (a) 80.0 cm (b) 76.9 cm
- 34.55 49.4 cm , 2.02 diopters
- 34.57 (a) far point: 202 cm , near point: 44.6 cm
 (b) 56.1 cm
- 34.59 (a) 6.06 cm (b) 4.12 mm
- 34.61 (a) -6.33 (b) 1.90 cm (c) 0.127 rad
- 34.63 (a) 0.661 m (b) 59.1
- 34.65 16.0 mm
- 34.67 (a) 20.0 cm (b) 39.0 cm
- 34.69 51 m/s
- 34.71 (a) 1.49 cm
- 34.73 (b) 2.4 cm ; -0.133
- 34.75 2.00
- 34.77 (a) converging, 52.5 cm from the lens
 (b) converging, 17.5 cm from the lens
- 34.79 converging, $+50.2 \text{ cm}$
- 34.81 (a) 58.7 cm , converging (b) 4.48 mm ; virtual
- 34.83 (a) 6.48 mm (b) no, behind the retina
 (c) 19.3 mm from the cornea; in front of the retina
- 34.85 (a) 6.49 cm (b) 6.76 cm $\leq s \leq 7.05 \text{ cm}$
 (c) 66.2 cm (d) 1.36 m/s , downward (e) 24.1 cm

- 34.87 (a) 30.9 cm (b) 29.2 cm
- 34.89 (b) first image: (i) 51.3 cm to the right of the lens
 (ii) real (iii) inverted
 second image: (i) 51.3 cm to the right of the lens
 (ii) real (iii) erect
- 34.91 -26.7 cm
- 34.93 7.06 cm to the left of the spherical mirror vertex,
 0.177 cm tall; 13.3 cm to the left of the spherical
 mirror vertex, 0.111 cm tall
- 34.95 134 cm to the left of the object
- 34.97 4.17 diopters
- 34.99 (d) 36.0 cm , 21.6 cm ; $d = 1.2 \text{ cm}$
- 34.101 (a) -16.6 cm (b) 20.0 cm to the right
- 34.103 (b) 10.0 cm (c) 90 cm in front of the glass, virtual
- 34.105 (a) $4f$
- 34.107 (b) 1.74 cm
- 34.109 Choice (d)
- 34.111 Choice (b)

Chapter 35

- 35.1 0.75 m , 2.00 m , 3.25 m , 4.50 m , 5.75 m ,
 7.00 m , 8.25 m
- 35.3 (a) 2.0 m (b) constructively
 (c) 1.0 m , destructively
- 35.5 (a) 11 (b) 2.50 m
- 35.7 1.14 mm
- 35.9 0.83 mm
- 35.11 (a) 39 (b) $\pm 73.3^\circ$
- 35.13 12.6 cm
- 35.15 1200 nm
- 35.17 (a) $0.750I_0$ (b) 80 nm
- 35.19 1670 rad
- 35.21 (a) 4.52 rad (b) $0.404I_0$
- 35.23 114 nm
- 35.25 (a) 55.6 nm (b) (i) 2180 nm
 (ii) 11.0 wavelengths
- 35.27 (a) 514 nm ; green (b) 603 nm ; orange
- 35.29 $0.11 \mu\text{m}$
- 35.31 0.570 mm
- 35.33 1.57
- 35.35 (a) 96.0 nm (b) no, no
- 35.37 (a) 1.58 mm (green), 1.72 mm (orange)
 (b) 3.45 mm (violet), 4.74 mm (green),
 5.16 mm (orange) (c) $9.57 \mu\text{m}$
- 35.39 1.730
- 35.41 761 m , 219 m , 90.1 m , 20.0 m
- 35.43 $6.8 \times 10^{-5} (\text{C}^\circ)^{-1}$
- 35.45 $1.33 \mu\text{m}$
- 35.47 600 nm , 467 nm ; no
- 35.49 (a) 1.54 (b) $\pm 15.0^\circ$
- 35.51 (a) 50 MHz (b) 237.0 m
- 35.53 (a) $\lambda/2$ (b) 6
 (c) $\pm 9.64 \text{ cm}$, $\pm 31.7 \text{ cm}$,
 $\pm 84.1 \text{ cm}$
 (d) 2, $\pm 11.3 \text{ cm}$ from the center
- 35.55 14.0
- 35.57 Choice (d)
- 35.59 Choice (c)

Chapter 36

- 36.1 506 nm
- 36.3 (a) 226 (b) $\pm 83.0^\circ$
- 36.5 9.07 m
- 36.7 (a) 63.8 cm
 (b) $\pm 22.1^\circ$, $\pm 34.3^\circ$, $\pm 48.8^\circ$, $\pm 70.1^\circ$
- 36.9 $\pm 16.0^\circ$, $\pm 33.4^\circ$, $\pm 55.6^\circ$
- 36.11 (a) 10.9 mm (b) 5.4 mm
- 36.13 (a) 580 nm (b) 0.128

- 36.15 (a) $\pm 13.0^\circ, \pm 26.7^\circ, \pm 42.4^\circ, \pm 64.1^\circ$
 (b) 2.08 W/m^2
 36.17 (a) $d = 2a$ (b) 0.101 W/m^2
 36.19 (a) 3 (b) 2
 36.21 (a) $0.0627^\circ, 0.125^\circ$ (b) $0.249I_0, 0.0256I_0$
 36.23 (a) 3.00 mm (b) 6.00 mm
 36.25 (a) 4830 lines/cm (b) 4; $\pm 37.7^\circ, \pm 66.5^\circ$
 36.27 (a) $2.40 \mu\text{m}$ (b) 417 slits/mm (c) $1.80 \mu\text{m}$
 36.29 (a) 2.69×10^4 (b) $m = 3$
 36.31 (a) 467 nm (b) 27.8°
 36.33 (a) 1820 slits (b) $41.0297^\circ, 0.0137^\circ$
 36.35 0.232 nm
 36.37 92 cm
 36.39 1.88 m
 36.41 220 m
 36.43 (a) 77 m (Hubble), 1100 km (Arecibo)
 (b) 1500 km
 36.45 $30.2 \mu\text{m}$
 36.47 (a) 78 (b) $\pm 80.8^\circ$ (c) $555 \mu\text{W/m}^2$
 36.49 1.68
 36.51 (b) $4.49 \text{ rad}, 7.73 \text{ rad}$ (c) $3.14 \text{ rad}, 6.28 \text{ rad},$
 9.42 rad ; no (d) $4.78^\circ, 6.84^\circ, 9.59^\circ$
 36.53 -0.033 mm ; decrease
 36.55 360 nm
 36.57 second
 36.59 (a) 65° (b) 0.16 nm (c) $4.4 \times 10^6 \text{ m/s}$
 (d) $2.6 \times 10^{16} \text{ Hz}$ (e) $3.3 \times 10^{-34} \text{ J} \cdot \text{s}$
 (f) $6.6 \times 10^{-34} \text{ J} \cdot \text{s}$ (g) yes

- 36.61 (a) 1.03 mm (b) 0.148 mm
 36.63 (a) $12.1 \mu\text{m}$ (b) $10.4 \text{ cm}, 15.2 \text{ cm}$
 36.65 (a) $10.6 \mu\text{m}$ (b) 532 nm/s (c) $1.60 \mu\text{m}$
 (d) 1.11 s (e) 92.7% (f) 2.88°
 36.69 Choice (d)
 36.71 Choice (a)

Chapter 37

- 37.1 bolt A
 37.3 $0.867c$; no
 37.5 (a) $0.998c$ (b) 126 m
 37.7 (a) 12.0 ms measured by the first officer
 (b) $0.997c$
 37.9 92.5 m
 37.11 (a) 0.66 km (b) $49 \mu\text{s}$; 15 km (c) 0.45 km
 37.13 (a) 3570 m (b) $90.0 \mu\text{s}$ (c) $89.2 \mu\text{s}$
 37.15 (a) $0.806c$ (b) $0.974c$ (c) $0.997c$
 37.17 (a) toward (b) $0.385c$
 37.19 $0.784c$
 37.21 $0.611c$
 37.23 (a) $0.159c$ (b) $\$172 \text{ million}$
 37.25 (a) 2.30 (b) 4.11
 37.27 $3.06p_0$
 37.29 (a) $0.866c$ (b) $0.608c$
 37.31 (a) $0.866c$ (b) $0.986c$
 37.33 (a) 0.450 nJ (b) $1.94 \times 10^{-18} \text{ kg} \cdot \text{m/s}$
 (c) $0.968c$

- 37.35 (a) 0.867 nJ (b) 0.270 nJ (c) 0.452
 37.37 (a) 5.34 pJ (nonrel), 5.65 pJ (rel), 1.06
 (b) 67.8 pJ (nonrel), 331 pJ (rel), 4.88
 37.39 (a) 2.06 MV (b) $0.330 \text{ pJ} = 2.06 \text{ MeV}$
 37.41 $0.700c$
 37.43 42.5 y
 37.45 (a) 1.0 kg (b) $A: 1.7 \text{ kg}, B: 1.8 \text{ kg}$
 37.47 (a) $\Delta = 9 \times 10^{-9}$ (b) $7000m$
 37.49 5.01 ns , clock on plane
 37.51 0.168 MeV
 37.53 (a) $1.08 \times 10^{14} \text{ J}$ (b) $2.70 \times 10^{19} \text{ W}$
 (c) $1.10 \times 10^{10} \text{ kg}$
 37.55 (a) $IHL\hat{j}$
 (b) $IHLB\hat{i}$
 (c) $\vec{\mu}' = \vec{\mu}/\gamma$
 (d) $\vec{B}' = B\gamma\hat{k}$ (e) $\vec{B}'_{\perp} = \vec{B}_{\perp}\gamma$
 37.57 $0.357c$; receding
 37.59 154 km/h
 37.61 $2.04 \times 10^{-13} \text{ N}$
 37.63 (a) $2.6 \times 10^{-8} \text{ s}$ (b) 0.97
 37.65 (a) $2.0 \times 10^{-18} \text{ kg}$ (b) $4.0 \times 10^4 \text{ m/s}^2$
 37.67 (a) $(0, 0), (-L\gamma, Lv\gamma/c^2), (-vT\gamma, T\gamma),$
 $((-L + vT)\gamma, (T + Lv/c^2)\gamma)$
 (c) cLT (d) cLT (e) yes
 37.69 (a) 2494 MeV (b) 2.526 times
 (c) 987.4 MeV , twice as much
 37.71 Choice (c)
 37.73 Choice (b)

CREDITS

Cover John Woodworth/Alamy

Front Matter John Woodworth/Alamy; iii: Hugh D. Young; iii: Roger A. Freedman; xiii: Brian A Jackson/Shutterstock; xiv: Blanscape/Shutterstock; xv: George Burba/Shutterstock; xvi: 06photo/Shutterstock; xvii: Tomatito/Shutterstock

Chapter 1 p. 1: Minerva Studio/Shutterstock; p. 2: Michele Perbellini/Shutterstock; p. 2: NASA; p. 5: Science History Images/Alamy Stock Photo; p. 6: Courtesy of Bruker Corporation; p. 6: DGLimages/Alamy Stock Photo; p. 6: ESB Basic/Shutterstock; p. 6: JPL/NASA; p. 6: NASA; p. 6: Sebastian Kaulitzki/Shutterstock; p. 8: Chronicle/Alamy Stock Photo; p. 10: Tyler Olson/Shutterstock

Chapter 2 p. 34: Vibe Images/Fotolia; p. 37: moodboard/Brand X Pictures/Getty Images; p. 37: Stefan Holm/Shutterstock; p. 46: NASA; p. 50: Richard Megna/Fundamental Photographs; p. 53: RUSS ROHDE/Getty Images; p. 53: Thomas_EyeDesign/E+/Getty Images

Chapter 3 p. 67: Feng Li/Getty Images Sport/Getty Images; p. 71: Luca Lozzi / Getty Images News/Getty Images; p. 72: Dominique Douieb/PhotoAlto Agency RF Collections/Getty Images; p. 75: Richard Megna /Fundamental Photographs; p. 77: Richard Megna /Fundamental Photographs; p. 84: hubblesite/STScI/NASA; p. 85: David Wall/Alamy; p. 86: Chris Putnam/123rf; p. 89: CERN

Chapter 4 p. 100: ilbusca/Getty Images; p. 105: FotoPrisma/Fotolia; p. 107: Wayne Eastep/The Image Bank/Getty Images; p. 110: FLUKY FLUKY/Shutterstock; p. 112: Cheryl A. Meyer/Shutterstock; p. 116: Kadmy/Fotolia; p. 119: Maksym Gorpenyuk/Shutterstock; p. 120: John W. McDonough/Sports Illustrated/Getty Images; p. 120: PCN Photography/Alamy Stock Photo; p. 120: Roy Pedersen/Fotolia; p. 127: www.autosnout.com

Chapter 5 p. 130: Brian A Jackson/Shutterstock; p. 139: NASA; p. 142: Efired/Shutterstock; p. 144: Alex Kosev/Shutterstock; p. 148: Eye of Science/Science Source; p. 149: dzphotovideo/Getty Images; p. 155: T-Photo/Shutterstock; p. 156: BrunoWeltmann/Shutterstock; p. 156: Comaniciu Dan/Shutterstock; p. 156: Jason Stitt/Shutterstock; p. 156: NASA/JPL/Space Science Institute; p. 166: Data from Rothschild, M. et al. "The Flying Leap of a Flea," Scientific American, Vol. 229, No. 5 (November 1973), pp. 92–101.; p. 169: From www.engineeringtoolbox.com

Chapter 6 p. 172: Brocreative/Shutterstock; p. 173: mariiya/Fotolia; p. 173: Steve Gschmeissner/Science Source; p. 181: Mikadun/Shutterstock; p. 186: Steve Gschmeissner/Science Source; p. 190: Anthony Hall/Fotolia; p. 190: Fox Photos/Stringer/Getty Images; p. 190: Gayvoronskaya_Yana/Shutterstock; p. 190: Keystone/Stringer/Shutterstock; p. 191: fandu/Fotolia; p. 200: www.sciencebuddies.org

Chapter 7 p. 203: Nagel Photography/Shutterstock; p. 204: Otvalo/Shutterstock; p. 205: Erni/Shutterstock; p. 205: Robert F. Bukaty/Apiimages; p. 207: hinnamsaisuy/Shutterstock; p. 212: threerocksimages/Shutterstock; p. 214: G. Ronald Austing/Science Source; p. 214: JNP/Shutterstock; p. 220: 135pixels/Shutterstock; p. 220: Kletr/Shutterstock; p. 223: LaiQuocAnh/Shutterstock; p. 224: Thorsten Eckert/Alamy; p. 235: www.engineershandbook.com

Chapter 8 p. 237: Colorful High Speed Photographs/Getty Images; p. 238: Alex Emanuel Koch/Shutterstock; p. 239: Willie Linn/Shutterstock; p. 240: Jim Cummins/Getty Images; p. 240: Vereshchagin Dmitry/Shutterstock; p. 242: Ted Kinsman/Science Source; p. 248: FPG/Getty Images; p. 251: David Leah/Getty Images; p. 256: Richard Megna /Fundamental Photographs; p. 258: Elliott Rusty Harold/Shutterstock; p. 259: NASA; p. 270: From www.chuckhawks.com

Chapter 9 p. 273: Blanscape/Shutterstock; p. 274: connel/Shutterstock; p. 278: Hybrid Medical Animation/Science Source; p. 285: Ammit Jack/Shutterstock; p. 285: Dan Rodney/Shutterstock; p. 288: David J. Phillip/Associated Press; p. 288: DenisNata/Shutterstock; p. 290: NASA; p. 300: NASA

Chapter 10 p. 303: Dan Rodney/Shutterstock; p. 307: 68/Ocean/Corbis; p. 311: Bruce MacQueen/Shutterstock; p. 311: David Lentink/Science Source; p. 311:

Robert Young/Fotolia; p. 313: gorillaimages/Shutterstock; p. 315: Chris DeRidder/Shutterstock; p. 316: Bjorn Heller/Shutterstock; p. 320: NHPA/Photoshot

Chapter 11 p. 339: nito/Shutterstock; p. 342: haveseen/Shutterstock; p. 344: Maridav/Shutterstock; p. 347: Andrew Bret Wallis/Getty Images; p. 347: Benjamin Marin Rubio/Shutterstock; p. 347: Djomas/Shutterstock; p. 347: Richard Carey/Fotolia; p. 350: Dante Fenolio/Science Source

Chapter 12 p. 366: Isabelle Kuehn/Shutterstock; p. 366: orlandin/Shutterstock; p. 367: travelview/Shutterstock; p. 370: Pascal, Blaise. 1653. Treatise on the Equilibrium of Liquids.; p. 370: SPL/Science Source; p. 371: Lisa F. Young/Shutterstock; p. 372: Andrey Jitkov/Shutterstock; p. 373: Archimedes; p. 375: Pali A/fotolia; p. 376: Pearson; p. 376: Popov Nikolay/Shutterstock; p. 377: Fribus Mara/Shutterstock; p. 379: Sunychka Sol/Shutterstock; p. 382: TechWizard/Shutterstock; p. 383: Edward Lara/Shutterstock; p. 383: Jurik Peter/Shutterstock; p. 383: VitaminCo/Shutterstock; p. 384: Harold Edgerton/MIT courtesy of Palm Press, INC.

Chapter 13 p. 395: NASA/JPL-Caltech/SSI/Cornell; p. 395: Newton published the law of gravitation in 1687. Philosophiae naturalis principia mathematica (MacLehose, 1726) [1871] Reprinted for Sir William Thomson and Hugh Blackburn.; p. 397: JPL/Cornell University/NASA; p. 397: NASA/JPL/University of Arizona; p. 399: Robert Nemiroff & Jerry Bonnell/NASA; p. 400: NASA; p. 400: swisshippo/Fotolia; p. 404: NASA; p. 405: Based on Newton, Isaac. 1728. A Treatise on the System of the World.; p. 405: NASA/Dembinsky Photo Associates/Alamy Stock Photo; p. 406: M.Showalter/NASA; p. 407: Kepler, Johannes. 1597. Mysterium Cosmographicum.; p. 409: NASA; p. 410: NASA; p. 416: John Michell, (1784). "On the Means of Discovering the Distance, Magnitude, & c. of the Fixed Stars, in Consequence of the Diminution of the Velocity of Their Light, in Case Such a Diminution Should be Found to Take Place in any of Them, and Such Other Data Should be Procured from Observations, as Would be Farther Necessary for That Purpose". Philosophical Transactions of the Royal Society 74 (0): 35–57.; p. 417: NASA; p. 418: NASA; p. 419: These images/animations were created by Prof. Andrea Ghez and her research team at UCLA and are from data sets obtained with the W. M. Keck Telescopes.; p. 426: ssd.jpl.nasa.gov; p. 427: nssdc.gsfc.nasa.gov/planetary/factsheet; p. 428: Based on S. Seager et al., (2007) "Mass–Radius Relationships for Solid Exoplanets" The Astrophysical Journal, vol. 669, no.2, p. 1297.; arXiv:0707.2895 [astro-ph].

Chapter 14 p. 429: blurAZ/Shutterstock; p. 430: Steve Byland/Shutterstock; p. 434: American Diagnostic Corporation; p. 446: Zurijeta/Shutterstock; p. 449: Christopher Griffin/Alamy Stock Photo; p. 452: symbiot/Shutterstock; p. 453: sonya etchison/Shutterstock

Chapter 15 p. 464: Walter D. Mooney/U.S. Geological Survey; p. 465: MARCO POLO COLLECTION/Balan Madhavan / Alamy; p. 466: imageBROKER/Alamy; p. 467: EpicStockMedia/Shutterstock; p. 476: Kodda/Shutterstock; p. 478: Christian Delbert/shutterstock; p. 483: Richard Megna /Fundamental Photographs; p. 487: Shmeliova Natalia/Shutterstock; p. 488: NSO/AURA/NSF

Chapter 16 p. 501: Eduard Kyslynsky/Shutterstock; p. 505: Anton Havelaar/123rf; p. 505: BERG, RICHARD E.; STORK, DAVID G., THE PHYSICS OF SOUND, 1st Ed., ©1982. Reprinted and Electronically reproduced by permission of Pearson Education, Inc., Upper Saddle River, New Jersey.; p. 505: Ollyy/Shutterstock; p. 506: deusexlupus/123rf; p. 506: Steve Thorne/Redferns/Getty Images; p. 508: Kretztechnik/Science Source; p. 511: Pius Lee/Shutterstock; p. 516: Digoarpi/Shutterstock; p. 519: B-D-S Piotr Marcinski/Shutterstock; p. 519: Martin Bough/Fundamental Photographs; p. 522: Roger A. Freedman; p. 523: Gino Santa Maria/Shutterstock; p. 526: cleanfotos/Shutterstock; p. 529: NASA; p. 30: NASA

Chapter 17 p. 541: huyangshu/Shutterstock; p. 545: David Thyberg/Shutterstock; p. 545: hofhauser/Shutterstock; p. 545: luckyraccoon/123rf.com; p. 545: Roger A. Freedman; p. 548: Evannovostro/Shutterstock; p. 550: Eric Schrader/Pearson Education, Inc.; p. 552: Hugh D. Young; p. 554: Roman Sigaev/Shutterstock; p. 556: Adam Hart-Davis/Science Source; p. 557: Richard Megna/Fundamental Photographs; p. 558: NASA; p. 558: Wavebreak Media Ltd/123rf; p. 562: DmitryND/Getty Images; p. 562: Paul Nicklen/ Getty Images; p. 564: Hugh D. Young; p. 565: Ted Kinsman/Science Source; p. 567: From https://data.giss.nasa.

C-2 Credits

gov/gistemp/graphs/; p. 567: Trends: A Compendium of Data on Global Change, U.S. Department of Energy's Carbon Dioxide Information Analysis Center, <http://cdiac.esd.ornl.gov/trends/co2/contents.htm>; p. 575: Nature Production/Nature Picture Library

Chapter 18 p. 579: Stockyimages/Shutterstock; p. 581: Topham/The Image Works; p. 586: Courtesy of Bruker Corporation; p. 590: Rich Iwasaki/Getty Images; p. 591: ARCTIC IMAGES/Alamy Stock Photo; p. 593: David Grossman/The Image Works; p. 599: Ray Coleman/Science Source; p. 600: Melissa King/Shutterstock; p. 606: NAO/AURA/NSF; p. 611: T.A. Rector/University of Alaska Anchorage, H. Schweiker/WIYN and NAO/AURA/NSF

Chapter 19 p. 613: Ken Paul/Getty Images; p. 614: John P. Surey; p. 614: Phumphao Sumrankong/Shutterstock; p. 614: Stock/Getty Images; p. 619: Diego Cervo/Shutterstock; p. 620: Liv friis-larsen/Shutterstock; p. 624: Rob Byron/Shutterstock; p. 624: Tom Branch/Science Source; p. 630: dean bertoncelj/Shutterstock

Chapter 20 p. 642: George Burba/Shutterstock; p. 644: James Morgan/Shutterstock; p. 645: Tyler Olson/Shutterstock; p. 653: Bill Bachman/Science Source; p. 658: Staff Sgt. Robert Zoellner; p. 659: cowardlion/Shutterstock; p. 664: DenisNata/Shutterstock; p. 665: Eric Schrader/Pearson Education; p. 670: Eric Schrader/Pearson Education; p. 677: Based on the Atlantic Ocean Atlas, part of the Woods Hole Oceanographic Institution Atlas Series, Vol. 1. (Woods Hole: Woods Hole Oceanographic Institution, 1960). From <https://commons.wikimedia.org/wiki/File:Thermocline.jpg>

Chapter 21 p. 678: Mikael Damkier/Shutterstock; p. 682: Clayton Stalter/The Image Works; p. 684: George Resch/Fundamental Photographs; p. 685: Knape/Getty Images; p. 690: Matt9122/Shutterstock; p. 701: otnaydur/Shutterstock; p. 701: Richard Megna / Fundamental Photographs; p. 704: Mark Smith/Science Source; p. 716: Data from Dominic Clarke et al., "Detection and Learning of Floral Electric Fields by Bumblebees," Science, April 5, 2013; Vol. 340, no. 6128, pp. 66–69.

Chapter 22 p. 718: U. S. Department of Energy; p. 723: Mark Webster/Getty Images; p. 725: Akg-images; p. 734: Dr. David Furness, Keele University/Science Source; p. 736: CTK/Alamy Stock Photo; p. 736: yevgeniy11/Shutterstock

Chapter 23 p. 747: Ted Kurihara Photography/Getty Images; p. 752: NASA; p. 754: Vladyslav Danilin/Fotolia; p. 755: StockLite/Shutterstock; p. 757: BSIP/Newscom; p. 762: Keystone/Getty Images; p. 765: Map from the USGS, U.S. Geological Survey Library; p. 768: 3D4Medical/Science Source

Chapter 24 p. 783: Stock image/Shutterstock; p. 784: Andrew Lambert Photography/Science Source; p. 786: Eric Schrader/Pearson Education; p. 787: otnaydur/Shutterstock; p. 791: National Technology and Engineering Solutions of Sandia, LLC.; p. 795: Cabeza de Marmore/Shutterstock; p. 798: Based on H Heller, M Schaefer, & K Schulten, "Molecular dynamics simulation of a bilayer of 200 lipids in the gel and in the liquid-crystal phases", J. Phys. Chem. p. 97:8343-60, 1993/Pearson Science; p. 800: Luis Molinero Martínez/123rf; iii: Caiimage/Chris Ryan/GettyImages

Chapter 25 p. 812: JGI/Tom Grill/Getty Images; p. 815: Eric Schrader/Pearson Education; p. 817: Stockbyte/Getty Images; p. 818: Dr. David Furness, Keele University/Science Source; p. 820: TFoxFoto/Shutterstock; p. 822: Laurence Gough/Shutterstock; p. 824: Andrew J. Martinez/Science Source; p. 824: Richard Megna /Fundamental Photographs; p. 843: Based on F. Vollrath and D. Edmonds, "Consequences of electrical conductivity in an orb spider's capture web," Naturwissenschaften (100:12, December 2013, pp. 1163–69).

Chapter 26 p. 844: 06photo/Shutterstock; p. 846: Claro Cortes/Reuters; p. 849: John P. Surey; p. 854: Yury Kosourov/Shutterstock; p. 855: Gary Ombler/Getty Images; p. 857: Richard Megna/Fundamental Photographs; p. 860: Mehau Kulyk/Science Source; p. 864: Evgeny Tomeev/Shutterstock; p. 864: Vladimir Arndt/Shutterstock

Chapter 27 p. 878: Michael Flippo/Fotolia; p. 879: harukaze-k/Fotolia; p. 880: Robert Smith/Getty Images; p. 882: LStockStudio/Shutterstock; p. 885: Richard Megna/Fundamental Photographs; p. 885: Richard Megna/Fundamental photographs; p. 888: Richard Megna/Fundamental Photographs; p. 889: NASA; p. 889: Photo courtesy of Berkeley Lab; p. 889: Roman Krochuk/Shutterstock; p. 898: Simon Fraser/Science Source; p. 903: heptagon.circle/Shutterstock; p. 905: Microgen/Shutterstock

Chapter 28 p. 918: Image courtesy of CERN; p. 922: Tristan3D/Shutterstock; p. 925: TaweePhoto/Shutterstock; p. 927: Jeremy Walker/Science Source; p. 929: James Steidl/Fotolia; p. 936: Eric Schrader; p. 941: Kenneth Scarberry

Chapter 29 p. 953: jabejon/Getty Images; p. 954: Simon Fraser/University of Durham/Science Source; p. 960: CTK/Alamy Stock Photo; p. 968: Mino Surkala/Shutterstock; p. 968: Susan & Allan Parker/Alamy Stock Photo; p. 970: NASA; p. 976: Richard Megna/Fundamental Photographs

Chapter 30 p. 988: EschCollection/GettyImages; p. 989: Petr Malyshev/Shutterstock; p. 992: Ron Chapple/Shutterstock; p. 994: Turley/Fotolia; p. 997: Ensuser/Shutterstock; p. 997: NASA

Chapter 31 p. 1018: Roger A. Freedman; p. 1021: Rubberball/Mike Kemp/Getty Images; p. 1028: Lawrence Migdale/Science Source; p. 1028: Mauro Fermariello/Science Source; p. 1036: Katharina Wittfeld/Shutterstock; p. 1037: Roger A. Freedman; p. 1037: Thomas Barrat/Shutterstock; p. 1038: egd/Shutterstock

Chapter 32 p. 1048: JLIImages/Alamy Stock Photo; p. 1049: Albert Einstein in James Clerk Maxwell, A Commemoration Volume (New York: The Macmillan Company, 1931), p. 71.; p. 1049: Bettmann/Getty Images; p. 1050: Jirapol Konroo/Shutterstock; p. 1050: Pablo Hidalgo/123rf; p. 1052: Ted Kinsman/Science Source; p. 1058: John A Davis/Shutterstock; p. 1060: Dudarev Mikhail/Shutterstock; p. 1062: Craig Holmes Premium/Alamy Stock Photo; p. 1063: Alexander Tsiras/Science Source; p. 1065: NASA; p. 1068: grzymkiewicz/Getty Images; p. 1075: "ICNIRP Statement on the 'Guidelines for Limiting Exposure to Time-Varying Electric, Magnetic, and Electromagnetic Fields (up to 300 GHz)," 2009; Health Physics 97(3): 257–258.

Chapter 33 p. 1077: Rozaliya/Getty Images; p. 1078: bradleym/Getty Images; p. 1078: NIH/Science Source; p. 1082: Mead Treadwell/NASA; p. 1082: Passakorn Umpornmaha/Shutterstock; p. 1082: Peter Herring/Image Quest Marine; p. 1082: Roger A. Freedman; p. 1085: Ken Kay /Fundamental Photographs; p. 1087: VEM/Science Source; p. 1089: mtlghma/Fotolia; p. 1091: Dougal Waters/Getty Images; p. 1091: Pictal/Agfotostock; p. 1093: Diane Hirsch /Fundamental Photographs; p. 1096: Nestor Rizhniak/Shutterstock; p. 1097: Cefo design/Shutterstock; p. 1097: Linda Hirst; p. 1097: Peter Aprahamian/Sharples Stress Engineers Ltd./Science Source; p. 1098: lauriek/Getty Images; p. 1098: Roger A. Freedman; p. 1101: Loskutnikov/Shutterstock

Chapter 34 p. 1110: Corbis/VCG/Getty Images; p. 1113: Eric Schrader; p. 1115: NASA; p. 1117: OlegD/Shutterstock; p. 1124: Vaclav Volrab/Shutterstock; p. 1126: Richard Megna/Fundamental Photographs, NYC; p. 1130: Hoby Finn/Getty Images; p. 1130: John Hedgecoe/TopFoto/The Image Works; p. 1136: John Hedgecoe/TopFoto/The Image Works; p. 1137: kak2s/Shutterstock; p. 1137: THK Photo Products; p. 1138: Omikron/Science Source; p. 1139: ethylalkohol/Shutterstock; p. 1142: Jan Zoetekouw/Shutterstock; p. 1144: Jan Hinsch/Science Source; p. 1147: NAOA

Chapter 35 p. 1159: Ilya Andriyanov/Shutterstock; p. 1162: Peter Dazeley/Getty Images; p. 1163: Roger A. Freedman; p. 1164: Pearson Education Inc.; p. 1170: RBFried/Getty Images; p. 1173: Bausch & Lomb Incorporated; p. 1174: Bausch & Lomb Incorporated; p. 1175: Vladimirkarp/Shutterstock; p. 1175: Vukusic, Pete; p. 1176: Chaiwut Siriphithakwong/Shutterstock

Chapter 36 p. 1185: Tomatito/Shutterstock; p. 1186: Hugh D. Young; p. 1186: Ken Kay/Fundamental Photograph; p. 1189: Pearson Education Inc.; p. 1191: Richard Megna/Fundamental Photographs; p. 1193: Roger A. Freedman; p. 1197: Dr. OGA/Shutterstock; p. 1198: National Optical Astronomy Observatory (NOAO); p. 1198: SPL/Science Source; p. 1199: Tek Image/Science Source; p. 1200: Science History Images/Alamy Stock Photo; p. 1202: Omikron/Science Source; p. 1203: Image Courtesy of National Radio Astronomy Observatory.; p. 1204: GIPhotoStock/Science Source; p. 1204: Pearson Education Inc.; p. 1204: Pearson Education, Inc; p. 1205: Michal Ninger/Shutterstock; p. 1207: Paul Silverman/Fundamental Photographs; p. 1208: GIPhotoStock/Science Source; p. 1213: NASA

Chapter 37 p. 1217: Courtesy of Brookhaven National Laboratory; p. 1220: Adimas/Fotolia; p. 1224: John F. Kennedy Space Center/NASA; p. 1227: Eva-Katalin/Getty Images; p. 1228: Jack Fields/Science Source; p. 1233: sirtravelalot/Shutterstock; p. 1236: John F. Kennedy Space Center/NASA; p. 1236: sirtravelalot/Shutterstock; p. 1241: Geng yuhe/Associated Press; p. 1244: Matt Cooper/Shutterstock

Rear Matter A-1: Adapted from the National Institute of Standards and Technology (<http://physics.nist.gov/cuu>); A-7: National Institute of Standards and Technology (<http://physics.nist.gov/cuu>); A-8: NASA (<http://solarsystem.nasa.gov/planets/>).

INDEX

A

- Absolute pressure, 370–371, 581
- Absolute temperature, 509
- Absolute temperature scale, 546, 658–659
- Absolute zero, 546, 659
- Absorption lines, 1197
- Ac adapters, 1037
- Acceleration
 - angular, 276–279, 278t, 305–308
 - around curve, 70
 - average, 40–41, 70–72. *See also* Average acceleration
 - calculating by integration, 53–55
 - centripetal, 82–83, 149
 - centripetal component of, 280–281
 - changing, finding position and velocity for, 53–55
 - circular motion and, 81–83
 - constant, 44–49. *See also* Constant acceleration
 - finding velocity and position from, 53–55
 - fluid resistance and, 146–149
 - free fall and, 2, 50–53, 113–114
 - human tolerance for, 46, 84
 - inertial reference frame and, 106–107, 110
 - instantaneous, 42–44, 70–72. *See also* Instantaneous acceleration
 - linear, 276, 278t
 - mass and, 108–109, 113–114, 116
 - net force and, 107–113
 - Newton's first law and, 103–107
 - Newton's second law and, 107–113
 - parallel component of, 72–74, 84
 - of particle in wave, 472–474
 - perpendicular component of, 72–74, 84
 - of projectile, 74–76
 - radial, 82, 149–150
 - of rocket, 256–259
 - of rolling sphere, 313
 - sign conventions for, 43, 43t
 - in simple harmonic motion, 435–437
 - tangential, 84, 280
 - uniform circular motion and, 81–83
 - units of, 112
 - as vector quantity, 40
 - vs. velocity, 40
 - weight and, 109, 113–114
 - of yo-yo, 310–311, 312
- Acceleration due to gravity, 50, 113–114, 691
- apparent weight and, 138–141, 414–415
- at different latitudes and elevations, 415t
- vs. gravitation, 396
- magnitude of, 397–398
- Acceleration vectors, 34, 69–74, 277
- Accelerators. *See* Particle accelerators
- Access resistance, 1047
- Accretion disk, 418
- Accuracy
 - in measurement, 8
 - vs. precision, 9
- ac source, 1019. *See also* Alternating-current circuits
- Action potentials, 987
- Action–reaction pairs, 115–119
 - gravitational forces as, 396
- Activation energy, 599
- Addition, of vectors, 11–14, 16–17
- Adiabatic, 625
- Adiabatic process, 624
 - Carnot cycle and, 653
 - for ideal gas, 629–631
- Air
 - dielectric strength of, 761, 798
 - as insulator, 562
 - ionization of, 761–762
- Air bags, 238
- Air conditioners, 651
- Air drag, 146–148
- Airplanes
 - banked curves and, 153
 - noise control for, 522
 - sonic boom from, 529, 530–531
 - wing lift and, 381–382
- Air pressure, 368
- Air resistance, projectile motion and, 74–76
- Airy disk, 1203, 1205
- Airy, George, 1203
- Alternating current, 816, 844, 1018–1047
 - applications of, 863–866
 - conversion from dc to, 1047
 - dangers of, 1036
 - measurement of, 1018–1021
 - rectified, 1019–1020
 - rectified average, 1020
 - root-mean-square value of, 1021–1022
- Alternating-current circuits
 - capacitors in, 1024–1026
 - in household systems, 844
 - impedance of, 1027–1029
 - inductors in, 1022–1023, 1026
 - L-R-C* series, 1027–1030
 - phase angle and, 1023, 1028–1029
 - phasors and, 1018–1021
 - power in, 1031–1033
 - resistance and reactance in, 1022–1026
 - resistors in, 1022, 1026
 - resonance in, 1034–1036
 - transformers and, 1036–1039
- Alternators, 959–960, 1018
- Ammeters, 825, 826, 854–855
 - voltmeters and, 856–857, 1021
- Ampere, 686, 814, 926–927
- Ampère, André, 880
- Ampere's law, 930–936. *See also* Maxwell's equations
 - applications of, 933–936
 - displacement current and, 971–972
 - electromagnetic waves and, 1049, 1054, 1059
 - generalization of, 971–972
 - general statement of, 932–933
 - for long straight conductor, 930–932
 - in symmetric situations, 933
- Amplitude
 - displacement, 502, 510–512
 - of electromagnetic waves, 1058
 - in interference, 1166–1167
 - of oscillation, 430
 - of pendulum, 449–451
 - pressure, 503–504, 510
 - in simple harmonic motion, 434–435
 - of sound waves, 502–505
 - of waves, 470
- Analyzers, 1092–1093
- Anesthetic gases, 640
- Angle of incidence, critical, 1086
- Angle of reflection, 1081
- Angle(s)
 - polarizing, 1094
 - radians and, 273, 281
- Angular acceleration, 276–279
 - angular velocity and, 276–277
 - calculation of, 276
 - constant, 277–279, 278t
 - vs. linear acceleration, 278
 - torque and, 305–308
 - as vector, 277
- Angular coordinates, 272–273
- Angular displacement, 273–274
 - torque and, 314–316
- Angular frequency, 430–431
 - of electromagnetic waves, 1058
 - vs. frequency, 434
 - natural, 452
 - period and, 430–431
 - in simple harmonic motion, 433–435
- Angular magnification, 1142–1143, 1146
- Angular momentum, 316
 - axis of symmetry and, 317–318
 - conservation of, 319–321
 - of electrons, 937–938
 - of gyroscopes, 322–324
 - precession and, 322–324
 - rate of change of, 316, 317
 - rotation and, 316–321
 - torque and, 316, 317–318
 - as vector, 317, 322
- Angular simple harmonic motion, 443
- Angular size, 1142
- Angular speed, 274
 - instantaneous, 280
 - precession, 323
 - rate of change of, 279–280
- Angular velocity, 273–277
 - angular acceleration and, 276–277
 - average, 273–274
 - calculation of, 274
 - instantaneous, 274
 - vs. linear velocity, 274
 - rate of change of, 280
 - as vector, 275
- Angular vs. linear kinematics, 279–282
- Antinodal curves, 1162
- Antinodal planes, 1067
- Antinodes, 483
 - displacement, 514–515
 - pressure, 515
- Antiparallel vectors, 11
- Aphelion, 408
- Apparent weight
 - acceleration due to gravity and, 138–141, 414–415
 - earth's rotation and, 414–415
 - magnitude of, 414–415
- Appliances, power distribution systems in, 863–866
- Archimedes's principle, 373
- Arecibo telescope, 1212
- Aristotle, 50
- Associative law, 12
- Astigmatism, 1140–1141
- Aston, Francis, 892
- Astronomical telescopes. *See* Telescopes
- Asymptote, 861
- Atmosphere, 348, 368
- Atmospheric pressure, 348, 368
 - elevation and, 583–584
 - measurement of, 371–372
- Atomic mass, 587, 680
 - measurement of, 892
- Atomic number, 681
- Atomic particles, 680–681. *See also* Electron(s); Neutrons, mass of; Proton(s)
- Atomic structure, 680
- Atom(s)
 - interactions between, 443–445
 - nucleus of, 680
 - structure of. *See* Atomic structure
 - Thomson's model of, 744
- Attenuator chains, 877
- Audible range, 501
- Automobiles
 - gas compression in, 582–586
 - ignition systems in, 997
 - power distribution systems in, 863, 865
 - vertical simple harmonic motion in, 443
- Average acceleration, 40–41, 70–72
- Average acceleration vector, 70–71
- Average angular acceleration, 276
- Average angular velocity, 273–274

I-2 Index

Average density, 367
Average power, 187, 479
Average speed, 38
Average velocity. *See also* Velocity
 vs. average speed, 38
 vs. instantaneous velocity, 37, 67–68
 motion in two or three dimensions and, 67–69
 straight-line motion and, 34–37
Average velocity vector, 67–69
Average x -acceleration, 40
Average x -velocity, 35–37
Avogadro's number, 587
Axis
 elliptical, 408
 optic, 1114
 polarizing, 1091
 semi-major, 408, 409
Axis of rotation, 275
 change in direction of, 275, 280–281
 fixed, 272–273
 moment of inertia for, 305–306
 moving, 308–314
 parallel-axis theorem and, 287–288
 through center of mass, 287–288
Axis of symmetry, angular momentum and, 317–318
 ax - t graphs, 44–45

B

Back emf, 903, 960
Back-of-the-envelope calculations, 10
Bainbridge's mass spectrometer, 892
Balance
 Cavendish (torsion), 397
 spring, 101
Ballistic pendulums, 247
Ballistic seed dispersal, 99
Banked curves, 153
Bar, 368
Bar magnets, 901
Barometers, mercury, 371–372
Baseball, curve ball in, 384
Batteries
 charging, 829
 vs. current source, 824
 energy in, 827
 terminal voltage of, 824
Beams, 1079
Beat frequency, 522–523
Beats, 522–524
Bednorz, Johannes, 818
Bell, Alexander Graham, 512
Bernoulli's equation, 378–383
Bimetallic strip thermometer, 543
Binary star systems, 418, 1251
Binoculars, 1146
Binomial theorem, 444, 445
Biot-Savart law, 922
Birefringence, 1096
Blackbody, 566
Black holes, 415–419
Blood flow, 65, 383
Blood pressure, 371f, 379f
Blue-ray discs, 1204
Body. *See* Human body
Bohr magneton, 937–938
Boiling, 555, 556–557
Boltzmann constant, 590
Boundary conditions, for waves, 481
Bound charges, 799–800
Bourdon pressure gauge, 372
Bouyant force, 373–374
Bragg condition, 1201
Bragg, Laurence, 1201
Bragg reflection, 1202, 1216
Bragg, William, 1201
Brahe, Tycho, 407
Breaking stress, 351–352
Brewster, David, 1094
Brewster's law, 1094

Bridge circuits, 849–854
Bright fringes, 1189
British system, 5–6, 5t. *See also* Units of measure
British thermal unit (Btu), 552
Brittle material, 351
Bulk modulus, 349
Bulk strain, 348–349
Bulk stress, 348–349
Buoyancy, 373–375

C

Cables, winding/unwinding, 285–286, 307–308
Calcite, birefringence in, 1096
Calculations
 calorimetry, 558–560
 units of measure in. *See* Units of measure
Calorie (cal), 552
Calorimetry calculations, 558–560
Calorimetry, phase changes and, 555–560
Cameras, 1135–1138
 flash unit of, 791
 focusing, 1135–1136
Cancer, 757f, 941f
Canine resonance, 453
Capacitance, 781, 782
 vs. amount of charge or voltage, 796
 calculation of, 782–784
 vs. coulombs, 782
 dielectrics and, 793–803
 equivalence, 787
 units for, 782
Capacitive reactance, 1025–1026
Capacitors, 781–811
 in ac circuits, 1024–1026, 1032
 applications of, 781, 791
 capacitance of, 782. *See also* Capacitance
 capacitive reactance of, 1025–1026
 charge storage in, 790–793
 charging, 859–860, 971–972
 cylindrical, 785–786
 dielectrics in, 793–803
 discharging, 861–862
 electric-field energy and, 791–792
 electrolytic double-layer, 796
 energy storage in, 790–793
 in networks, 789
 in parallel, 787–788, 846–847
 parallel-plate, 783, 794
 vs. resistors, 846
 in series, 786–787, 846–847
 spherical, 785
 symbols for, 782
 in vacuum, 782–784, 791
Carbon dating, 156
Carbon dioxide, greenhouse effect and, 567
Carnot cycle, 653–659
 efficiency of, 657–658
 entropy and, 662–663
 of heat engine, 653–659
 for ideal gas, 654–655
 Kelvin temperature scale and, 658–659
 of refrigerator, 657
 reversibility of, 657–658
 second law of thermodynamics and, 657–658
Cars. *See* Automobiles
Cathode-ray tubes, for magnetic field measurement, 882–883
Cavendish balance, 397
Cavendish, Henry, 397
Celestial mechanics, 395
Cell membrane, 768, 798f, 811
Cell polarity, 1047
Celsius temperature scale, 543, 546
Center of curvature, 1114
Center of gravity, 338–341
Center of mass, 253–256
 center of gravity and, 338–341
 combined rotational-translational motion and, 308–313
 external forces and, 255–256
 motion of, 254–256
 planetary motion and, 410–411
 torque and, 306
Centigrade scale, 543
Centimeter, 5t, 112
Central force, 409
Centrifugal force, 150
Centripetal acceleration, 82–83, 149
Centripetal component of acceleration, 280–281
Čerenkov radiation, 1249–1250
cgs metric system, 112
Charge distribution, 695
 electric fields and, 718–721, 727–728, 738t.
 See also Gauss's law
 static, 752
Charged particles, motion in magnetic fields, 888–893
Charging by induction, 683–684
 polarization and, 684
Chemical bonds, 682, 701
Chemical reactions. *See* Reaction(s)
Chokes, 992–993
Chromatic resolving power, 1198–1200, 1204
Circle
 circumference of, 273
 reference, 433
Circuit breakers, 864
Circuit diagrams, 824–825
Circuit(s), 812
 alternating-current. *See* Alternating-current circuits
 bridge, 849–854
 complete, 815, 822
 critically damped, 1007
 diagrams of, 824–825
 direct-current. *See* Direct-current circuits
 electromotive force and, 822–827
 energy in, 828–832
 incomplete, 822
 inductors in, 992–995. *See also* Inductance
 junctions in, 849–850
 Kirchhoff's rules for, 849–854
 L-*C*, 1002–1006
 loops in, 849–850
 L-*R*-*C* series, 1007–1008, 1027–1030
 open, 864
 oscillating, 1002–1006
 overdamped, 1007–1008
 overloaded, 864–865
 potential changes around, 827
 power in, 828–832
 R-*C*, 858–863
 relaxation times of, 860–861
 R-*L*, 998–1002
 self-inductance and, 992–995
 short, 826–827, 864–865
 time constants for, 860–861, 1000
 underdamped, 450, 1007–1008
Circular apertures, diffraction and, 1203–1205
Circular motion, 81–84, 432–434
 acceleration and, 81–83
 dynamics of, 149–154
 net force and, 108
 nonuniform, 84, 154
 vs. projectile motion, 83
 uniform, 81–83. *See also* Uniform circular motion
Circular orbits, 405–406, 409
Circular polarization, 1095–1096
Classical mechanics, 100
Clausius statement, 652
Climate change, 566–567
Closed orbits, 404
Closed surface, electric flux through, 719
Coaxial cable, 936
Coefficient of kinetic friction, 142
Coefficient of linear expansion, 547–558
Coefficient of performance, 649
Coefficient of resistivity, temperature, 818, 820
Coefficient of rolling friction, 146
Coefficient of static friction, 142–143
Coefficient of volume expansion, 548–549
Coherent waves, 1160

- Cohesion, in trees, 367f
- Coils
- inductance of. *See* Inductance
 - magnetic fields of, 927–930
 - magnetic torque on, 899–900
 - Tesla, 990–991
- Cold reservoirs, 644–645
- Collisions, 245–252
- classification of, 248–249
 - conservation of momentum and, 245–249
 - elastic, 245–246, 249–252
 - inelastic, 245–249
 - kinetic energy in, 245–246
- Combined translation and rotation, 308–313
- Combustion, heat of, 558
- Comet Halley, orbit of, 410
- Commutative law, 12
- Commutators, 902–903, 960
- Complete circuits, 815, 822
- Completely inelastic collisions, 245–249
- Components, of vectors, 14–18, 20–21, 23–24, 102–103
- Compound microscope, 1143–1145
- Compressibility, fluid, 349, 376
- Compression, 347
- fluid density and, 468
- Compression ratio, 647
- Compressive strain, 347
- Compressive stress, 347
- Concave mirrors, 1114–1118
- Concentration of particles, in current, 814
- Condensation, 556–557
- Condensed matter. *See* Liquid(s); Solids
- Condenser microphones, 783
- Conduction, 561–564
- Conduction current, 971
- electrical, 817
 - microscopic model of, 832–834
 - thermal, 561–562, 817
- Conductivity, 817
- Conductors, 682–683
- in capacitors, 782
 - conductivity of, 817
 - current density in, 815–816
 - current flow in, 813–814
 - diodes of, 821
 - electric charge on, 729, 734–737
 - electric fields at, 692, 736–737
 - electron motion in, 813
 - equipotential surfaces and, 766–767
 - holes in, 904
 - interaction force between parallel, 926–927
 - magnetic fields of, 923–926
 - magnetic force of, 926–927
 - magnetic forces on, 893–896
 - metallic, 832–834
 - nonohmic (nonlinear), 817
 - ohmic (linear), 817
 - particle concentration in, 814
 - resistance of, 819–822, 824, 827
 - resistivity of, 816–819
 - semiconductors, 817, 821, 904
 - superconductors, 818, 964
 - thermal, 542
- Conservation laws, universal, 681
- Conservation of angular momentum
- net torque and, 319–321
 - of planets, 408–409
- Conservation of electric charge, 681–682
- Kirchhoff's junction rule and, 850
- Conservation of electrostatic force, 850
- Conservation of energy, 171, 218
- with electric force, 751
 - in simple harmonic motion, 438–441
 - total mechanical energy, 203–204
- Conservation of mass and energy, 1241
- Conservation of mechanical energy, 748
- Conservation of momentum, 241–245, 1237
- collisions and, 245–249
 - conservation of components and, 242
- Conservative forces, 215–219
- elastic collisions and, 245
 - work done by, 748
- Consonance, 524
- Constant acceleration, 44–49
- due to gravity, 50–53
 - equations of motion with, 47
 - of freely falling bodies, 50–53
 - of projectile, 74
- Constant angular acceleration, 278t
- rotation with, 277–279
- Constant forces, 172–173
- Constant linear acceleration, 278t
- Constant-pressure process, 624, 626–628
- Constant-temperature process, 624–625
- Constant torque, 314–315
- Constant-volume gas thermometers, 545
- Constant-volume process, 624
- Constructive interference, 484, 520–521, 1160–1162, 1164–1165
- in holography, 1206–1207
 - in x-ray diffraction, 1202
- Contact force, 101, 116, 141
- Contact lenses, 1139–1141
- Continuity equation, 376–377
- Convection, 564
- Conventional current, 814
- Converging lenses, 1127–1129
- Converging mirrors, 1114–1118
- Convex mirrors, 1118–1120
- Cooling, evaporative, 558
- Coordinates
- angular, 272–273
 - spacetime, 1232
- Copernicus, Nicolaus, 407
- Cornea, 1138, 1158
- Corona discharge, 761–762
- Correspondence principle, 1243
- Coulomb, 686–687
- vs. capacitance, 782
- Coulomb, Charles Augustin de, 685
- Coulomb's law, 586, 685–690
- electric force and, 678, 685–690
 - Gauss's law and, 725
 - proportionality constant in, 686–687
 - statement of, 685
 - superposition of forces and, 687
- Critical angle, 1086
- Critical damping, 450
- Critical fields, 975
- Critically damped circuits, 1007
- Critical point, 600, 601
- Critical temperature, 585, 975
- Cross product, 22–24
- Cryopreservation, 578
- Crystal lattice, 586
- Crystalline lens, 1139
- Cube, electric flux through, 724
- Curie constant, 939
- Curie, Pierre, 939
- Curie's law, 939
- Current, 812, 813
- alternating, 816
 - capacitor, 1024–1026
 - in circuits, 822–831
 - concentration of particles in, 814
 - conduction, 971
 - conventional, 814
 - vs. current density, 815
 - direct, 816, 844. *See also* Direct-current circuits
 - direction of, 813–814, 819, 894
 - displacement, 970–974
 - drift velocity and, 813, 814–816
 - eddy, 969–970, 1038
 - electric charge in, 813–814
 - electric field and, 813–814
 - electromotive force and. *See* Inductance
 - electron motion in, 813
 - full-wave rectifier, 1019–1020
 - heat, 561
 - induced, 954–955, 962–964
 - inductance and, 988. *See also* Inductance
 - inductor, 1022–1023, 1026
 - Kirchhoff's rules for, 849–854
 - lagging, 1033
 - measurement of, 854–858
 - notation for, 859
 - Ohm's law and, 816, 819–821
 - rectified average, 1020
 - resistance and, 819–822
 - resistor, 1022, 1026
 - root-mean-square, 1021–1022
 - sinusoidal, 1018–1021. *See also* Alternating current
 - time-varying, 859
 - units of, 686, 814
 - "using up," 824
 - voltage and, 819–822
- Current amplitude, 1019
- Current-carrying conductor, magnetic forces on, 893–896
- Current density, 815
- vs. current, 815
 - resistivity and, 816
 - vector, 815
- Current flow, direction of, 813–814
- Current loops. *See also* Magnetic dipoles
- force and torque on, 897–902
 - magnetic fields of, 927–930
 - magnetic moment of, 898, 929
 - in magnetization, 937–938
- Curve ball, 384
- Curve(s)
- acceleration around, 70
 - antinodal, 1162
 - banked, 153
 - gravitational potential energy and, 206–210
 - magnetization, 940
 - motion along, 185–187
 - nodal, 1162
 - resonance, 518–519, 1035
 - response, 1035
 - work–energy theorem for, 181
- Cycles, 430–431
- Cyclic process, in heat engines, 644
- Cyclotron, 888, 910
- Cyclotron frequency, 888
- Cylinders, moment of inertia of, 285t, 289–290
- Cystic fibrosis, 685f
- D**
- Damped harmonic motion, 1007
- Damped oscillations, 449–451
- energy in, 451
 - with periodic driving force, 452
- Damping, 450
- critical, 450
- Dark fringes, 1187–1189
- d'Arsonval galvanometers, 854, 857, 899, 1019–1020
- dc circuits. *See* Direct-current circuits
- Deceleration, 43
- Decibel scale, 512–513
- Deformation
- Hooke's law and, 345, 349, 351
 - plastic, 351–352
 - reversible, 351
 - stress and strain and, 345–351
- Degrees, 543, 545
- Degrees of freedom, 595
- Density, 366–368
- average, 367
 - of common substances, 367t
 - current, 815
 - displacement current, 972
 - of earth, 401
 - energy, 791, 1061–1062
 - fluid, 373–374, 468
 - linear charge, 695
 - linear mass, 474

I-4 Index

Density (*continued*)

- magnetic energy, 996–997
- magnetic flux, 887
- mass-to-volume ratio and, 367
- measurement of, 366
- vs. pressure, 581
- surface charge, 695
- volume charge, 695

Depth, fluid pressure and, 369–370

Derivatives

- instantaneous velocity and, 37
- partial, 220–221, 472–473

Destructive interference, 484, 520, 522, 1160–1162, 1164–1165

Dewar flask, 566

Dewar, James, 566

Diamagnetism, 939–940, 975–976

Diamond, 1087, 1088

Diatomic molecules, 595

Dichroism, 1091

Dielectric breakdown, 793–794, 798

Dielectric constant, 794

Dielectric function, 1061

Dielectrics, 781, 793–803

- electromagnetic waves in, 1060–1061
- Gauss's law and, 801
- permittivity of, 795–796
- polarization of, 795–796, 799–800

Dielectric strength, 798

- of air, 761, 798

Diesel cycle, 648–649

Diesel engines, 648–649

- adiabatic compression in, 631

Difference tone, 524

Differential principle, 238

Diffraction, 1185–1216

- bright fringes in, 1189
- with circular apertures, 1203–1205
- dark fringes in, 1187–1189
- Fraunhofer, 1186–1187
- Fresnel, 1186–1187
- holography and, 1206–1207
- Huygens's principle and, 1186–1187
- image formation and, 1204
- intensity in, 1190–1194
- vs. interference, 1187
- multiple-slit, 1194–1196
- near-field, 1186
- resolving power of, 1204
- single-slit, 1187–1193
- of sound, 1185, 1193
- x-ray, 1200–1203

Diffraction gratings, 1196–1200

Diffraction minima, 1194

Diffuse reflection, 1080, 1111

Digital multimeters, 857

Dimagnetism, 939–940

Dimensionally consistent units, 6

Diode(s), 821

Diopters, 1141

Dipoles

- electric, 696, 701–705, 799–800, 899. *See also* Electric dipole(s)
- magnetic, 898

Direct current, 816

- conversion to ac, 1047
- dangers of, 1036

Direct-current circuits, 844–877

- in automobiles, 863
- Kirchhoff's rules for, 849–854
- measuring instruments for, 854–858
- in power distribution systems, 863–866
- R-C*, 858–863
- resistors in series and parallel in, 844–847

Direct-current generators, 960–962

Direct-current motors, 902–904

Direction, 10

- of force, 10
- of vectors, 10, 14–16
- of waves, 471, 473

Disk, electric flux through, 723

Dispersion, 1083, 1088–1090

Displacement, 10–11, 17. *See also* Position

- angular, 273
- average velocity and, 34–37
- in oscillation, 429
- in simple harmonic motion, 432, 435–437
- superposition principle and, 482
- wave pulse and, 481–482
- work and, 172–175, 176–178

Displacement amplitude, 502

- sound intensity and, 510–512

Displacement current, 970–974

Displacement nodes/antinodes, 514–515

Dissipative forces, 216

Dissonance, 524

Distance

- image, 1112
- object, 1111
- relativity of, 1227–1231
- units of, 112

Distribution function, 598

Diverging lenses, 1129

DNA

- base pairing in, 710, 773
- x-ray diffraction of, 1199, 1202

DNA Spring, 234

Doppler, Christian, 524

Doppler effect

- for electromagnetic waves, 528–529, 1235–1236
- for sound waves, 524–529, 1235–1236

Doppler radar, 529

Doppler shift, 1236

Dot product, 19–21

Drag, 146–149

Drift velocity, 813

- current and, 814–816
- Hall effect and, 904–905

Driven oscillation, 451–453

Driving force, 451–453

Ductile material, 351

Dulong and Petit, rule of, 555, 596–597

DVD players and DVDs, 1197, 1204

Dynamics, 34, 100. *See also* Force(s); Mass; Motion

- of circular motion, 149–154
- fluid, 366, 378–381
- of particles, 134–141
- of rotational motion, 302–336

Dyne, 112

E

Ear, human, 502, 504, 519

Earth

- density of, 401
- magnetic fields of, 878, 879, 882
- rotation of, 414–415
- surface temperature of, 567
- weight of objects and, 399–400

Eccentricity, orbital, 408

Eddy currents, 969–970

- in transformers, 1038

Edison, Thomas, 1018

Efflux, speed of, 380–381

Einstein, Albert, 87, 415, 1217. *See also* Relativity

Elastic, 210, 346

Elastic collisions

- momentum conservation and, 245–246
- one-dimensional, 249–250
- relative velocity and, 250–251
- two-dimensional, 252

Elastic deformations, Hooke's law and, 345, 349, 350, 351

Elastic hysteresis, 351

Elasticity, 351–352

Elastic limit, 351

Elastic modulus, 345

- bulk, 348–349
- shear, 350
- Young's, 346–347

Elastic potential energy, 210–215, 219–220

- gravitational potential energy and, 211, 212

Electrical conductivity, 817

Electrical energy, 188

- units of, 188

Electrical oscillation, 1002–1006

- in *L-C* circuits, 1004–1005

Electrical power, 188. *See also* Power

Electric charge, 679–682

- attraction and repulsion and, 679
- in bees, 716
- bound, 799–800
- in capacitors, 782. *See also* Capacitors
- in closed surface, 719
- on conductors, 729, 734–737
- conservation of, 681–682, 850
- density of, 695
- distribution of. *See* Charge distribution
- electric dipole and, 701–705
- electric field and, 718–721, 727–728. *See also* Gauss's law
- flux and, 718–721. *See also* Electric flux
- free, 800
- induced, 682–684, 799–800
- magnetic force on, 881–882
- vs. magnetic poles, 879–880
- magnitude of, 685
- negative, 679
- in nerve cells, 734, 877
- notation for, 859
- point. *See* Point charges
- positive, 679
- quantized, 682
- structure of matter and, 680–681
- superposition of forces and, 687
- time-varying, 859
- typical values for, 687

Electric circuits. *See* Circuit(s)

Electric constants

- vs. dielectric constant, 794
- fundamental, 686–687

Electric current. *See* Current

Electric dipole moment, 702

Electric dipole(s), 696, 701–705, 799–800

- field of, 696, 704–705
- force on, 702–703, 704
- potential energy of, 703–704
- torque on, 702–703, 704, 899

Electric-field energy, 791

Electric field lines, 700–701

- electromagnetic waves and, 1050
- equipotential surfaces and, 765–766
- point charges and, 727–728, 1050

Electric field(s), 690–699

- calculation of, 694–699, 768
- of capacitor, 782
- of charged conducting sphere, 730
- charge distribution and, 718–721, 727–728, 738t. *See also* Gauss's law
- at conductors, 692, 736–737
- current and, 813–814
- direction of, 692, 700
- of earth, 737
- electric dipole and, 696, 701–705
- electric forces and, 690–694
- electric potential and, 754, 755–756, 767–769
- of electromagnetic waves, 1050, 1058, 1066–1067
- energy storage in, 790–793
- vs. flows, 699
- flux of, 718–725. *See also* Electric flux
- Gauss's law for, 718–746, 930, 1049, 1052–1053
- of hollow sphere, 733
- in hydrogen bomb, 739–740
- induced, 967–969
- line integral of, 932–933
- magnetic fields and, 971–972

- magnitude of, 692, 700
 - molecular orientation in, 799–800
 - nodal/antinodal planes of, 1067
 - nonelectrostatic, 955, 968–969
 - of parallel conducting plates, 732, 783
 - of plane sheet of charge, 731
 - of point charge, 691–693
 - resistivity and, 816–819
 - superposition of, 695
 - of symmetric charge distributions, 738
 - test charge for, 690–691
 - uniform, 700
 - of uniform line charge, 731
 - of uniformly charged sphere, 732–733
 - units for, 691, 756
 - in vacuum, 791
 - as vector quantity, 768
 - work done by, 748–753. *See also* Electric potential energy
- Electric flux
 - calculation of, 721–725
 - charge and, 718–721
 - enclosed charge and, 719–721
 - fluid-flow analogy for, 721–722
 - Gauss's law for, 718–746
 - of nonuniform electric field, 723
 - outward vs. inward, 719
 - of uniform electric field, 722
- Electric force, 154–156
 - conservation of energy with, 751
 - Coulomb's law and, 678, 685–690
 - direction of, 685
 - electric field and, 690–694
 - vs. electric potential energy, 750
 - vs. gravitational force, 686, 687
 - potential energy and, 220
 - on uncharged particles, 684
 - units of, 686–687
 - vector addition of, 689
 - work done by, 748–753, 754
- Electric lines
 - hot side, 863, 864
 - household, 863–866
 - neutral side of, 863, 864
- Electric motors
 - direct-current, 902–904
 - magnetic force in, 893
- Electric potential, 747–780. *See also* Potential
 - calculation of, 755, 760–764
 - electric circuits and, 828–832
 - electric field and, 754, 755–756, 767–769
 - electric potential energy and, 747–753
 - equipotential surfaces and, 764–767
 - field lines and, 765–766
 - maximum, 761–762
 - of a with respect to b, 754
 - as scalar quantity, 754, 768
 - units of, 754, 756
 - work done by, 754
- Electric potential energy, 747–753
 - alternative concepts of, 752–753
 - in capacitors, 782. *See also* Capacitors
 - electric-field energy and, 791
 - vs. electric force, 750
 - electric potential and, 756
 - of several point charges, 752
 - of two point charges, 749–750
 - in uniform field, 748–749
- Electric toothbrush, 989f
- Electrocardiography, 755f
- Electrodynamics, quantum, 1078
- Electrolytic double-layer capacitors, 796
- Electromagnetic energy flow, 1062–1064
- Electromagnetic induction, 953–987
 - changing magnetic flux and, 955
 - eddy currents and, 969–970
 - experiments in, 954–955
 - Faraday's law and, 953, 955–962
 - induced electric fields and, 967–969
 - Lenz's law and, 962–964
- Maxwell's equations and, 953, 972–974
 - motional electromotive force and, 965–967
 - superconductors and, 964, 975–976
- Electromagnetic interaction, 154–156
- Electromagnetic momentum flow, 1064–1065
- Electromagnetic radiation, 565–567, 1050–1051.
 - See also* Electromagnetic wave(s)
- Electromagnetic spectrum, 1051–1052
- Electromagnetic wave equation, 1055–1057
- Electromagnetic wave(s), 1048–1076
 - amplitude of, 1058–1059
 - angular frequency of, 1058
 - applications of, 1051–1052
 - in dielectrics, 1060–1061
 - direction of, 1055
 - Doppler effect for, 528–529, 1235–1236
 - electric fields of, 1050, 1058, 1066–1067
 - energy in, 1062–1064
 - frequency of, 1058
 - generation of, 1050–1051
 - intensity of, 1063
 - magnetic fields of, 1058, 1066–1067
 - magnitude of, 1055
 - in matter, 1060–1061
 - Maxwell's equations and, 1048–1052
 - momentum of, 1064–1065
 - plane, 1052–1057
 - polarization of, 1055. *See also* Polarization
 - power in, 479
 - Poynting vector of, 1062–1064
 - properties of, 1055
 - radiation pressure and, 1065
 - reflected, 1066–1069
 - right-hand rule for, 1054
 - and safety, 1075
 - sinusoidal, 1057–1061
 - speed of, 1054, 1057–1058, 1068
 - standing, 1050, 1066–1069
 - superposition of, 1066
 - transverse, 1053, 1057
 - units for, 1050
 - wave functions for, 1058, 1059
 - wave number for, 1058
- Electromagnetism, 678, 880
- Electrometers, 794
- Electromotive force (emf), 822–827
 - back, 903
 - current and. *See* Inductance
 - of electric motor, 903
 - Hall, 904–905
 - induced, 903, 954–955
 - measurement of, 858
 - motional, 965–967
 - self-induced, 992–995, 1022–1024
 - sinusoidal alternating, 1018
 - source of, 823, 825, 829–830
 - theory of relativity and, 1218
 - in transformers, 1037
- Electromyography, 855f
- Electron(s)
 - angular momentum of, 937–938
 - charge of, 680–681, 686–687. *See also* Electric charge
 - charge-to-mass ratio for, 891–892
 - in magnetic fields, 888–889
 - mass of, 680, 892
 - orbital motion of, 813
 - spin of, 938
- Electron spin, 938
- Electron volts, 756–757
- Electrostatic force
 - conservation of, 850
 - line integral for, 747, 932–933
- Electrostatic precipitators, 777
- Electrostatics, 679
- Electrostatic shielding, 735–736
- Electrostatics problems, 687
- Electroweak interactions, 156
- Elements, isotopes of, 892
- Elevation, atmospheric pressure and, 583–584
- Elliptical orbits, 408–409
- Elliptical polarization, 1095–1096
- emf. *See* Electromotive force (emf)
- Emissivity, 565
- Endoscopes, 1087
- Energy activation, 599
 - conservation of, 171, 218
 - conversion of, 218
 - costs of, 865
 - in damped oscillations, 451
 - electrical, 188
 - in electric circuits, 828–832
 - electric-field, 791
 - in electromagnetic waves, 1062–1064
 - equipartition of, 595
 - internal. *See* Internal energy
 - kinetic. *See* Kinetic energy
 - in *L-C* circuits, 1002–1006
 - of locomotion, 200
 - magnetic-field, 995–998
 - molecular, 585–587
 - potential, 201–234. *See also* Potential energy
 - power and, 865
 - relativistic kinetic, 1239–1240
 - rest, 1240–1241
 - in simple harmonic motion, 438–441
 - total, 171, 1240
 - units for, 756–757
 - in wave motion, 478–481
 - work and, 171–187. *See also* Work
- Energy density, 791, 1061–1062
- Energy diagrams, 222–223
- Energy-flow diagrams
 - for heat engines, 645
 - for refrigerators, 649
- Energy flow, electromagnetic, 1062–1064
- Energy-mass conservation law, 1241
- Energy of locomotion, 200
- Energy storage, in capacitors, 790–793. *See also* Capacitors
- Energy transfer
 - heat and, 552–555
 - rates of, 561
- Engines. *See* Heat engines
- Engine statement, of second law of thermodynamics, 651
- Entropy, 659–665
 - calculation of, 666–667
 - Carnot cycle and, 662–663
 - of coiled polymers, 668
 - in cyclic processes, 663–664
 - internal energy and, 661
 - in irreversible processes, 664
 - in living organisms, 664
 - microscopic interpretation of, 665–668
 - randomness and, 642, 660
 - reversibility of, 661–662, 664–665
 - second law of thermodynamics and, 664–665, 668
- Equation(s)
 - Bernoulli's, 378–381
 - continuity, 376–377
 - electromagnetic wave, 1055–1057
 - ideal-gas, 580–584
 - lensmaker's, 1129–1130
 - Maxwell's. *See* Maxwell's equations
 - of motion with constant acceleration, 47
 - of parabola, 76
 - of simple harmonic motion, 432–434
 - of state, 579–585, 601
 - van der Waals, 584
 - wave. *See* Wave equation
- Equilibrium, 105, 337–365
 - center of gravity and, 338–341
 - first condition of, 336–337
 - for mechanical waves, 465
 - Newton's first law and, 105
 - phase, 556, 600
 - potential energy and, 222–223
 - rigid-body, 338, 341–345
 - rotation and, 338

- Equilibrium (*continued*)
 second condition of, 338
 stable, 222
 static, 338
 thermal, 541–543
 torque and, 338
 unstable, 222
 weight and, 338–341
 weight lifting and, 344
- Equilibrium processes, 643
- Equipartition, of energy, 595
- Equipotential surfaces, 764–767
 conductors and, 766–767
 vs. Gaussian surface, 766–767
- Equipotential volume, 766–767
- Equivalent capacitance, 787
- Equivalent resistance, 845, 846
- Erect image, 1113
- Errors, measurement, 8
- Escape speed, 403, 406, 416
- Estimates, order-of-magnitude, 10
- Ether, light propagation through, 1176, 1218
- Evaporation, 558
- Event horizons, 417, 418
- Event, in frame of reference, 1220
- Exoplanets, 428
- Experiments
 theories and, 2
 thought, 1220–1222
- Extended objects, 1111
 gravitational potential energy for, 287, 310
 image formation by lenses and, 1127–1129
 image formation by mirrors and, 1112–1113
- External forces, 103–104, 241
 center-of-mass motion and, 255–256
 torque and, 306
- Extracorporeal shock wave lithotripsy, 530
- Eye, 1138–1142
 index of refraction of, 1139
 resolution of, 1214
 structure of, 1138–1139
- Eyeglasses, corrective, 1139–1142
- Eyepiece, microscope, 1144
- F**
- Fahrenheit scale, 544
- Farad, 782, 784
- Faraday cage, 735–736
- Faraday disk dynamo, 966–967
- Faraday, Michael, 700, 782, 880
- Faraday number, 811
- Faraday's icepail experiment, 735
- Faraday's law of induction, 953, 955–962, 966–967
 electromagnetic waves and, 1049, 1053, 1059. *See also* Maxwell's equations
- Far point, of eye, 1139
- Farsightedness, 1139–1142
- Fat measurement, 1028f
- Fermi, Enrico, 10
- Ferromagnetism, 940–941, 994
- Fiber optics, 1087
- Field, 399
- Field lines. *See* Electric field lines; Magnetic field lines
- Field point, 691–692, 918–919
- Filters
 high-pass, 1025
 low-pass, 1024
 polarizing, 1090, 1091–1093, 1094
- First condition for equilibrium, 337–338
- First law of thermodynamics, 613–641
 cyclic processes and, 620–623
 internal energy and, 618–623
 isolated systems and, 620–623
- Fission, nuclear, 1241
- Fixed-axis rotation, 272–273, 277
- Flash unit, of camera, 791
- Flow line, 376. *See also* Fluid flow
- Flow rate, 376–377
- Flow tubes, 376
- Fluid density, 366–368
 buoyancy and, 373–375
 compression and, 468
 measurement of, 374
 rarefaction and, 468
- Fluid dynamics, 366
 viscosity and, 382–384
- Fluid flow, 376–378
 Bernoulli's equation and, 378–381
 continuity equation and, 376–377
 laminar, 376, 382–383
 measurement of, 381
 pressure and, 378–381
 rate of, 376–377
 speed of, 378–381
 steady, 376
 turbulent, 376, 383
- Fluid mechanics, 366–394
 Bernoulli's equation and, 378–381
 buoyancy and, 373–375
 density and, 366–368, 373–374
 fluid flow and, 376–378
 pressure and, 368–373
 surface tension and, 375
- Fluid pressure, 348, 368–373
 depth and, 369–370
 measurement of, 371–372
 Pascal's law and, 368, 369–370
- Fluid resistance, 146–148
- Fluid(s), 366
 compressibility of, 349
 ideal, 376
 incompressibility of, 376, 378
 motion of, 376–377
 speed of sound in, 506–507
 viscous, 382–384
- Fluid statics, 366
- Fluorescent light, 994, 1078
- Flux. *See* Electric flux; Magnetic flux
- f -number, 1136–1137
- Focal length
 of camera lens, 1136
 of microscope lens, 1144–1145
 of mirror, 1115–1116
 of telescope lens, 1145–1146
 of thin lens, 1127, 1129–1131
- Focal point
 of microscope lens, 1144–1145
 of mirror, 1115–1116
 of thin lens, 1127
 virtual, 1119
- Foci (focus), 408
- Fog, 558
- Food, energy value of, 558
- Force constant, 182
- Forced convection, 564
- Forced oscillations, 451–453, 518
- Force fields, 399
- Force per unit area, 347
- Force(s), 101–103
 acting at a distance, 399
 action–reaction pairs and, 115–119
 buoyant, 373
 central, 409
 centrifugal, 150
 components of, 220–222
 conservative, 215–219
 constant, 172–173
 contact, 101, 116, 141
 on dancer's body, 127
 direction of, 10. *See also* Vector(s)
 dissipative, 216
 driving, 451–453
 electric. *See* Electric force
 electromotive, 822–827
 electrostatic, 850
 equal and opposite, 115–119
 external, 103–104, 241
- fluid resistance, 146–149
- free-body diagrams for, 119–120, 134
- friction, 101, 104, 141–149. *See also* Friction
- fundamental, 154–156
 gravitational. *See* Gravitation
 as interaction, 100–103, 926–927
 intermolecular, 585–587
 internal, 241, 306
 line of action of, 303
 long-range, 101, 116
 magnetic, 154–156, 881–882
 magnitude of, 10, 14–16, 101
 mass and, 108–109
 measurement of, 101, 112
 net. *See* Net force
 nonconservative, 215–219
 normal, 101, 141
 particle interactions and, 154–156
 periodic driving, 451–453
 potential energy and, 219–222
 power and, 187–189
 vs. pressure, 348, 368
 properties of, 101
 restoring, 430
 strong interactions, 155
 strong nuclear, 155, 680
 superposition of, 101–103, 398, 687
 tension, 101. *See also* Tension
 tidal, 418
 torque of, 302–308
 units of, 5, 101
 weak interactions, 155–156
 weight as, 101
- Force vectors, 10, 101–103
- Fossil fuels, climate change and, 567
- Fourier analysis, 505
- Fourier series, 488
- Fractional error, 8
- Fracture, 351–352
- Frame of reference, 85
 event in, 1220
 inertial, 106–107, 110, 1217, 1218, 1219–1220
 relative velocity and, 85
 simultaneity and, 1220–1222
- Franklin, Benjamin, 679
- Franklin, Rosalind, 1202
- Fraunhofer diffraction, 1186–1187
- Fraunhofer, Joseph von, 1186
- Free-body diagrams, 119–120, 134
- Free charges, 800
- Free convection, 564
- Free expansion, 618, 625
- Free fall
 acceleration and, 2, 50–53, 113–114
 fluid resistance and, 146–149
- Frequency, 430
 angular, 430–431
 vs. angular frequency, 434
 beat, 522–524
 fundamental, 487
 normal-mode, 486–490
 period and, 430–431
 of standing waves, 487
- Fresnel, Augustin Jean, 1186
- Fresnel diffraction, 1186–1187
- Friction, 104, 141–149
 and climbing shoes, 170
 coefficients of, 142–143
 fluid resistance and, 146–149
 kinetic, 141–143, 216
 magnitude of, 142
 rolling, 146, 313–314
 static, 142–143
 stick-slip phenomenon and, 143
- Friction force, 101, 104, 141–149
- Full-wave rectifier current, 1019–1020
- Fundamental electric constants, 686–687
- Fundamental forces, 154–156
- Fundamental frequency, 487

- Fuses, 864
Fusion. *See* Nuclear fusion
- G**
Galilean coordinate transformation, 1219–1220
Galilean velocity transformation, 87, 1220
Galileo Galilei, 2, 50, 100, 1077
Gallium, melting temperature of, 557
Galvanometers, d'Arsonval, 854, 857, 899, 1019–1020
Gas
 bulk modulus of, 349
 density and, 366–367, 370
 heat capacities of, 594–596
 ideal. *See* Ideal gas
 intermolecular forces in, 585–587
 isotherms and, 584–585
 kinetic energy of, 594
 mass of, 581–582
 molecular speed in, 597–599
 molecules in, 585–587
 monatomic, 595
 polyatomic, 595
 pV-diagrams for, 584–585
 sound waves in, 508–509
 thermal conductivities of, 561t
 volume of, 582
Gas constant, 581–582
Gaseous phase, 555–556
Gasoline engines, 647–649
Gas pressure
 molecular collisions and, 588–589
 molecular kinetic energy and, 589–590
 temperature and, 546
Gas thermometers, 545–546, 582, 659
Gauge pressure, 370–371
Gauges, pressure, 371–372
Gauss, 882
Gauss, Carl Friedrich, 725
Gaussian surface, 727–728
 vs. equipotential surface, 766–767
Gauss's law, 718–746
 applications of, 729–733
 charge and electric flux and, 718–721
 conductors with cavities and, 734, 766–767
 dielectrics and, 801
 for electric fields, 930, 1049, 1052–1053. *See also* Maxwell's equations
 experimental testing of, 735–736
 general form of, 727–728
 for magnetic fields, 930, 1049, 1052–1053. *See also* Maxwell's equations
 for magnetism, 886–887
 overview of, 727–728
 point charge inside nonspherical surface and, 726–727
 point charge inside spherical surface and, 725–726
 qualitative statement of, 720
 solid conductors and, 729–733
Geiger counter, 777
Gemstones, 1087, 1088
General theory of relativity, 415, 1243–1244. *See also* Relativity
Generators
 alternating-current, 1018
 direct-current, 960–962
 energy conversion in, 962
 homopolar, 967
 slidewire, 961–962
Geometric optics, 1079, 1110–1158
 cameras and, 1135–1138
 eye and, 1138–1142
 index of refraction and, 1080
 magnifiers and, 1142–1143
 microscopes and, 1143–1145
 reflection at plane surface and, 1110–1113
 reflection at spherical surface and, 1114–1122
 refraction at plane surface and, 1110–1113
 refraction at spherical surface and, 1122–1126
 sign rules for, 1112
 telescopes and, 1145–1147
 thin lenses and, 1127–1135
Global positioning systems, 1182, 1244
Global warming, 566–567
g (magnitude of acceleration due to gravity), 50, 113–114
GPS systems, 1074, 1182, 1244
Gradient, 768
 potential, 767–769
 potential energy, 221
Gram, 4, 5t, 112
Grand unified theory (GUT), 156
Gran Telescopio Canarias, 1147
Graphical method, for image location, 1120–1122
Graphs
 concavity vs. curvature of, 43
 parabolic, 46
 of sound waves, 503
 v_x - t , 42–44, 44t
 of wave function, 470–471
 x - t , 36–37, 39–40
 a_x - t , 44–45
 x - t , 44t
Grating spectrographs, 1198–1200
Gravitation, 154–156, 395–428
 acceleration due to. *See* Acceleration due to gravity
 action-reaction pairs and, 396
 black holes and, 415–419
 as conservative force, 402
 on cosmic scale, 399
 escape speed and, 403, 406
 as fundamental force, 154–155
 general theory of relativity and, 1243–1244
 measurement of, 397–398
 Newton's law of, 395–399
 satellite orbits and, 404–407
 specific gravity and, 367
 spherically symmetric bodies and, 396–397
 spherical mass distributions and, 411–413
 superposition of forces and, 398
 weight and, 399–402
 work done by, 402–404
Gravitational constant, 396
 calculation of, 397–398
Gravitational field, 691
Gravitational force(s), 101, 154–155, 412
 vs. electric force, 686, 687
 vs. gravitational potential energy, 403
 per unit mass, 691
 on point mass inside spherical shell, 413
 on point mass outside spherical shell, 411–412
 between spherical mass distributions, 412
Gravitational interaction, 154–155
Gravitational potential energy, 201–210, 402–404
 elastic potential energy and, 211, 212
 for extended bodies, 287, 310
 vs. gravitational force, 403
 motion along curves and, 206–210
 as negative value, 402–403
 nongravitational forces and, 204–205
Gravitational red shift, 418, 1243
Gravitational torque, 339–340
Gravity, 101
 acceleration due to. *See* Acceleration due to gravity
 weight and, 101
Greenhouse effect, 567
Ground-fault interrupters, 865
Grounding wires, 864–865
Guitars, string vibration in, 519–520
Gyroscopes, 322–324
- H**
Hall, Edwin, 904
Hall effect, 904–905
Halley's Comet, 410
Harmonic analysis, 488
Harmonic content, 488, 505
Harmonic motion, damped, 1007
Harmonic oscillators, 432
Harmonics, 487
Harmonic series, 487
Hearing, 501, 505, 506, 513, 1162f
Heat, 552–567
 added in thermodynamic process, 617–618
 of combustion, 558
 as energy in transit, 594
 energy transfer and, 552–555. *See also* Heat transfer
 of fusion, 556–557
 global warming and, 566–567
 mechanical energy and, 552
 melting and, 556–557
 phase changes and, 556–557
 quantity of, 552–555
 sign rules for, 614
 specific, 552–553
 steam, 558
 of sublimation, 558
 vs. temperature, 552
 units of measure for, 552, 553
 of vaporization, 556–558
Heat calculations, 558–560
Heat capacity, 594–597
 of gases, 594–596, 626–628, 627t
 molar, 554–555, 594–597
 point-molecule model of, 594–595
 ratio of, 627–628
 of solids, 596–597
 temperature variation of, 597
 vibration and, 595–596
Heat current, 561
Heat engines, 644–647
 Carnot, 653–659, 662–663. *See also* Carnot cycle
 Carnot cycle
 energy-flow diagrams and, 645
 hot and cold reservoirs and, 644–645
 internal-combustion, 615, 624, 647–649
 refrigerators as, 649–651
 thermal efficiency of, 645
Heat flow, 552
Heat pumps, 651
Heat transfer, 552–555, 561–567
 by conduction, 561–564
 by convection, 564
 mechanisms of, 561–568
 by radiation, 565–567
Henry, 990
Henry, Joseph, 880, 990
Hertz, 430, 1050
Hertz, Heinrich, 430, 1050, 1068, 1078
High-pass filters, 1025
Hohmann transfer orbit, 427
Hole conduction, 904
Holes, in semiconductors, 813
Holography, 1206–1207
Homopolar generators, 967
Hooke, Robert, 182
Hooke's law, 182, 183, 345, 349
 elastic deformations and, 345, 350, 351
 limits of, 351
 simple harmonic motion and, 431
Horsepower, 189
Hot reservoirs, 644–645
Hot side of line, 863, 864
House wiring systems, 863–866, 1037
 h (Planck's constant), 937
Hubble Space Telescope, 1115, 1174, 1212
Human body
 blood flow, 383
 blood flow in heart, 65
 DNA Spring, 234
 energy of locomotion, 200
 forces on dancer's body, 127
 lung volume in, 33
 magnetic fields of, 882f

I-8 Index

- Human body (*continued*)
moment of inertia, 336
as thermodynamic system, 619
- Huygens's principle, 1099–1101
diffraction and, 1186–1187
- Hydrometers, 374
- Hyperopia, 1139–1142
- Hysteresis, 941
- Hysteresis loops, 941
- I**
- Ibn Sahl, 1081
- Ice, melting of, 555–560
- Icepail experiment, Faraday's, 735
- Ideal fluid, 376
- Ideal gas, 581
adiabatic process for, 629–631
Carnot cycle for, 654–655
heat capacities of, 626–628
internal energy of, 625
isothermal expansion of, 616
kinetic-molecular model of, 588–594
volume of, 582
- Ideal-gas constant, 581–582
- Ideal-gas equation, 580–584
- Ideal-gas temperature scale, 545, 546, 659
- Idealized models, 3
- Image
erect, 1113
inverted, 1113
in optics, 1111
real, 1111
virtual, 1111, 1132
- Image distance, 1112
- Image formation
by cameras, 1135–1138
by diffraction, 1204
by lenses, 1127–1129
by reflection, 1110–1113. *See also* Mirrors
by refraction, 1122–1126
- Image point, 1110, 1111
- Impedance, 1027–1029
- Impulse, 236–237
- Impulse–momentum theorem, 236–238, 475–476
- Incident waves, 484
- Incubators, 566
- Index of refraction, 1060–1061, 1080
birefringence and, 1096
dispersion and, 1088–1090
of eye, 1139
of gemstones, 1087
laws of reflection and refraction and, 1081–1083
of lens, 1129–1130
of reflective/nonreflective coatings, 1174
total internal reflection and, 1085–1088
transparency and, 1082
wave aspects of light and, 1080, 1083
- Induced charges, 683
molecular model of, 799–800
polarization and, 684, 795–796
- Induced current, 954–955
direction of, 962–964
magnitude of, 964
- Induced electric fields, 967–969
- Induced emf, 903, 954–955. *See also* Electromagnetic induction
applications of, 955
direction of, 957–958
magnetic flux and, 954–955, 958
- Inductance, 988–1017
magnetic-field energy and, 995–998
mutual, 988–991
 R - L circuits and, 998–1002
self-inductance, 992–995
- Induction, 683
- Inductive reactance, 1023–1024
- Inductors, 992–995
in ac circuits, 1022–1023, 1026, 1032
current and rate of change in, 1000
energy stored in, 995–998
inductive reactance of, 1023–1024
vs. resistors, 996
- Inelastic collisions, 245–249
- Inertia, 104
law of, 106
mass and, 108, 113
moment of. *See* Moment of inertia
rotational, 283
- Inertial frame of reference, 106–107, 1217, 1218, 1219–1220
Newton's first law and, 106–107
Newton's second law and, 110
simultaneity and, 1220–1222
theory of relativity and, 1217
- Inertial mass, 108, 113. *See also* Mass
- Infrared radiation, 1051–1052
- Infrasound, 501
- Instantaneous acceleration, 42–44, 70–72. *See also* Acceleration
angular, 276
- Instantaneous acceleration vector, 70–71. *See also* Acceleration vectors
- Instantaneous angular acceleration, 276
- Instantaneous angular speed, 280
- Instantaneous angular velocity, 274
- Instantaneous power, 187–188
in waves, 478–479
- Instantaneous speed, 38. *See also* Speed
angular, 280
- Instantaneous velocity. *See also* Velocity
vs. average velocity, 37, 67–68
vs. instantaneous speed, 38
motion in two or three dimensions and, 37–40
straight-line motion and, 37–40. *See also* Velocity
- Instantaneous velocity vector, 67–69
- Instantaneous x -acceleration, 42
- Instantaneous x -velocity, 37
- Insulators, 542, 682–684
- Integral principles, 238
- Integral(s)
line, 185, 186, 747
moment of inertia, 289
surface, 723
- Integration, velocity and position by, 53–55
- Intensity
of electromagnetic radiation, 1063
inverse-square law for, 480
pressure amplitude and, 510–511
in single-slit diffraction, 1190–1194
sound, 510–514
wave, 480
- Intensity maxima, 1191–1192
- Interactions. *See* Particle interactions
- Interference, 481–482, 520–522, 1159–1184
amplitude in, 1166–1167
coherent sources and, 1159–1162
constructive, 484, 520–521, 1160–1162, 1164–1165, 1202
destructive, 484, 520, 522, 1160–1162, 1164–1165
vs. diffraction, 1187
in holography, 1206–1207
Michelson interferometer and, 1175–1176
Michelson-Morley experiment and, 1176
Newton's rings and, 1173–1174
nodal/antinodal curves and, 1162
in noise reduction, 522
path difference and, 1168–1169
phase difference and, 1168–1169
phase shifts and, 1170–1171
during reflection, 1170–1171
reflective/nonreflective coatings and, 1174
sinusoidal waves and, 1160
sound waves and, 520–522, 1184
standing waves and, 484, 1160, 1162
superposition and, 1159–1162
in thick films, 1171–1172
in thin films, 1170–1175
in three dimensions, 1160
of traveling waves, 520
in two dimensions, 1160
two-source/slit, 1163–1165, 1189
water waves and, 1163
Young's experiment for, 1163–1165, 1175
- Interference fringes, 1164, 1170, 1174
Newton's, 1173–1174
- Interference maxima, 1194
- Interference patterns, 1162
intensity in, 1166–1169
- Interferometer, 1175–1176
- Intermolecular forces, 585–587
- Internal-combustion engines, 615, 624, 647–649
Diesel cycle in, 648–649
otto cycle in, 647–648
- Internal energy, 218, 613, 618, 619, 661
change in, 620–623, 627–628
of cyclic processes, 620–623
entropy and, 661
first law of thermodynamics and, 618–623
of ideal gas, 625
of isolated systems, 620–623
notation for, 619
potential energy and, 618
temperature and, 625
- Internal forces, 241
torque and, 306
- Internal resistance, 824, 825, 827
- International Bureau of Weights and Measures, 4
- International Space Station, 404f, 406
- International System (SI) units, 4, 5. *See also* Units of measure
- Interplanetary dust, 1074
- Inverse-square law, for intensity, 480
- Inverse-square law, for sound intensity, 511
- Inverted image, 1113
in camera lens, 1137
- Ionization, 681
corona discharge and, 761–762
- Ions, 681
- Irreversible process, 642–643
- Isobar, 625
- Isobaric process, 624
- Isochor, 625
- Isochoric process, 624
- Isolated systems, 241
internal energy of, 620–623
- Isotherm, 584–585, 625
- Isothermal expansion, of ideal gas, 616
- Isothermal process, 624–625
Carnot cycle and, 653
- Isotopes, 892
- Isotropic, 1079
- J**
- Joule, 172, 177, 202, 552, 553
- Joule, James, 172, 552
- Joule per coulomb, 754
- Junctions, in circuits, 849–850
- Jupiter, magnetic field of, 970
- K**
- Kaons, 1252
- Kelvin, 546
- Kelvin–Planck statement, 651
- Kelvin temperature scale, 545–546, 655, 658–659
- Kepler, Johannes, 407–408
- Kepler's first law, 408
- Kepler's second law, 408–409
- Kepler's third law, 409, 418
- Kilocalorie (kcal), 552
- Kilogram, 4, 5, 109
- Kilohm, 820

Kilometer, 5, 5t
 Kilowatt, 5, 188
 Kilowatt-hour, 188, 865
 Kinematics, 34. *See also* Motion
 linear vs. angular, 279–282
 Kinetic energy, 177
 in collisions, 245–246
 in composite systems, 181–182
 conservative forces and, 215–219
 equipartition of, 595
 gas pressure and, 589–590
 heat capacities and, 594–597
 molecular, 585–587, 589–590, 594, 625
 moment of inertia and, 282–285
 vs. momentum, 238–239
 physical meaning of, 180
 potential energy and, 201, 202, 215–216
 relativistic, 1239–1240
 rotational, 282–287, 308–309
 as scalar quantity, 177
 in simple harmonic motion, 438–441
 torque and, 315
 units of, 177
 with varying forces, 181–187
 work–energy theorem and, 176–181
 Kinetic friction, 141–143
 coefficient of, 142
 as nonconservative force, 216
 Kinetic friction force, 142
 Kinetic-molecular model, of ideal gas, 588–594
 Kirchhoff, Gustav Robert, 849
 Kirchhoff's rules, 849–854, 972
 Kundt's tube, 514

L

Lagging current, 1033
 Laminar flow, 376, 382–383
 Large Hadron Collider, 1249
 Laser light, 1051, 1078, 1160, 1162
 Laser printers, 680, 762
 Laser(s), 1162
 in holography, 1206
 light emission, 1078
 Latent heat of fusion, 556–557
 Lateral magnification, 1113, 1117
 vs. angular magnification, 1143
 of camera, 1136
 of microscope, 1144
 Laue, Max von, 1200
 Law of Biot and Savart, 922
 Law of conservation of energy, 171, 203–204, 218
 Law of inertia, 106. *See also* Newton's first law of motion
 Law of reflection, 1081–1083
 Law of refraction, 1081–1083
 Laws, physical, 1–2
 L-C circuits, 1002–1006
 Leaning Tower of Pisa, 2
 Length
 proper, 1228
 relativity of, 1227–1231
 units of, 4, 5
 Length contraction, 1228
 Lenses, 1127
 of camera, 1135–1138
 corrective, 1139–1142
 of eye, 1138–1139
 magnifying, 1142–1143
 of microscopes, 1143
 nonreflective coatings for, 1174
 properties of, 1127
 reflective coatings for, 1174
 properties of, 1127
 of telescopes, 1145–1146
 thin, 1127–1135. *See also* Thin lenses
 Lensmaker's equation, 1129–1130
 Lenz, H. F. E., 962
 Lenz's law, 962–964, 965
 Lever arm, 303
 Light, 1077–1109
 beams of, 1079
 coherent, 1160
 diffraction of, 1185–1216
 dispersion of, 1083, 1088–1090
 Doppler effect for, 524–529, 1235–1236
 early studies of, 1077–1078
 fluorescent, 994, 1078
 Huygens's principle and, 1099–1101
 intensities of, 1082
 interference and, 1163–1165. *See also* Interference
 laser, 1051, 1078, 1160, 1162
 monochromatic, 1051, 1078, 1160
 natural, 1091
 as photons. *See* Photons
 polarized, 1090–1097
 rays of, 1079
 reflection of, 1079–1085. *See also* Reflection
 refraction of, 1079–1085. *See also* Refraction
 scattering of, 1097–1098
 speed of, 4, 1052–1057, 1078, 1218–1219, 1252
 total internal reflection of, 1085–1088
 unpolarized, 1091
 visible, 1051
 as wave and particle. *See* Wave–particle duality
 wave fronts of, 1078–1079
 wavelengths of, 1051–1052
 Lightning rods, 762
 Limit of resolution, 1204
 Linear acceleration, 276
 vs. angular acceleration, 278t
 constant, 278t
 in rigid-body rotation, 280–282
 Linear charge density, 695
 Linear conductors, 817
 Linear expansion, 547–558, 548t
 Linear mass density, 474
 Linear momentum, 236, 316
 Linear polarization, 1055, 1090, 1092
 of electromagnetic wave, 1055
 Linear speed, in rigid-body rotation, 279–280
 Linear superposition, 482
 Linear velocity, 276
 vs. angular velocity, 274
 Linear vs. angular kinematics, 279–282
 Line integral, 185, 186, 747
 of electric fields, 932–933
 of electrostatic force, 747, 933
 of magnetic fields, 933
 Line of action, 303
 Liquid crystal display (LCD), 1097, 1137
 Liquid crystals, 1097f
 Liquid phase, 555–556
 Liquid(s)
 compressibility of, 349
 molecular speed in, 599
 molecules in, 586
 phases of, 599–602
 Longitudinal waves, 465. *See also* Mechanical waves;
 Wave(s)
 periodic, 468
 sound, 468
 wave function for, 473
 Long-range forces, 101, 116
 Loops, in circuits, 849–850
 Lorentz factor, 1224, 1228
 Lorentz transformations, 1231–1235
 coordinate, 1231–1232
 velocity, 1232–1233
 Loudness, 505
 Loudspeakers, 895, 1026
 Low-pass filters, 1024
 L-R-C parallel circuits, resonance in, 1035
 L-R-C series circuits, 1007–1008
 with ac source, 1027–1030
 impedance in, 1028–1029, 1035
 phase angle and, 1028–1029
 power in, 1033
 resonance in, 1034–1036

M

Mach number, 530
 Macroscopic properties
 vs. microscopic properties, 579
 theories of matter and, 588
 Macroscopic state, 666
 Magnetic ballast, 994
 Magnetic bottles, 889
 Magnetic declination, 879
 Magnetic dipole moment. *See* Magnetic moment
 Magnetic dipoles, 898, 901
 force and torque on, 897–902
 of magnets, 901
 in nonuniform magnetic fields, 900
 potential energy for, 899
 Magnetic domains, 940
 Magnetic energy density, 996–997
 Magnetic-field energy, 995–998
 Magnetic field lines, 879, 884–887
 for current element, 922
 direction of, 919
 magnetic flux and, 884–887
 vs. magnetic lines of force, 884
 for moving charge, 919–920
 with no endpoints, 886
 Magnetic field(s), 878–917
 on axis of coil, 928–929
 calculation of, 922
 of circular current loops, 927–930
 critical, 975
 of current element, 921–923
 direction of, 881–882
 of earth, 878, 879, 882
 of electromagnetic waves, 1052–1057, 1058–1060,
 1066–1069
 Gauss's law for, 886–887, 930, 1049, 1052–1053
 Hall effect and, 904–905
 of human body, 882f
 line integral of, 933
 of long cylindrical conductor, 934
 of long straight conductor, 930–932, 934
 magnitude of, 882, 887
 measurement of, 882–883, 975
 motion in, 888–893
 of motors, 893–896
 of moving charge, 880, 918–921
 nodal/antinodal planes of, 1067
 notation for, 880
 of solenoid, 935–936
 sources of, 918–952, 971–972
 of straight current-carrying conductor, 923–926
 superposition of, 921
 test charges for, 882–883
 vector, 880, 919
 Magnetic flux, 884–887
 calculation of, 955
 Faraday's law and, 955–962
 Gauss's law for magnetism and, 886–887
 induced electric fields and, 967–969
 induced emf and, 954–955
 Lenz's law and, 964
 Meissner effect and, 975
 superconductivity and, 975–976
 in transformers, 1037
 units for, 886
 Magnetic flux density, 887
 Magnetic force(s)
 on current-carrying conductors, 893–896
 on current loops, 897–902
 direction of, 881–882
 in electric motors, 893–896
 as fundamental force, 154–155
 Hall effect and, 904–905
 in loudspeakers, 895
 magnitude of, 882
 between parallel conductors, 926–927
 units for, 882
 Magnetic inclination, 879
 Magnetic lines of force, 884

I-10 Index

- Magnetic materials, 901, 937–941
 Bohr magneton, 937–938
 diamagnetic, 939–940
 ferromagnetic, 940–941
 paramagnetic, 938–939
 relative permeability of, 938
- Magnetic moment, 898, 901
 alignment of, 937–941
 of current loop, 898, 929
 direction of, 898
 magnitude of, 938
 notation for, 938
 vector, 898
- Magnetic monopoles, 880
- Magnetic poles, 879–880
 vs. electric charge, 879–880
- Magnetic resonance imaging (MRI), 898f, 917, 929f, 976, 1017
- Magnetic susceptibility, 939
- Magnetic torque, 897–902
- Magnetic variation, 879
- Magnetism, 878–880
 electron motion and, 880
 Gauss's law for, 886–887, 930
- Magnetization, 901, 922, 937–941
 Bohr magneton and, 937–938
 diamagnetism and, 939–940
 ferromagnetism and, 940–941, 994
 paramagnetism and, 938–939
 saturation, 940
- Magnetization curve, 940
- Magneton, Bohr, 937–938
- Magnetotactic bacteria, 952
- Magnetrons, 888
- Magnet(s)
 attracting unmagnetized objects, 901
 bar, 901
 magnetic dipoles of, 901
 magnetic moment of, 901
 permanent, 878–879, 937–938
- Magnification
 angular, 1142–1143, 1146
 lateral, 1113, 1117, 1136, 1143, 1145
- Magnifiers, 1142–1143
- Magnitude, of vectors, 10, 12, 14–16
- Malus, Étienne-Louis, 1092
- Malus's law, 1093
- Manometers, 371
- Marconi, Guglielmo, 1050
- Mars, gravity on, 401
- Mass, 108, 113
 acceleration and, 108–109, 113–114, 116
 of atom, 587, 680, 891–892
 center of, 253–256
 density and, 366
 of electron, 680, 891–892
 force and, 108–109
 of gas, 581–582
 inertial, 108, 113
 measurement of, 108–109, 114
 molar, 554–555, 580, 587
 of molecule, 587, 891–892
 of neutron, 680
 Newton's second law and, 108–109, 110, 113–115
 of proton, 680
 rest, 1237–1239
 of star, 1251
 terminal speed and, 146–149
 units of, 4–5, 109, 112
 weight and, 109, 113–115
- Mass-energy conservation law, 1241
- Mass per unit length, 474
- Mass spectrograph, 912
- Mass spectrometers, 892
- Mass-to-volume ratio, density and, 367
- Matter
 condensed. *See* Liquid(s); Solids
 molecular properties of, 585–588
 phases of, 599–602
- Maxwell–Boltzmann distribution, 598–599
- Maxwell, James Clerk, 972, 1049, 1078
- Maxwell's equations, 880, 953, 972–974
 electromagnetic waves and, 1048–1052, 1218
 Huygens's principle and, 1099–1101
 in optics, 1082
- Maxwell's wave theory, 1048–1052
- Mean free path, 593
- Mean free time, 832
- Measurement
 accuracy in, 8, 9
 precision in, 9
 significant figures in, 8–9
 uncertainty in, 8
 units of, 3–7. *See also* Units of measure
- Mechanical energy
 conservation of, 203–204, 438–441, 748
 conservative vs. nonconservative forces, 215–219
 heat and, 552
 in simple harmonic motion, 438–441
 total, 203–204
- Mechanical waves, 464–500. *See also* Wave(s)
 amplitude of, 470
 boundary conditions for, 481
 direction of, 471, 473
 energy of, 478–481
 energy transport by, 465–466
 equilibrium state for, 465
 incident, 484
 intensity of, 480
 interference in, 481
 longitudinal. *See* Longitudinal waves
 mathematical description of, 469–474
 normal-mode patterns of, 486–490
 periodic, 466–469. *See also*
 Periodic waves
 power of, 478–480
 propagation of, 465–466
 sinusoidal, 466–467. *See also* Sinusoidal waves
 sound, 468
 speed of. *See* Wave speed
 standing, 483–490
 superposition of, 482, 488
 transverse. *See* Transverse waves
 traveling, 483–485
 types of, 464–466
 wave equation for, 473, 477
 wavelength of, 467
- Mechanics, 34
 celestial, 395
 classical (Newtonian), 100, 1218–1219
 fluid. *See* Fluid mechanics
 quantum. *See* Quantum mechanics
- Medium, 464–465
- Megawatt, 188
- Megohm, 820
- Meissner effect, 975
- Melting, 556–557
- Membrane potential, 768, 811
- Mercury barometers, 371–372
- Metallic conduction, 832–834
- Metals
 breaking stresses of, 352t
 as conductors, 561
 elastic moduli of, 347t
 thermal conductivities of, 561t
- Meter, 4
- Method of components, 14
- Michelson, Albert, 1176
- Michelson interferometer, 1175–1176
- Michelson–Morley experiment, 1176, 1218
- Microampere, 814
- Microcoulomb, 687
- Microfarad, 784
- Microgram, 5t
- Micrographs, 1145
- Micrometer, 5t
- Microphones, condenser, 783
- Microscopes
 dynamic atomic force, 463
 optical system of, 1143–1145
 resolving power of, 1204
- Microscopic state, 666
- Microscopic vs. macroscopic properties, 579
- Microsecond, 5t
- Microwaves, 1051
- Milliampere, 814
- Millibar, 368
- Milligram, 5t
- Millikan, Robert, 892
- Millikan's oil-drop experiment, 779
- Millimeter, 5t
- Millisecond, 5t
- Mirages, 1100
- Mirrors. *See also* Reflection
 concave, 1114–1118
 converging, 1114–1118
 convex, 1118–1120
 graphical methods for, 1120–1122
 image formation by, 1110–1113
 inverted image, 1113
 parabolic, 1116
 plane, 1111–1112
 spherical, 1114–1122
- Mitchell, John, 416
- Models, 3
 idealized, 3
- Molar heat capacity, 554–555, 594–597, 626–628, 627t
- Molar mass, 554–555, 580, 587
- Molar specific heat, 554–555
- Molecular bonds, 595–596
- Molecular collisions, 592–594
 gas pressure and, 588–589
- Molecular kinetic energy, 585–587, 594–595
 gas pressure and, 589–590
 temperature and, 625
- Molecular mass, 587
 measurement of, 892
- Molecular speed, 591–592, 597–599
 Maxwell–Boltzmann distribution and, 598–599
- Molecular vibration, 443–445, 586, 595–596
- Molecular weight, 554–555, 580
- Molecules
 gas, 586, 588–589
 intermolecular forces and, 585–587
 in liquids, 586
 vs. moles, 588
 polar, 799–800
 polyatomic, 595
 in solids, 586
- Moles, 554, 587, 588
- Moment arm, 303
- Moment of inertia, 282–285
 calculation of, 284–285, 289–290, 447
 of cylinder, 285t, 289–290
 in human body and sports, 336
 parallel-axis theorem and, 287–288
 of rectangular plate, 285t
 in simple harmonic motion, 443
 of slender rod, 285t
 of sphere, 285t, 290
 torque and, 305–306
- Momentum, 235–271
 angular. *See* Angular momentum
 and archerfish, 271
 collisions and, 245–252
 components of, 236
 conservation of, 241–245, 249–251, 1237
 electromagnetic, 1064–1065
 impulse and, 236–241
 impulse–momentum theorem and, 236–238, 475–476

- vs. kinetic energy, 238–239
- linear, 236, 316
- magnitude of, 236
- net force and, 236
- Newton's second law and, 236
- Newton's third law and, 241–242
- rate of change of, 236
- relativistic, 1237–1239
- rocket propulsion and, 256–259
- in simple harmonic motion, 441
- total, 241–242, 254
- transverse, 475–476
- units of, 236
- as vector, 236, 239
- wave speed and, 475–476
- Monatomic gases, 595
- Monochromatic light, 1051, 1078, 1160
- Monopoles, magnetic, 880
- Moon, 399, 400f, 403, 406
- Morley, Edward, 1176
- Motion
 - along curve. *See* Curve(s)
 - of center of mass, 254–256
 - circular, 149–154, 432–434. *See also* Circular motion
 - Kepler's laws of, 407–409
 - Newton's laws of, 100–170. *See also*
 - Newton's laws of motion
 - orbital, 404–407
 - periodic, 429–463. *See also* Oscillation
 - planetary, 407–411
 - projectile, 74–81. *See also* Projectile motion
 - rotational. *See* Rotation/rotational motion
 - of satellites, 404–407
 - simple harmonic, 431–445
 - straight-line, 34–65. *See also* Straight-line motion
 - translational, 302, 308–314, 595
 - in two or three dimensions, 66–99. *See also*
 - Motion in two or three dimensions
- Motional electromotive force, 965–967
- Motion diagrams, 39–40
- Motion in two or three dimensions, 66–99
 - acceleration vector, 69–74
 - circular, 81–84
 - frame of reference for, 85
 - position and velocity vectors, 66–69
 - projectile, 74–81
 - relative velocity, 84–89
- Motors, electric, 893–896, 902–904
- Moving-axis rotation, 308–314
- MRI (magnetic resonance imaging), 898f, 917, 929f, 976, 1017
- Müller, Karl, 818
- Multimeters, digital, 857
- Multiplication, of vectors, 16
- Muons, 1247–1248
- Muscle fibers, work done by, 173
- Musical instruments
 - pipe organs, 516–520
 - standing waves and, 488–489
 - string, 486–490
 - wind instruments, 518
- Music, sound waves in, 505–506
- Mutual inductance, 988–991
- Myopia, 1139–1142
- N**
- Nanocoulomb, 687
- Nanometer, 5t
- Nanoparticles, 941f
- Nanosecond, 5t
- Natural angular frequency, 452
- Natural light, 1091
- Near-field diffraction, 1186
- Near point, of eye, 1139
- Nearsightedness, 1139–1142
- Negative ions, 681
- Negative work, 173–175, 177
- Neon, 892
- Nerve cells, 734f, 818f, 877
- Net force, 102–103, 241–242
 - acceleration and, 107–113
 - center-of-mass motion and, 255–256
 - circular motion and, 108
 - momentum and, 236–238
 - Newton's first law and, 104–105, 106–107
 - Newton's second law and, 107–113
 - torque and, 305, 316, 317
 - zero, 104–105, 106–107
- Net torque, 315
- Neutral side of line, 863, 864
- Neutrons, mass of, 680
- Newtonian mechanics, 100, 1218–1219
- Newtonian synthesis, 411
- Newton, Isaac, 100, 1077
- Newton-meter, 172, 303
- Newton (N), 5, 101, 109, 112
- Newton's first law of motion, 100, 103–107
 - application of, 119, 129–134
 - equilibrium and, 105, 129–130
 - inertia and, 104, 106
 - net force and, 104–105
 - statement of, 104
- Newton's law of gravitation, 395–399
- Newton's laws of motion, 100–128
 - application of, 119–120
 - first law, 103–107. *See also* Newton's first law of motion
 - free-body diagrams for, 119–120, 134
 - inertial frame of reference and, 106–107
 - Kepler's laws and, 408–409, 411
 - relativity and, 1237–1239, 1243–1244
 - second law, 107–115. *See also* Newton's second law of motion
 - third law, 115–119. *See also* Newton's third law of motion
- Newton's rings, 1173–1174
- Newton's second law of motion, 107–115
 - acceleration and, 107–113
 - application of, 110–111, 119, 134–141
 - component equations for, 110
 - fluid resistance and, 146–149
 - inertial reference frames and, 110
 - mass and, 108–109, 110, 113–115
 - momentum and, 236
 - net force and, 107–113
 - relativity and, 1237–1239, 1243–1244
 - rotational analog of, 306, 312
 - statement of, 109–110
 - weight and, 109, 113–115
- Newton's third law of motion, 115–119
 - action–reaction pairs and, 115–119
 - application of, 116–118, 119
 - fluid resistance and, 146–149
 - gravitational forces and, 398
 - momentum and, 241–242
 - statement of, 115
 - tension and, 118
- Nodal curves, 1162
- Nodal planes, 1067
- Nodes, 483
 - displacement, 514–515
 - pressure, 515
- Noise, 506
- Noise reduction, wave interference in, 522
- Nonconservative forces, 215–219
- Nonelectrostatic fields, 955, 968–969
- Nonlinear conductors, 817
- Nonreflective coatings, 1174
- Nonuniform circular motion, 84, 154
- Normal force, 101, 141
- Normal mode, 487
- North (N) pole, 879
- Nuclear fission, 1241
- Nuclear fusion, 791, 927
- Nuclear power plants, 558, 1241
- Nuclear reactions, fission, 1241
- Nuclear reactors, 1241
- Nucleus, atomic, 680
- Nuclides, radioactive. *See* Radioactivity
- O**
- Object distance, 1111
- Object, in optics, 1110–1111
- Objective, microscope, 1144
- Object point, 1111
- Ocean thermal energy conversion, 677
- Oculars, 1144
- Oersted, Hans Christian, 880
- Ohm, 820
- Ohm, Georg Simon, 816
- Ohmic conductors, 817
- Ohmmeters, 857
- Ohm's law, 816, 819–821
- 1 newton per coulomb, 691
- Onnes, Heike Kamerlingh, 818
- Open circuits, 864
- Open orbits, 404
- Operational definition, 4
- Optical fibers, 1087
- Optic axis, of mirror, 1114
- Optics, 1077. *See also* Light
 - geometric. *See* Geometric optics
 - image in, 1111
 - object in, 1110–1111
 - physical, 1079, 1159
- Orbital eccentricity, 408
- Orbital period, 409
- Orbital speed, 409
- Orbit(s)
 - center of mass and, 410–411
 - circular, 405–406, 409
 - closed, 404
 - of Comet Halley, 410
 - elliptical, 408–409
 - open, 404
 - satellite, 404–407
 - sector velocity and, 408
 - semi-major axis of, 408, 409
- Order-of-magnitude estimates, 10
- Organ pipes, 516–520
- Oscillation, 429–463
 - amplitude of, 430
 - damped, 449–451
 - displacement in, 429
 - driven, 451–453
 - electrical, 1002–1006
 - forced, 451–453, 518
 - frequency of, 430–431
 - in harmonic oscillators. *See*
 - Harmonic oscillators
 - molecular vibration and, 443–445
 - overview of, 429–431
 - of pendulum, 446–449
 - of periodic waves, 466–469
 - resonance and, 453, 518–520
 - simple harmonic motion and, 431–445. *See also*
 - Simple harmonic motion (SHM)
 - in spring, 429–430
- Oscillation cycle, 430–431
- Oscillation period, 430–431
- Otto cycle, 647–648
- Overdamped circuits, 1007–1008
- Overdamping, 450
- Overloaded circuits, 864–865
- Overtones, 487
- P**
- Pacemakers, 860f
- Parabola, equation for, 76
- Parabolic graphs, 46
- Parabolic mirrors, 1116
- Parabolic trajectory, 76
- Parallel-axis theorem, 287–288
- Parallel connection, 787–788
- Parallel-plate capacitors, 783
 - dielectrics in, 794
- Parallel resistors, 845
- Parallel vectors, 11
- Paramagnetism, 938–939

I-12 Index

- Paraxial approximation, 1115
Paraxial rays, 1115
Partial derivatives, 220–221, 472–473
Particle accelerators
 cyclotrons and, 888, 910
Particle interactions
 electromagnetic, 154–156
 fundamental types of, 154–156
 gravitational, 154–156
 strong, 155
 weak, 155–156
Particle motion vs. wave motion, 467
Particle(s), 3
 photons as. *See* Photons; Wave–particle duality
 straight-line motion and, 34–35
Particle speed vs. wave speed, 510
Particle velocity vs. wave velocity, 510
Pascal, 346, 368
Pascal, Blaise, 346
Pascal’s law, 369–370
Path difference, and diffraction, 1190–1191, 1195, 1197, 1201
Paths, in thermodynamic system, 617–618
Pendulum
 ballistic, 247
 periodic motion of, 446–449
 physical, 447–449
 simple, 446–447
Pendulum bob, 446
Percent error, 8
Perihelion, 408
 precession of, 1243
Period
 frequency and, 430–431
 orbital, 409
 oscillation, 430
 in simple harmonic motion, 434–435
 of uniform circular motion, 83
Periodic driving force, damped oscillation and, 452
Periodic motion, 429–463. *See also* Oscillation
 amplitude of, 430
 displacement in, 429
 frequency of, 430–431
 in harmonic oscillators, 432
 molecular vibration and, 443–445
 overview of, 429–431
 of pendulum, 446–449
 resonance and, 453
 simple harmonic motion and, 431–445. *See also* Simple harmonic motion (SHM)
 of spring, 429–430
 of waves, 466–469
Periodic waves, 466–469. *See also*
 Mechanical waves
 longitudinal, 468. *See also* Longitudinal waves
 mathematical description of, 469–474
 sinusoidal, 466–467, 469–474. *See also*
 Sinusoidal waves
 transverse, 466–468. *See also* Transverse waves
Permanent magnets, 878–879, 937–938
Permanent set, 351
Permeability, 938
Permittivity, of dielectric, 795–796
Phase, 555–556
 wave, 471
Phase angle, 436, 1023, 1028–1029
Phase change, 556–557
Phase diagrams, 600
Phase equilibrium, 556, 585, 600
Phase shifts, interference and, 1170–1171
Phases of matter, 599–602
 critical point and, 600, 601
 molecular interactions and, 599
 pVT -surfaces and, 601–602
 sublimation and, 600
 triple point and, 600
Phase transitions, 599–602
Phase velocity, 471
Phasor, 433, 1018–1021
Phasor diagrams, 1019
Phosphor, 1078
Photocopying machines, 762
Photoelasticity, 1096–1097
Photoelectric effect, 1078
Photography. *See* Cameras
Photomicrographs, 1145
Photons, 1078, 1241
 wave–particle duality and. *See* Wave–particle duality
Physical optics, 1079, 1159
Physical pendulum, 447–449
Physical quantities, 3–4
 units of, 3–7. *See also* Units of measure
Physical theories, 1–2
Physics
 as experimental science, 1
 nature of, 1–2
 overview of, 1
 problem-solving strategies for, 2–3
 as process, 2
Picoampere, 814
Picofarad, 784
Pions, 1241
Pipe organs, 516–520
Pitch, 505
 π , value of, 8
Planck’s constant, 937
Plane, 66
Plane mirrors
 graphical methods for, 1120–1122
 image formation by, 1110–1113
Plane of incidence, 1081
Plane surface
 reflection at, 1110–1113
 refraction at, 1124–1125
Planetary motion, 407–411. *See also*
 Orbit(s)
 center of mass and, 410–411
 Kepler’s laws of, 407–411
Planets. *See also specific planets*
 conservation of angular momentum and, 408–409
 detection of, 256
Plane waves, electromagnetic, 1052–1057
Plastic, 346, 351
Plastic deformation, 351–352
Pluto, 406f
Point charges, 685
 electric dipole and, 696, 701–705
 electric fields of, 718–721. *See also*
 Electric charge
 electric potential energy of, 749–752
 electromagnetic waves from, 1050
 force between, 688
 inside closed surface, 719
 inside nonspherical surface, 726–727
 inside spherical surface, 725–726
 magnetic field lines for, 919–920
 superposition of, 687
Point mass
 inside spherical shell, 413
 outside spherical shell, 411–412
Point-molecule model, of gas heat capacity, 594–595
Point objects, 1111
 image formation for, 1114–1115, 1122–1126, 1204
 resolution of, 1204
Polarization, 684, 781, 799–800, 1090–1097
 charged bodies and, 684, 800
 circular, 1095–1096
 of dielectrics, 795–796, 799–800
 electric field lines and, 701
 of electromagnetic waves, 1055, 1090–1097
 elliptical, 1095–1096
 induced charges and, 684, 799–800
 of light waves, 1090–1097
 linear, 1055, 1090, 1092
 partial, 1094
 photoelasticity and, 1096–1097
 by reflection, 1094–1095
Polarizers, 1090
Polarizing angle, 1094
Polarizing axis, 1091
Polarizing filters, 1090, 1091–1093, 1094
Polar molecules, 799–800
Polaroid filters, 1091
Polyatomic molecules, 595
Porro prism, 1086
Position. *See also* Displacement
 as function of time
 with changing acceleration, 53–55
 with constant acceleration, 44–47
 by integration, 53–55
 potential energy and, 202
Position vectors, 67–69
Positive ions, 681
Positive work, 173–175, 177
Positrons, motion in magnetic fields, 889
Potential. *See* Electric potential
Potential difference, 756. *See also* Voltage
 around circuits, 827
 capacitance and, 782, 784
 measurement of, 854
 notation for, 859
 resistance and, 845
 time-varying, 859
Potential energy, 201–234, 748
 around circuits, 827
 of capacitor, 790–793
 conservative forces and, 215–219, 223
 elastic, 210–215, 219–220
 electric, 747–753. *See also* Electric potential energy
 of electric dipole, 703–704
 electric forces and, 220
 energy diagrams and, 222–223
 equilibrium and, 222–223
 force and, 219–222
 gradient of, 221
 gravitational, 201–210, 287, 310, 402–404
 intermolecular forces and, 585–587
 kinetic energy and, 201, 202, 215–216
 for magnetic dipoles, 899
 of molecules, 586
 position and, 202
 potential wells and, 586
 in simple harmonic motion, 438–441
 units of, 202
 work and, 748
Potential gradient, 767–769
Potential wells, 586
Potentiometers, 858
Pound (pound-force), 112
Pounds per square inch, 346
Pounds per square inch absolute (psia), 370
Pounds per square inch gauge (psig), 370
Power, 187–189
 average, 187, 479
 of corrective lens, 1141
 electrical, 188
 in electric circuits, 828–832, 1031–1033
 for electric motors, 903
 energy and, 865
 force and, 187–189
 instantaneous, 187–188
 measurement of, 188, 856–857
 rotational motion and, 314–316
 of sound waves, 510–511
 velocity and, 188
 of waves, 478–480
 work and, 478–480
Power distribution systems, 863–866

- Power factor, 1032–1033
 Power plants, nuclear, 1241
 Powers-of-10 notation, 9
 Poynting vector, 1062–1064
 Precession, 322–324
 of perihelion, 1243
 Precession angular speed, 323
 Precision, vs. accuracy, 9
 Prefixes, for units of measurement, 5
 Presbyopia, 1139
 Pressure, 348–349
 absolute, 370–371
 atmospheric, 348, 368
 bulk stress/strain and, 348–349
 vs. density, 581
 on elephant in water, 393–394
 fluid flow and, 378–381
 in fluids, 348, 368–373
 vs. force, 348, 368
 gauge, 370–371
 measurement of, 371–372
 radiation, 1065
 reciprocal, 349
 residential water, 380
 as scalar quantity, 348
 speed and, 378–381
 units of, 346, 349
 Pressure amplitude, 503–504
 sound intensity and, 510–511
 Pressure gauges, 371–372
 Pressure nodes/antinodes, 515
 Primary windings, 1036
 Principal maxima, 1196
 Principal rays
 for lenses, 1131–1132
 for mirrors, 1120–1122
 Principle-ray diagrams, 1120–1122, 1132–1134
 Principle(s)
 conservation of momentum, 242
 differential, 238
 integral, 238
 physical, 1–2
 Printers, laser, 680, 762
 Prism
 dispersion by, 1088
 Porro, 1086
 Prism binoculars, 1146
 Problem solving
 idealized models for, 3
 strategies for, 2–3
 Product
 scalar, 19–21, 22
 of vectors, 19–24
 Projectile, 74
 Projectile motion, 74–81
 acceleration and, 74–76
 air resistance and, 74–76
 vs. circular motion, 83
 components of, 74–76
 trajectory and, 74–76
 velocity and, 74–76
 Projectors, 1137
 Propagation speed, 465–466, 467
 Proper length, 1228
 Proper time, 1223–1225, 1226
 Proton(s)
 charge of, 686
 mass of, 680
 Psia (pounds per square inch absolute), 370
 Psig (pounds per square inch gauge), 370
pV-diagrams, 584–585
pV-isotherms, 584–585
pVT-surfaces, 601–602
- Q**
 Quanta, 1078. *See also* Photons
 Quantity of heat, 552–555
 Quantum electrodynamics, 1078
 Quantum mechanics, 595–596, 597, 974
 Quarks, 680
 Quarter-wave plates, 1096
 Quasars, 1213
- R**
 Radar, Doppler, 529
 Radial acceleration, 82, 149–150
 Radians, 273, 281
 Radiation
 absorption of, 565–566
 applications of, 566
 blackbody, 566
 cancer and, 757f, 941f
 Čerenkov, 1249–1250
 electromagnetic, 565–567, 1050–1051. *See also*
 Electromagnetic wave(s)
 global warming and, 566–567
 solar, 566–567
 Stefan–Boltzmann law/constant and, 565
 thermal, 1078
 x. *See* X rays
 Radiation pressure, 1065
 Radiator, ideal, 566
 Radioactivity
 beta decay, 115
 charged particles and, 891
 electromagnetic waves and, 1048
 Radiocarbon dating, 155–156
 Radios
 transmitters and receivers for, 1050–1051
 tuning, 1035–1036
 Radio waves, 1049, 1051
 Radius of curvature
 for lens, 1129–1130
 for spherical surface, 1112, 1124
 Radius, Schwarzschild, 416–417, 418
 Rainbows, 1089–1090
 Randomness, in thermodynamic
 processes, 643
 Range of validity, 2
 Rarefaction, 468
 Ratio of heat capacities, 508–509,
 627–628
 Rayleigh, Lord, 1204
 Rayleigh's criterion, 1204
 Rays, 1079, 1131–1132
 paraxial, 1115
 principal, 1120–1122, 1131–1132
R-C circuits, 858–863, 877
 Reaction(s)
 activation energy for, 599
 chemical, 599
 nuclear. *See* Nuclear reactions, fission
 Real image, 1111
 Reciprocal pressure, 349
 Rectified alternating current, 1019–1020
 Rectified average current, 1020
 Rectifiers, 1019–1020
 Red shifts, gravitational, 418, 1243
 Reference circle, 433
 Reference frame, inertial, 106–107, 110
 Reference point, 433
 Reference standards, 4
 Reflected waves, 481
 sinusoidal, 482
 Reflecting telescopes, 1147
 Reflection, 1079–1085
 Bragg, 1202, 1216
 diffuse, 1080, 1111
 of electromagnetic waves, 1068
 Huygens's principle and, 1099
 image formation and, 1110–1113
 interference during, 1170–1171
 law of, 1081–1083
 phase shifts during, 1170–1171
 at plane surface, 1110–1113
 polarization by, 1094–1095
 specular, 1080, 1111
 at spherical surface, 1114–1122
 total internal, 1085–1088
 in x-ray diffraction, 1201–1202
 Reflection gratings, 1197
 Reflective coatings, 1174
 Reflector, ideal, 566
 Refracting telescopes, 1145–1146
 Refraction, 1079–1085
 in eye, 1139
 Huygens's principle and, 1100–1101
 index of. *See* Index of refraction
 law of, 1081–1083
 at plane surface, 1110–1113
 at spherical surface, 1122–1126
 Refractive index. *See* Index of refraction
 Refrigerators, 649–651
 Carnot, 657
 practical, 650–651
 workless, 651
 Refrigerator statement, of second law of
 thermodynamics, 652
 Relative permeability, 938
 Relative velocity, 84–89. *See also* Velocity
 elastic collisions and, 250–251
 frame of reference for, 85, 1233f
 Galilean velocity transformation and, 87
 in one dimension, 85
 in two or three dimensions, 86–87
 Relativistic momentum, 1237–1239
 Relativistic work and energy, 1239–1242
 Relativity, 1217–1252
 aging and, 1226
 Doppler effect and, 528, 1235–1236
 Einstein's postulates for, 1218–1219
 Galilean coordinate transformation and,
 1219–1220
 general theory of, 415, 1176, 1243–1244
 inertial frame of reference and, 1217, 1218,
 1219–1220
 invariance of physical laws and,
 1217–1220
 of length, 1227–1231
 Lorentz transformations and, 1231–1235
 Newtonian mechanics and, 1218–1219,
 1237–1239, 1243–1244
 principle of, 1217–1218
 of simultaneity, 1220–1222
 special theory of, 1217–1244
 speed of light and, 1218–1219
 of time intervals, 1222–1227
 twin paradox and, 1226
 Relaxation time, 860–861
 Resistance, 819–822
 equivalent, 845, 846
 internal, 824, 825, 827
 measurement of, 854–858
 Resistance thermometer, 544, 545
 Resistivity
 conductor resistance and, 816–819
 of metal, 832–834
 temperature and, 817–818
 Resistors, 820–821
 in ac circuits, 1022, 1026, 1031
 vs. capacitors, 846
 in dc circuits, 844–847
 energy dissipation in, 829
 equivalent resistance and, 845
 vs. inductors, 996
 in parallel, 845, 846–847
 power in, 1031
 power input to, 828–829
 power rating of, 829
 in series, 844–845
 shunt, 855
 Resolving power (resolution), 1204
 of camera lens, 1205
 chromatic, 1198–1200, 1204
 in diffraction, 1204

- Resolving power (*continued*)
 of grating spectrograph, 1198–1200
 limit of, 1204
- Resolving power (resolution) (co)
 Rayleigh's criterion and, 1204
- Resonance, 453, 518–520
 in ac circuits, 1034–1036
 in alternating-current circuits, 1034–1036
 in mechanical systems, 453
- Resonance angular frequency, 1034
- Resonance curve, 518–519, 1035
- Resonance frequency, 1034
- Response curves, 1035
- Rest energy, 1240–1241
- Rest mass, 1237–1239
- Restoring force, 430
 in pendulum, 452
 in simple harmonic motion, 432
- Resultant, 11
- Reversed image, 1113
- Reversible processes, 643
- Right-handed system, 24
- Right-hand rule, 22, 919
- Rigid body, 272
- Rigid-body equilibrium, 338, 341–345
- Rigid-body rotation, 272–301. *See also*
 Rotation/rotational motion
 about moving axis, 308–314
 angular acceleration in, 276–279
 angular momentum and, 317–318
 angular velocity in, 273–277
 around fixed axis, 272–273
 dynamics of, 302–336
 kinetic energy in, 282–287
 linear acceleration in, 280–282
 linear speed in, 279–280
 moment of inertia and, 282–285
 with translational motion, 308–314
- R-L* circuits, 998–1002
 current decay in, 1001–1002
 current growth in, 999–1000
- rms speed, 591–592
- Rocket propulsion, 256–259
- Roller coasters, 84
- Rolling friction, 146, 313–314
- Rolling without slipping, 309–310
- Röntgen, Wilhelm, 1200
- Root-mean-square current, 1021–1022
- Root-mean-square speed, 591
- Root-mean-square values, 1021–1022
- Root-mean-square voltage, 863
- Rotational inertia, 283
- Rotational kinetic energy, 283, 308–309
- Rotation/rotational motion, 272–336
 about axis of symmetry, 317–318
 of American eels, 301
 angular acceleration and, 276–279,
 305–308
 angular momentum and, 316–321
 angular velocity and, 273–277
 with constant angular acceleration, 277–279
 coordinates for, 272–273
 direction of, 272–273, 302
 dynamics of, 302–336
 of earth, 414–415
 energy in, 282–287
 equilibrium and, 338
 fixed-axis, 272–273, 277
 of gyroscopes, 322–324
 linear acceleration in, 280–282
 linear speed in, 279–280
 moving-axis, 308–314
 Newton's second law of motion and, 306, 312
 power and, 314–316
 precession and, 322–324
 rigid-body, 272–301. *See also* Rigid-body rotation
 in rolling without slipping, 309–310
 torque and, 302–308
 with translational motion, 308–314
- units of, 273
 work and, 314–316
- Rotors, 902–903
- Rounding, significant figures in, 9
- Rule of Dulong and Petit, 555, 596–597
- Rutherford backscattering spectrometry, 780
- R value, 562
- S**
- Satellite orbits, 404–407
- Saturation magnetization, 940
- Scalar product, 19–21, 22
- Scalar quantities, 10
- Scattering of light, 1097–1098
- Schwarzschild, Karl, 416
- Schwarzschild radius, 416–417, 418
- Scientific notation, 9
- Scuba tank, air mass in, 583
- Second, 4
- Secondary rainbows, 1089–1090
- Secondary windings, 1036
- Second condition for equilibrium, 338
- Second law of thermodynamics, 642–677
 Carnot cycle and, 657–658
 Clausius statement of, 652
 engine statement of, 651
 entropy and, 664–665, 668
 Kelvin–Planck statement of, 651
 refrigerator statement of, 652
- Seconds, 4, 430
- Seconds per cycle, 430
- Sector velocity, 408
- Self-induced emf, inductive reactance and,
 1022–1024
- Self-inductance, 988, 992–995. *See also*
 Inductance
- Semiconductor(s), 904
 conduction in, 813–814
 diodes of, 821
 holes in, 813, 904
 metallic conduction and, 833–834
 moving charges in, 817–818
 resistivity of, 817
- Semi-major axis, 408, 409
- Series connection, 786–787
- Series motors, 903
- Series, resistors in, 844–845
- Shear modulus, 350
- Shear strain, 350
- Shear stress, 350
- Shielding, 746
- SHM. *See* Simple harmonic motion (SHM)
- Shock waves, 529–531
- Short circuit, 826–827, 864
- Shunt motors, 903
- Shunt resistors, 855
- Significant figures, 8–9
- Simple harmonic motion (SHM), 431–445.
See also Oscillation
 acceleration in, 435–437
 amplitude in, 434–435
 angular, 443
 applications of, 442–445
 circular motion and, 432–434
 displacement in, 435–437
 vs. electric oscillation, 1004
 energy in, 438–441
 equations of, 432–434
 as model of periodic motion, 432
 momentum in, 441
 period in, 434–435
 velocity in, 435–437
 vertical, 442–443
- Simple pendulum, 446–447
 vs. physical pendulum, 447
- Simultaneity, relativity of, 1220–1222
- Single-slit diffraction, 1187–1193. *See also*
 Diffraction
- Sinusoidal alternating emf, 1018
- Sinusoidal current, 1018–1021. *See also*
 Alternating current
- Sinusoidal electromagnetic waves, 1057–1061
- Sinusoidal waves, 466–467. *See also*
 Mechanical waves; Sound waves; Wave(s)
 electromagnetic, 1018–1021
 energy of, 478–481
 interference and, 1160
 particle velocity/acceleration in, 472–474
 reflected, 482
 standing, 483–490. *See also* Standing waves
 traveling, 483–485
 wave function for, 469–472
- SI units, 4, 5. *See also* Units of measure
- Sledding, Newton's first law and, 105
- Slidewire generators, 961–962
- Slug, 112
- Snell's law, 1081
- Snell, Willebrord, 1081
- Sodium doublet, 1199
- Solar radiation, 566–567
- Solenoids, 899, 901, 935–936
- Solid phase, 555–556
- Solids
 heat capacities of, 596–597
 molecules in, 586
 phases of, 599–602
 sound waves in, 507–508
 thermal conductivities of, 561t
- Sonar waves, 508
- Sound, 501
 infrasonic, 501
 loudness of, 505
 pitch of, 505
 resonance and, 518–520
 timbre of, 505
 ultrasonic, 501
- Sound intensity, 510–514
 decibel scale for, 512–513
 hearing loss and, 505, 506, 513
 inverse-square law for, 511
 representative values for, 513t
- Sound intensity level, 512, 513t
- Sound waves, 468, 501–540. *See also*
 Mechanical waves; Wave(s)
 audible range of, 501
 beats and, 522–524
 diffraction of, 1185, 1193
 direction of, 502
 displacement amplitude of, 502, 510–512
 Doppler effect and, 524–529, 1235–1236
 in fluid, 506–507
 frequency of, 505–506
 in gas, 508–509
 graphing of, 503
 harmonic content of, 488
 interference of, 520–522, 1184
 musical, 505–506
 perception of, 505–506
 pipe organs and, 516–520
 power of, 510–511
 pressure amplitude of, 503–504, 510–511
 as pressure fluctuations, 502–505
 shock, 529–531
 in solids, 507–508
 speed of, 468, 477, 506–509
 standing, 514–518
 superposition of, 482
 wind instruments and, 518
- Source of emf, 823
 internal resistance of, 825
 power input to, 829–830
 power output of, 829
- Source point, 691–692, 918–919
- South (S) pole, 879
- Space radiation shielding, 746
- Spacetime, 1232
- Spacetime coordinates, 1232

- Space travel, 409f, 427
 biological effects of, 409, 1226
 twin paradox and, 1226
 Spark plugs, 997
 Special theory of relativity, 87, 1217–1244.
 See also Relativity
 Specific gravity, 367
 Specific heat, 552–553
 molar, 554–555
 Spectrographs, grating, 1198–1200
 Spectroscopy, 1195
 Specular reflection, 1080, 1111
 Speed, 38
 air drag and, 146–148
 angular, 274, 279–280, 323
 average, 38
 of efflux, 380–381
 of electromagnetic waves, 1054, 1057–1058, 1068
 escape, 403, 406, 416
 instantaneous, 38
 of light, 4, 1051, 1078, 1218–1219, 1252
 measurement of, 4
 molecular, 591–592, 597–599
 orbital, 409
 of projectile, 75–76
 relativistic vs. nonrelativistic, 1224
 of rocket, 256–259
 root-mean-square (rms), 591–592
 of sound waves, 506–509
 supersonic, 530
 terminal, 147–149
 units of, 6
 vs. velocity, 38, 67, 280
 wave, 465–466, 467, 471, 474–478
 work and, 176–178
 of yo-yo, 310–311, 312
 Spheres
 electric field of, 730, 733
 electric flux through, 724
 gravitation and, 396–397
 mass distributions and, 411–413
 moment of inertia of, 285t, 290
 point charge inside, 725–726
 rolling, acceleration of, 313
 Spherical aberration, 1115, 1130
 Spherical mirrors
 concave, 1114–1118
 convex, 1118–1120
 extended objects in, 1116–1118
 focal point/length of, 1115–1116
 graphical methods for, 1120–1122
 image formation by, 1114–1122
 Spherical surface
 radius of curvature for, 1112, 1124
 reflection at, 1114–1122
 refraction at, 1122–1126
 Spherical symmetry
 gravitation and, 396–397
 Spiderweb conductivity, 843
 Spin, electron, 938
 Spring balance, 101
 Spring constant, 182
 Spring(s)
 elastic potential energy of, 210–215
 ideal, 430
 oscillation in, 429–430. *See also* Oscillation
 simple harmonic motion in, 431–445. *See also*
 Simple harmonic motion (SHM)
 work done on/by, 181–183
 SQUIDS (superconducting quantum interference devices), 976
 Stable equilibrium, 222
 Stable isotope ratio analysis (SIRA), 913
 Standards, reference, 4
 Standing waves, 483–490
 complex, 488
 electromagnetic, 1050, 1066–1069
 on fixed string, 486–490
 frequencies of, 486
 harmonics and, 487
 interference and, 484, 1160, 1162
 nodes and antinodes and, 483
 sound, 514–518
 string instruments and, 488–489
 Stars
 binary, 418, 1251
 mass of, 1251
 supernova, 1224
 systems of, 399, 418
 State(s), of matter, 555–556
 State variables, 579
 Static charge distribution, 752
 Static equilibrium, 338
 Static friction force, 142–143
 Steady flow, 376
 Steam heat, 558
 Stefan–Boltzmann constant, 565
 Stefan–Boltzmann law, 565
 Step-down transformer, 1037f
 Step-up transformer, 1036f, 1037
 Stick-slip phenomenon, 143
 Straight-line motion, 34–65
 with average acceleration, 40–41
 with average velocity, 34–37
 with changing acceleration, 53–55
 with constant acceleration, 44–49
 displacement and, 34–37
 finding position and velocity by integration, 53–55
 of freely falling bodies, 50–53
 graphing of, 36–37, 39–40, 42–44
 with instantaneous acceleration, 42–44
 instantaneous velocity and, 37–40
 motion diagrams of, 39–40
 relative velocity and, 85
 time and, 34–37
 work–energy theorem for, 181–185
 Strain, 345–351
 bulk, 348–349
 compressive, 347
 deformation and, 345–351
 elasticity and, 351
 elastic modulus and, 345–351
 shear, 350
 stress, 350
 tensile, 346–348
 volume, 348–349
 Streamline, 376, 700
 Strength
 tensile, 351–352
 ultimate, 351–352
 Stress, 345–351
 breaking, 351–352, 352t
 bulk, 348–349
 compressive, 347
 deformation and, 345–351
 elasticity and, 351–352
 elastic modulus and, 345–351
 shear, 350
 tensile, 346–348, 550–551
 thermal, 550–551
 units of, 346
 volume, 348–349
 Stress-strain diagram, 351f
 String instruments, standing waves and, 488–489
 String, standing waves on. *See* Standing waves
 Strong interactions, 155
 Strong nuclear force, 155, 680
 Sublimation, heat of, 558, 600
 Subtraction, of vectors, 11–14
 Sun, 997f
 Sunglasses, polarized, 1093, 1094
 Sunsets, 1097
 Superconducting quantum interference devices (SQUIDS), 976
 Superconductivity, 818, 975–976
 Superconductors, 964
 type-I, 975–976
 type-II, 976
 Supercooling, 558
 Superheating, 558
 Supermassive black holes, 418
 Supernovas, 1224
 Superposition
 of electric fields, 695
 of forces, 101–103, 398, 687
 of magnetic fields, 921
 principle of, 482, 687, 695, 1159–1162
 of waves, 482, 488
 Supersonic speed, 530
 Surface charge density, 695
 Surface integral, 723
 Surface tension, 375
 Symmetry properties, of systems, 729
 Systems
 isolated, 241
 symmetry properties of, 729
T
 Tangential acceleration, 84
 Tangential component of acceleration, 280
 Target variables, 3
 Telephoto lens, 1136, 1138
 Telescopes, 1115
 Gran Telescopio Canarias, 1147
 Hubble Space Telescope, 1115f, 1174, 1212
 optical system of, 1145–1147
 resolving power of, 1204
 Very Large Array, 1203
 Temperature, 541–551
 absolute, 509
 body, 544f, 571
 boiling, 556–557
 critical, 585, 975
 gas pressure and, 546
 vs. heat, 552
 internal energy and, 625
 macroscopic definition of, 542
 melting, 556–557
 molecular kinetic energy and, 625
 resistivity and, 817–818
 as scalar quantity, 10
 vs. temperature interval, 544
 thermal equilibrium and, 541–543
 units of measure for, 543, 546
 Temperature coefficient of resistivity, 817–818, 820
 Temperature gradient, 561
 Temperature interval, 544
 Temperature scale(s), 543–544
 absolute, 546, 658–659
 Celsius, 542
 conversion between, 544
 Fahrenheit, 544
 ideal-gas, 545, 546, 659
 Kelvin, 545–546, 655, 658–659
 Temporal artery thermometer, 544
 Tendons are nonideal springs, 184
 Tensile strain, 346–348
 elasticity and, 351–352
 plasticity and, 351–352
 Tensile strength, 351–352
 Tensile stress, 346–348
 elasticity and, 351–352
 plasticity and, 351–352
 thermal stress and, 550–551
 Tension, 346, 347
 Newton's third law and, 118
 static friction and, 143
 surface, 375
 Tension force, 101
 Terminal speed, 147–149
 Terminal voltage, 824
 Tesla, 882
 Tesla coils, 990–991

I-16 Index

- Tesla, Nikola, 882
- Test charge
- for electric fields, 690–691
 - for magnetic fields, 882–883
- Test mass, 691
- Theory, 2
- range of validity of, 2
- Theory of Everything (TOE), 156
- Theory of relativity. *See* Relativity
- Thermal conductivity, 561–562, 817
- Thermal conductors, 542
- Thermal efficiency, of heat engine, 645, 648
- Thermal equilibrium, 541–543
- Thermal expansion, 547–551
- linear, 547–558
 - in object with hole, 548
 - volume, 548–549
 - of water, 550
- Thermal properties of matter, 579–612
- Thermal radiation, 1078
- Thermal resistance, 562
- Thermal stress, 550–551
- Thermistors, 818
- Thermodynamic processes, 614
- adiabatic, 624, 629–631, 653
 - directions of, 642–643
 - disorder in, 643
 - equilibrium in, 643
 - heat added in, 617–618
 - in heat engines, 644–647
 - infinitesimal changes of state in, 623
 - intermediate states (paths) in, 617–618
 - isobaric, 624
 - isochoric, 624
 - isothermal, 624–625, 653
 - reversibility of, 642–643, 653
 - types of, 623–625
 - work done in, 614
- Thermodynamics, 541
- applications of, 614
 - first law of, 613–641. *See also* First law of thermodynamics
 - second law of, 642–677. *See also* Second law of thermodynamics
 - sign rules for, 614
 - third law of, 659
 - zeroth law of, 542–543
- Thermodynamic systems, 613–614
- human body as, 619
 - internal energy of. *See* Internal energy
 - paths in, 617–618
 - work done in, 614
- Thermometers, 543–544
- bimetallic strip, 543
 - gas, 545–546, 582, 659
 - resistance, 545
 - temporal artery, 544
- Thermos bottles, 566
- Thick-film interference, 1171–1172
- Thin-film interference, 1170–1175
- Thin-lens equation, 1129–1130
- Thin lenses, 1127–1135
- converging, 1127–1129
 - diverging, 1129
 - focal length of, 1127, 1129–1131
 - focal point of, 1127
 - graphical methods for, 1131–1132
 - image formation by, 1131–1134
 - index of refraction of, 1129–1130
 - positive, 1127
 - properties of, 1127
 - radius of curvature for, 1129–1130
- Third law of thermodynamics, 659
- Thomson, J. J., 744, 891
- Thomson's atomic model, 744
- Thomson's e/m experiment, 891–892
- Thought experiments, 1220–1222
- Three-wire line, 865
- Tidal forces, 418
- Timbre, 505
- Time
- mean free, 832
 - power and, 187
 - proper, 1223–1225, 1226
 - spacetime and, 1233
 - straight-line motion and, 34–37
 - units of, 4, 5t, 6
- Time constant, for circuit, 860–861, 1000
- Time dilation, 418, 1223–1226
- Time intervals
- measurement of, 1224
 - relativity of, 1222–1227
- Toroidal solenoid, magnetic field of, 936
- Torque, 302–308
- angular acceleration and, 305–308
 - angular displacement and, 314–316
 - angular momentum and, 316, 317–318
 - application of, 304–305
 - calculation of, 303
 - center of mass and, 306
 - constant, 314–315
 - on current loops, 897–902
 - direction of, 302
 - on electric dipole, 899
 - equilibrium and, 338
 - friction and, 313
 - gravitational, 339–340
 - of internal vs. external forces, 306
 - kinetic energy and, 315
 - magnetic, 897–902
 - magnitude of, 304
 - measurement of, 303
 - moment of inertia and, 305–306
 - net, 315
 - net force and, 305, 316, 317
 - positive vs. negative, 303
 - unit of, 303
 - as vector, 304–305
 - weight and, 306
 - work done by, 314–316
- Torr, 372
- Torricelli's theorem, 381
- Torsion balance, 397
- Torsion constant, 443
- Total energy, 171, 1240
- Total internal reflection, 1085–1088
- Total mechanical energy, 203–204
- Total momentum, 241–242, 254
- Total work, 175, 238
- Tractive resistance, 146
- Traffic light sensors, 994
- Trajectory, 74–76, 701
- Trampolines, 212
- Transcranial magnetic stimulation, 954f, 987
- Transformers, 990, 1036–1039
- Translational motion, 302
- molecular kinetic energy and, 595
 - with rotational motion, 308–314
 - vibrational, 595–596
- Transmission gratings, 1197
- Transverse waves, 465. *See also* Mechanical waves; Wave(s)
- electromagnetic, 1053, 1057
 - periodic, 466–468
 - speed of, 474–478
 - wave function for, 472–474
- Traveling waves, 483–485
- interference of, 484, 520
- Triple point, 600
- True weight, 414
- Tsunamis, 1210
- Tug-of-war, 254–255, 365
- Tuning forks, 434
- Turbulent flow, 376, 383
- Twin paradox, 1226
- Tyrannosaurus rex, physical pendulum and, 449

U

- Ultimate strength, 351–352
- Ultrasonic sound, 501
- Ultrasound, 508, 530, 540
- Uncertainty
- fractional (percent), 8
 - in measurement, 8
- Underdamping, 450, 1007–1008
- Uniform circular motion, 81–83, 149–154
- centripetal acceleration and, 82–83
 - dynamics of, 149–154
 - net force and, 108
 - vs. nonuniform circular motion, 84
 - period of, 83
 - vs. projectile motion, 83
 - in vertical circle, 153–154
- Units of measure, 3–7. *See also* Measurement
- for acceleration, 112
 - for amplitude, 430
 - for angular frequency, 430–431
 - for angular velocity, 273–274
 - in British system, 5–6, 5t
 - in calculations, 6–7
 - for capacitance, 782
 - in cgs metric system, 112
 - consistency for, 6
 - conversion of, 6–7
 - derived, 5
 - dimensionally consistent, 6
 - for electric current, 686–687, 814
 - for electric field, 691, 756–757
 - for electric force, 686–687
 - for electric potential, 754, 756–757
 - for electromagnetic waves, 1050
 - in equations, 6–7
 - for force, 5, 101, 112
 - for frequency, 430
 - for heat, 552, 553
 - for kinetic energy, 177
 - for length, 4, 5
 - for magnetic flux, 886
 - for magnetic force, 882
 - for mass, 4, 108–109, 112
 - for momentum, 236
 - for mutual inductance, 990–991
 - for period, 430
 - prefixes for, 5
 - for pressure, 348, 368
 - for resistance, 820
 - significant figures and, 8–9
 - system, 4, 5–6
 - for temperature, 543, 546
 - for time, 4, 5t, 6
 - for torque, 303
 - uncertainty and, 8
 - for velocity, 37
 - for volume, 7
 - for weight, 114
 - for work, 172
- Unit vectors, 18–19
- Universal conservation laws, 681
- Unpolarized light, 1091
- Unstable equilibrium, 222
- U.S. National Institute of Standards and Technology, 5

V

- Vacancies, 813
- Vacuum
- capacitors in, 782–784, 791

- electric-field energy in, 791
 - electric fields in, 791
 - permittivity of, 796
 - Vacuum bottles, 566
 - Validity, range of, 2
 - Van Allen radiation belts, 889
 - Van de Graaff electrostatic generator, 735, 761
 - van der Waals equation, 584
 - van der Waals interactions, 444, 584
 - Vaporization, 556
 - heat of, 556–558
 - Variables
 - state, 579
 - target, 3
 - Vector current density, 815
 - Vector field, 692
 - Vector product, 19, 22–24
 - Vector quantity, 10–11, 100–101
 - Vector(s), 10–24
 - acceleration, 34, 69–74, 277. *See also* Acceleration vectors
 - addition of, 11–14, 16–17
 - angular momentum, 317, 322
 - angular velocity, 275
 - antiparallel, 11
 - components of, 14–18, 20–21, 23–24, 102–103
 - direction of, 10, 14–16
 - displacement, 10–11, 17
 - force, 10, 101–103
 - force of, 10
 - magnitude of, 10, 12, 14–16
 - momentum, 236, 239
 - multiplication of, 16
 - negative of, 11
 - notation for, 11
 - parallel, 11
 - position, 67–69
 - Poynting, 1062–1064
 - products of, 19–24
 - right-hand rule for, 22, 919
 - subtraction of, 11–14
 - torque, 304–305
 - unit, 18–19
 - velocity, 34, 67–69, 275
 - Vector sum, 11, 16–17
 - Velocity
 - vs. acceleration, 40
 - angular, 273–277, 280
 - average, 34–37, 67–69. *See also* Average velocity
 - drift, 813, 814–816
 - as function of time
 - with changing acceleration, 53–55
 - with constant acceleration, 44–47
 - Galilean velocity transformation and, 87
 - graphing of, 36–37, 39–40
 - instantaneous, 37–40, 67–69. *See also* Instantaneous velocity
 - linear, 274, 276
 - Lorentz transformation for, 1232–1233
 - magnitudes of, 36t, 37. *See also* Speed
 - Newton's first law and, 103–107
 - of particle in wave, 472–474
 - phase, 471
 - power and, 188
 - of projectile, 74–76
 - relative, 84–89. *See also* Relative velocity
 - sector, 408
 - sign conventions for, 36, 43, 43t
 - in simple harmonic motion, 435–437
 - vs. speed, 38, 67, 280
 - speed and direction as, 10
 - straight-line motion and, 34–40, 53–55
 - units of, 37
 - Velocity selectors, 891
 - Velocity vectors, 34, 67–69, 275
 - Venturi meter, 381
 - Verne, Jules, 403
 - Vertex, of mirror, 1114
 - Vertical circle, uniform circular motion in, 153–154
 - Vertical simple harmonic motion, 442–443
 - Very Large Array Telescopes, 1203
 - Vibration, 430
 - heat capacities and, 595–596
 - molecular, 443–445, 586, 595–596
 - Virtual focal point, 1119
 - Virtual image, 1111, 1132
 - Virtual object, 1132
 - Viscosity, 376, 382–384
 - Visible light, 1051
 - Vision
 - in animals, 1098f, 1109, 1139, 1142, 1158, 1205f
 - defects in, 1139–1142
 - normal, 1139
 - polarized light and, 1109
 - ultraviolet, 1052
 - Volt
 - electron, 756–757
 - unit of potential, 754
 - Voltage, 754. *See also* Potential difference
 - in ac circuits, 1034
 - capacitor, 1024–1026
 - current and, 820–821
 - Hall, 904–905
 - household, 863
 - inductor, 1022–1023
 - measurement of, 854–858
 - resistor, 1022, 1026
 - root-mean-square, 863
 - sinusoidal, 1019
 - terminal, 824
 - transformers and, 1036–1039
 - Voltage amplitude, 1019, 1022
 - Voltmeters, 754, 825, 826, 855–857
 - ammeters and, 856–857, 1021
 - digital, 857
 - Volt per meter, 756
 - Volume
 - density and, 366–368
 - equipotential, 766–767
 - of gas, 582
 - units of, 7
 - Volume change, work done during, 585, 615–616
 - Volume charge density, 695
 - Volume expansion, 548–549
 - Volume strain, 348–349
 - Volume stress, 348–349
 - v_x - t graphs, 42–44, 44t
- W**
- Water
 - supercooled, 558
 - thermal expansion of, 550
 - Water pressure, in home, 380
 - Water waves, interference and, 1163
 - Watt, 188
 - Watt, James, 188
 - Wave equation, 473, 477
 - for electromagnetic waves, 1055–1057
 - for mechanical waves, 473, 477
 - Wave fronts, 1078–1079
 - Wave function, 469
 - additive property of, 482
 - for electromagnetic waves, 1058, 1059
 - graphing of, 470–471
 - for longitudinal waves, 473
 - for mechanical waves, 469–472
 - for sinusoidal waves, 469–472
 - superposition principle and, 482
 - for transverse waves, 472–474
 - Wave intensity, 480
 - Wave interference. *See* Interference
 - Wavelengths, 467
 - frequency and, 1058
 - of light, 1051–1052
 - measurement of, 1175–1176
 - Wave motion vs. particle motion, 467
 - Wave number, 470, 1058
 - Wave–particle duality, 1078
 - index of refraction and, 1083
 - Maxwell's wave theory and, 1048–1052
 - Wave phase, 471
 - Wave pulse, 466
 - Wave(s)
 - coherent, 1160
 - electromagnetic, 1048–1076
 - light as. *See* Wave–particle duality
 - mechanical, 464–500. *See also* Mechanical waves
 - medium for, 464–465
 - polarization of. *See* Polarization
 - radio, 1049, 1051
 - reflected, 481
 - shock, 529–531
 - sonar, 508
 - sound, 468, 501–540, 1184
 - transverse, 1053, 1057. *See also* Transverse waves
 - on vocal folds, 499–500
 - Wave speed, 465–466, 467, 471
 - calculation of, 477–478
 - impulse–momentum theory and, 475–476
 - vs. particle speed, 510
 - on a string, 474–477
 - for transverse waves, 474–478
 - Wave velocity vs. particle velocity, 510
 - Weak interactions, 155–156
 - Weber, 886
 - Weber, Wilhelm, 886
 - Weight, 101
 - acceleration and, 109, 113–114
 - apparent, 138–141, 414–415
 - equilibrium and, 338–341
 - gravitation and, 399–402
 - mass and, 109, 113–115
 - measurement of, 114
 - molecular, 554, 580
 - torque and, 306
 - true, 414
 - Weightlessness
 - apparent, 138–141, 405
 - true, 405
 - Weight lifting, equilibrium and, 344
 - Westinghouse, George, 1018
 - Wheatstone Bridge, 875
 - Wide-angle lens, 1136
 - Windings, 1036–1037
 - Wind instruments, 518
 - Windows, insulating, 612
 - Wire(s)
 - Ampere's law for, 930–936
 - interaction force between, 926–927
 - magnetic field of, 923–926, 930–932, 935
 - Wiring systems
 - automobile, 863, 865
 - household, 863–866, 1037
 - Work, 171–200
 - change in speed and, 176–178
 - in composite systems, 181–182
 - displacement and, 172–175, 176–178
 - done by conservative force, 748
 - done by constant forces, 172–173
 - done by electric fields, 748–753. *See also* Electric potential energy
 - done by electric force, 749–750, 754
 - done by electric potential, 754
 - done by electromotive force, 823–824
 - done by fluid flow, 378–381
 - done by gravitation, 402–404
 - done by torque, 314–316
 - done by varying forces, 181–187
 - done by waves, 478–480
 - done by working substance, 645
 - done during heat transfer, 651

Work (*continued*)
 done during volume change, 585, 615–616
 done in thermodynamic process, 617
 done in thermodynamic system, 614
 done on/by springs, 181–183
 done to charge capacitor, 790–793
 kinetic energy and, 176–181
 negative, 173–175, 177
 positive, 173–175, 177
 potential energy and, 748
 power and, 187–189, 478–480
 rate of, 187–189
 relativistic kinetic energy and, 1239–1242
 as scalar quantity, 173
 sign rules for, 173–175, 177, 614
 total, 175, 238
 units of, 172
 zero, 173–175
Work–energy theorem, 176–181, 748
 for composite systems, 181–182
 for motion along curve, 185–187
 for straight-line motion, 181–185
 for varying forces, 181–187
Working substance, 644, 645
Workless refrigerators, 651

X
X-ray diffraction, 1199–1203
X rays, 898, 1052, 1200. *See also* Radiation
 x - t graphs, 36–37
 velocity on, 39–40, 44t

Y
Yeager, Chuck, 530
Young’s interference experiment,
 1163–1165, 1175
Young’s modulus, 346–347
Young, Thomas, 1163

Z
Zero, absolute, 659
Zero height, 204
Zero net force, 104–105, 106–107
Zeroth law of thermodynamics, 542–543
Zero work, 173–175
Z machine, 791
Zoom lenses, 1137

Astronomical Data[†]

Body	Mass (kg)	Radius (m)	Orbit radius (m)	Orbital period
Sun	1.99×10^{30}	6.96×10^8	—	—
Moon	7.35×10^{22}	1.74×10^6	3.84×10^8	27.3 d
Mercury	3.30×10^{23}	2.44×10^6	5.79×10^{10}	88.0 d
Venus	4.87×10^{24}	6.05×10^6	1.08×10^{11}	224.7 d
Earth	5.97×10^{24}	6.37×10^6	1.50×10^{11}	365.3 d
Mars	6.42×10^{23}	3.39×10^6	2.28×10^{11}	687.0 d
Jupiter	1.90×10^{27}	6.99×10^7	7.78×10^{11}	11.86 y
Saturn	5.68×10^{26}	5.82×10^7	1.43×10^{12}	29.45 y
Uranus	8.68×10^{25}	2.54×10^7	2.87×10^{12}	84.02 y
Neptune	1.02×10^{26}	2.46×10^7	4.50×10^{12}	164.8 y
Pluto [‡]	1.30×10^{22}	1.19×10^6	5.91×10^{12}	248.0 y

[†]Source: NASA (<http://solarsystem.nasa.gov/planets/>). For each body, “radius” is its average radius and “orbit radius” is its average distance from the sun or (for the moon) from the earth.

[‡]In August 2006, the International Astronomical Union reclassified Pluto and similar small objects that orbit the sun as “dwarf planets.”

Prefixes for Powers of 10

Power of ten	Prefix	Abbreviation	Pronunciation
10^{-24}	yocto-	y	<i>yoc</i> -toe
10^{-21}	zepto-	z	<i>zep</i> -toe
10^{-18}	atto-	a	<i>at</i> -toe
10^{-15}	femto-	f	<i>fem</i> -toe
10^{-12}	pico-	p	<i>pee</i> -koe
10^{-9}	nano-	n	<i>nan</i> -oe
10^{-6}	micro-	μ	<i>my</i> -crow
10^{-3}	milli-	m	<i>mil</i> -i
10^{-2}	centi-	c	<i>cen</i> -ti
10^3	kilo-	k	<i>kil</i> -oe
10^6	mega-	M	<i>meg</i> -a
10^9	giga-	G	<i>jig</i> -a or <i>gig</i> -a
10^{12}	tera-	T	<i>ter</i> -a
10^{15}	peta-	P	<i>pet</i> -a
10^{18}	exa-	E	<i>ex</i> -a
10^{21}	zetta-	Z	<i>zet</i> -a
10^{24}	yotta-	Y	<i>yot</i> -a

Examples:

1 femtometer = 1 fm = 10^{-15} m
1 picosecond = 1 ps = 10^{-12} s
1 nanocoulomb = 1 nC = 10^{-9} C
1 microkelvin = 1 μ K = 10^{-6} K

1 millivolt = 1 mV = 10^{-3} V
1 kilopascal = 1 kPa = 10^3 Pa
1 megawatt = 1 MW = 10^6 W
1 gigahertz = 1 GHz = 10^9 Hz

Fundamental Physical Constants*

Name	Symbol	Value
Speed of light in vacuum	c	$2.99792458 \times 10^8 \text{ m/s}$
Magnitude of charge of electron	e	$1.602176634 \times 10^{-19} \text{ C}$
Gravitational constant	G	$6.67408(31) \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2$
Planck's constant	h	$6.62607015 \times 10^{-34} \text{ J} \cdot \text{s}$
Boltzmann constant	k	$1.380649 \times 10^{-23} \text{ J/K}$
Avogadro's number	N_A	$6.02214076 \times 10^{23} \text{ molecules/mol}$
Gas constant	R	$8.314462618 \dots \text{ J/mol} \cdot \text{K}$
Mass of electron	m_e	$9.10938356(11) \times 10^{-31} \text{ kg}$
Mass of proton	m_p	$1.672621898(21) \times 10^{-27} \text{ kg}$
Mass of neutron	m_n	$1.674927471(21) \times 10^{-27} \text{ kg}$
Magnetic constant	μ_0	$1.25663706 \times 10^{-6} \text{ Wb/A} \cdot \text{m}$ (approximate) $\cong 4\pi \times 10^{-7} \text{ Wb/A} \cdot \text{m}$
Electric constant	$\epsilon_0 = 1/\mu_0 c^2$	$8.854187817 \times 10^{-12} \text{ C}^2/\text{N} \cdot \text{m}^2$ (approximate)
	$1/4\pi\epsilon_0$	$8.987551787 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$ (approximate)

Other Useful Constants*

Mechanical equivalent of heat		4.186 J/cal (15° calorie)
Standard atmospheric pressure	1 atm	$1.01325 \times 10^5 \text{ Pa}$
Absolute zero	0 K	-273.15°C
Electron volt	1 eV	$1.6021766209(98) \times 10^{-19} \text{ J}$
Atomic mass unit	1 u	$1.660539040(20) \times 10^{-27} \text{ kg}$
Electron rest energy	$m_e c^2$	0.5109989461(31) MeV
Volume of ideal gas (0°C and 1 atm)		22.413962(13) liter/mol
Acceleration due to gravity (standard)	g	9.80665 m/s^2

*Source: National Institute of Standards and Technology (<http://physics.nist.gov/cuu>). Numbers in parentheses show the uncertainty in the final digits of the main number; for example, the number 1.6454(21) means 1.6454 ± 0.0021 . Values shown without uncertainties are exact. The exact values of the magnitude of the charge of the electron, Planck's constant, the Boltzmann constant, Avogadro's number, and the gas constant are from the redefinitions adopted in 2018. As consequences of these redefinitions, the values of the magnetic constant and electric constant now have fractional uncertainties of about 2×10^{-10} . As of this writing (2018) it was expected that updated values of the magnetic and electric constants, as well as of all other constants with uncertainties, were to be announced in May 2019. These updated values will be available at <http://physics.nist.gov/cuu>.