Infection Prevention and Control

HAND HYGIENE

Clean hands at the required moments using the correct technique.

PPE

Select and remove personal protective equipment according to exposure risk.

SHARPS

Use safe handling and dispose of sharps immediately in approved containers.

ISOLATION

Follow transmission-based precautions and posted instructions.

EXPOSURE

Wash or flush the area, report immediately and follow occupational health guidance.

ENVIRONMENT

Maintain clean equipment, safe waste handling and respiratory hygiene.

Healthcare-Associated Infections

COMMON ROUTES

Hands, equipment, devices, droplets, aerosols and contaminated environments may transmit infection.

PATIENT RISK

Invasive devices, surgery, immune suppression and prolonged stays can increase risk.

PREVENTION

Reliable hand hygiene, aseptic technique and device-care practices reduce avoidable infection.

Break the chain of infection at every clinical contact.

Hand Hygiene

ALCOHOL HAND RUB

Use on visibly clean hands and rub all surfaces until dry.

SOAP AND WATER

Use when hands are visibly soiled and when the local policy specifically requires it.

TECHNIQUE

Cover palms, backs of hands, between fingers, thumbs, fingertips and wrists.






Perform hand hygiene at the required moments before and after patient contact and aseptic tasks.

How to Handwash?

WASH HANDS WHEN VISIBLY SOILED! OTHERWISE, USE HANDRUB

Duration of the entire procedure: 40-60 seconds

0



Wet hands with water;

1



Apply enough soap to cover all hand surfaces;

2



Rub hands palm to palm;

3



Right palm over left dorsum with interlaced fingers and vice versa;

4



Palm to palm with fingers interlaced;

5



Backs of fingers to opposing palms with fingers interlocked;

6



Rotational rubbing of left thumb clasped in right palm and vice versa;

7



Rotational rubbing, backwards and forwards with clasped fingers of right hand in left palm and vice versa;

8




Rinse hands with water;

9



Dry hands thoroughly with a single use towel;

10



Use towel to turn off faucet;

11



Your hands are now safe.

World Health Organization | Patient Safety — A World Alliance for Safer Health Care | SAVE LIVES Clean Your Hands

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this document. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

WHO acknowledges the Hôpitaux Universitaires de Geneva (HUG), in particular the members of its Infection Control Programme, for their active participation in developing this material.

May 2009

How to Handrub?

RUB HANDS FOR HAND HYGIENE! WASH HANDS WHEN VISIBLY SOILED

Duration of the entire procedure: 20-30 seconds

1a



1b



Apply a palmful of the product in a cupped hand, covering all surfaces;

2



Rub hands palm to palm;

3



Right palm over left dorsum with interlaced fingers and vice versa;

4



Palm to palm with fingers interlaced;

5



Backs of fingers to opposing palms with fingers interlocked;

6




Rotational rubbing of left thumb clasped in right palm and vice versa;

7



Rotational rubbing, backwards and forwards with clasped fingers of right hand in left palm and vice versa;

8



Once dry, your hands are safe.


World Health Organization | Patient Safety — A World Alliance for Safer Health Care | SAVE LIVES Clean Your Hands

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this document. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.
WHO acknowledges the Hôpitaux Universitaires de Genève (HUG), in particular the members of the Infection Control Programme, for their active participation in developing this material.

May 2009

Your 5 Moments for Hand Hygiene

1. BEFORE TOUCHING A PATIENT
2. BEFORE CLEAN / ASEPTIC PROCEDURE
3. AFTER BODY FLUID EXPOSURE RISK
4. AFTER TOUCHING A PATIENT
5. AFTER TOUCHING PATIENT SURROUNDINGS



Standard Precautions

HAND HYGIENE

Clean hands at the correct time using the correct method.

RISK-BASED PPE

Select gloves, gown, mask and eye protection for the expected exposure.

RESPIRATORY HYGIENE

Apply source control and cough etiquette.

ASEPTIC PRACTICE

Protect key sites and key parts during procedures.

SAFE EQUIPMENT

Clean, disinfect or sterilize reusable equipment correctly.

WASTE AND LINEN

Handle sharps, waste, specimens and linen without exposing others.

Transmission-Based Precautions

CONTACT

Hand hygiene, gloves and gown as indicated. Dedicate or disinfect equipment.

DROPLET

Wear a medical mask and follow patient placement and transport precautions.

AIRBORNE

Use the required respirator, complete a fit check and follow airborne room requirements.

Check the isolation sign before entry and ask Infection Control when requirements are unclear.

CONTACT PRECAUTIONS



Visitors: Report to nurse before entering

Visitantes: Favor de notificar a la(el) enfermera(o) de su presencia antes de entrar a la habitación. Gracias.

ممنوع دخول هذه الغرفة بدون إذن من الممرضة

(In addition to Standard Precautions)

ALL STAFF: On entering the room, you must wear:		YES	NO
	**Gown** (upon contact with patient or environment)	✓	
	**Gloves**	✓	
	**Alcohol Hand Sanitizer**	✓	
	**Surgical Mask** (use as per standard precautions)		X

- PRIVATE ROOM Recommended – Cohort if private room not available.
- Utilize dedicated equipment (i.e., stethoscope, blood pressure cuff, etc.) or clean between patient use.
- Remove gown and gloves before leaving the patient environment.
- Hand hygiene must be performed immediately after gloves removed.
  - Utilize alcohol hand gel if hands are not visibly soiled.
  - Wash with antimicrobial soap and water for at least 15 seconds if hands are visibly soiled.

During Transport: Cover wound and drainage area.

DROPLET PRECAUTIONS



Visitors: Report to nurse before entering

Visitantes: Favor de notificar a la(el) enfermera(o) de su presencia antes de entrar a la habitación. Gracias.

ممنوع دخول هذه الغرفة بدون إذن من الممرضة

(In addition to Standard Precautions)

ALL STAFF: On entering the room, you must wear:		YES	NO
	**Surgical Mask**	✓	
	**Alcohol Hand Sanitizer**	✓	
	**Gown** (use as per standard precautions)		X
	**Gloves** (use as per standard precautions)		X

- PRIVATE ROOM Recommended – Cohort if private room not available.
- Hand hygiene must be performed immediately after gloves removed.
  - Utilize alcohol hand gel if hands are not visibly soiled.
  - Wash with antimicrobial soap and water for at least 15 seconds if hands are visibly soiled.
- Visitors Wear Surgical Mask

During Transport: Patient wears surgical mask.

- *(Vent dependent patients should have filter on ambu bag)*

AIRBORNE PRECAUTIONS

STOP

Visitors: Report to nurse before entering

Visitantes: Favor de notificar a la(el) enfermera(o) de su presencia antes de entrar a la habitación. Gracias.

ممنوع دخول هذه الغرفة بدون إذن من الممرضة

(In addition to Standard Precautions)

ALL STAFF: On entering the room, you must wear:		YES	NO
	**N95 Mask or P.A.P.R.** (Powered Air-Purifying Respirator)	✓	
	**Alcohol Hand Sanitizer**	✓	
	**Gown** (use as per standard precautions)		X
	**Gloves** (use as per standard precautions)		X

- Negative Pressure Room Required.
- Keep room door closed.
- Visitors must wear a mask.
- Non-immune persons should not enter the room of patients with known or suspected measles or chicken pox (varicella).
- Hand hygiene must be performed immediately after gloves removed.

During Transport: Patient wears surgical mask.
(Vent dependent patients should have filter on ambu bag)

Sharps and Occupational Exposure

- Encourage bleeding gently and wash the exposed skin with soap and water. Flush mucous membranes with water.
- Do not scrub the wound or use unsafe chemicals.
- Report the exposure immediately to the supervisor and Occupational Health.
- Follow urgent risk assessment, baseline testing and post-exposure prophylaxis when indicated.
- Document the incident and attend all required follow-up appointments.

Sharps Items

- Needles should NEVER be re – capped.
- All sharp items are placed in a puncture – proof , biohazard-labeled, sharps container.
- Containers are not to be over – fill sharps containers must be less than 3/4

