

References

- [1] Shion Chen, Yuki Nakaguchi, and Sachio Komamiya. Testing Bell's Inequality using Charmonium Decays. 2013.
- [2] Shion Chen. Testing bell's inequality in charm factory. In *IHEP seminar*, 2013.
- [3] Shion Chen. Testing quantum entanglement in collider experiments. In *KEK theory seminar*, 2013.
- [4] Shion Chen. Test of Bell's Inequality and Entanglement Measurement in Collider Experiments. Master's thesis, The University of Tokyo, 2014.
- [5] Shion Chen. Testing quantum mechanics in collider experiments. In *EM-FCSC International School of Sub-nuclear Physics 2015*, 2015.
- [6] Shion. Chen et al. Test beam analysis of SiW ECAL physics prototype in 2011 FNAL. In *CALICE Collaboration Meeting, LAPP, Annecy, France.*, 2013.
- [7] Shion. Chen et al. Test of Power Pulsing with the HBU-LED System. In *CALICE Collaboration Meeting, LAPP, Annecy, France.*, 2013.
- [8] Shion. Chen et al. Simulation Study on SiW ECAL optimization. In *CALICE Collaboration Meeting, LAPP, Annecy, France.*, 2013.
- [9] Tatsuhiko Tomita et al. A study of silicon sensor for ILD ECAL. In *International Workshop on Future Linear Colliders (LCWS13) Tokyo, Japan, November 11-15, 2013*, 2014.
- [10] M. S. Amjad et al. Beam test performance of the SKIROC2 ASIC. *Nucl. Instrum. Meth.*, A778:78–84, 2015.
- [11] Chihiro Kozakai, Shion Chen, Daniel Jeans, Yoshio Kamiya, and Sachio Komamiya. Robustness of a SiECAL used in Particle Flow Reconstruction. In *International Workshop on Future Linear Colliders (LCWS13) Tokyo, Japan, November 11-15, 2013*, 2014.
- [12] M. Chefdeville et al. Shower development of particles with momenta from 15 GeV to 150 GeV in the CALICE scintillator-tungsten hadronic calorimeter. *JINST*, 10(12):P12006, 2015.
- [13] Y. Sudo et al. Development of a Highly Granular Silicon-Tungsten ECAL for the ILD. *Nucl. Part. Phys. Proc.*, 273-275:2554–2556, 2016.
- [14] The ATLAS collaboration. Search for squarks and gluinos in events with an isolated lepton, jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector. 2016.

- [15] The ATLAS collaboration. Search for supersymmetry with two and three leptons and missing transverse momentum in the final state at $\sqrt{s} = 13$ TeV with the ATLAS detector. 2016.
- [16] Morad Aaboud et al. Measurement of $W^{\pm}W^{\pm}$ vector-boson scattering and limits on anomalous quartic gauge couplings with the ATLAS detector. 2016.
- [17] Morad Aaboud et al. Measurement of jet activity produced in top-quark events with an electron, a muon and two b -tagged jets in the final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. 2016.
- [18] Morad Aaboud et al. Measurements of $\psi(2S)$ and $X(3872) \rightarrow J/\psi\pi^+\pi^-$ production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. 2016.
- [19] Morad Aaboud et al. Measurements of charge and CP asymmetries in b -hadron decays using top-quark events collected by the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV. 2016.
- [20] Morad Aaboud et al. Measurement of the ZZ production cross section in pp collisions at $\sqrt{s} = 8$ TeV using the $ZZ \rightarrow \ell^-\ell^+\ell'^-\ell'^+$ and $ZZ \rightarrow \ell^-\ell^+\nu\bar{\nu}$ channels with the ATLAS detector. 2016.
- [21] Morad Aaboud et al. Search for triboson $W^{\pm}W^{\pm}W^{\mp}$ production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. 2016.
- [22] Georges Aad et al. Performance of algorithms that reconstruct missing transverse momentum in $\sqrt{s} = 8$ TeV proton-proton collisions in the ATLAS detector. 2016.
- [23] Morad Aaboud et al. Measurement of W boson angular distributions in events with high transverse momentum jets at $\sqrt{s} = 8$ TeV using the ATLAS detector. 2016.
- [24] Morad Aaboud et al. Measurements of long-range azimuthal anisotropies and associated Fourier coefficients for pp collisions at $\sqrt{s} = 5.02$ and 13 TeV and p +Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector. 2016.
- [25] Morad Aaboud et al. Search for anomalous electroweak production of WW/WZ in association with a high-mass dijet system in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. 2016.
- [26] Morad Aaboud et al. Search for dark matter in association with a Higgs boson decaying to b -quarks in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. 2016.
- [27] Morad Aaboud et al. A measurement of material in the ATLAS tracker using secondary hadronic interactions in 7 TeV pp collisions. 2016.

- [28] Morad Aaboud et al. Measurement of the inclusive cross-sections of single top-quark and top-antiquark t -channel production in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. 2016.
- [29] Morad Aaboud et al. Measurement of the $t\bar{t}Z$ and $t\bar{t}W$ production cross sections in multilepton final states using 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. 2016.
- [30] Morad Aaboud et al. Luminosity determination in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector at the LHC. 2016.
- [31] Morad Aaboud et al. Measurement of W^+W^- production in association with one jet in proton–proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Phys. Lett.*, B763:114–133, 2016.
- [32] Morad Aaboud et al. Search for dark matter produced in association with a hadronically decaying vector boson in pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector. *Phys. Lett.*, B763:251–268, 2016.
- [33] Morad Aaboud et al. Study of hard double-parton scattering in four-jet events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment. 2016.
- [34] Morad Aaboud et al. Search for Minimal Supersymmetric Standard Model Higgs bosons H/A and for a Z' boson in the $\tau\tau$ final state produced in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector. *Eur. Phys. J.*, C76(11):585, 2016.
- [35] Morad Aaboud et al. Dark matter interpretations of ATLAS searches for the electroweak production of supersymmetric particles in $\sqrt{s} = 8$ TeV proton-proton collisions. *JHEP*, 09:175, 2016.
- [36] Morad Aaboud et al. A measurement of the calorimeter response to single hadrons and determination of the jet energy scale uncertainty using LHC Run-1 pp -collision data with the ATLAS detector. 2016.
- [37] Morad Aaboud et al. Measurement of the $b\bar{b}$ dijet cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. 2016.
- [38] Morad Aaboud et al. Search for new phenomena in different-flavour high-mass dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Eur. Phys. J.*, C76(10):541, 2016.
- [39] Morad Aaboud et al. Measurement of top quark pair differential cross-sections in the dilepton channel in pp collisions at $\sqrt{s} = 7$ and 8 TeV with ATLAS. 2016.
- [40] Morad Aaboud et al. Measurement of the total cross section from elastic scattering in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Phys. Lett.*, B761:158–178, 2016.

- [41] Morad Aaboud et al. Search for heavy resonances decaying to a Z boson and a photon in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. 2016.
- [42] Morad Aaboud et al. Search for squarks and gluinos in events with hadronically decaying tau leptons, jets and missing transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV recorded with the ATLAS detector. 2016.
- [43] Morad Aaboud et al. Search for new resonances decaying to a W or Z boson and a Higgs boson in the $\ell^+\ell^-b\bar{b}$, $\ell\nu b\bar{b}$, and $\nu\bar{\nu}b\bar{b}$ channels with pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. 2016.
- [44] Morad Aaboud et al. Measurement of exclusive $\gamma\gamma \rightarrow W^+W^-$ production and search for exclusive Higgs boson production in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector. *Phys. Rev.*, D94(3):032011, 2016.
- [45] Morad Aaboud et al. Search for high-mass new phenomena in the dilepton final state using proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Lett.*, B761:372–392, 2016.
- [46] Morad Aaboud et al. Search for Higgs and Z Boson Decays to $\phi\gamma$ with the ATLAS Detector. *Phys. Rev. Lett.*, 117(11):111802, 2016.
- [47] Morad Aaboud et al. Measurement of jet activity in top quark events using the $e\mu$ final state with two b -tagged jets in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *JHEP*, 09:074, 2016.
- [48] Morad Aaboud et al. Search for supersymmetry in a final state containing two photons and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions at the LHC using the ATLAS detector. *Eur. Phys. J.*, C76(9):517, 2016.
- [49] Morad Aaboud et al. Search for bottom squark pair production in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Eur. Phys. J.*, C76(10):547, 2016.
- [50] Morad Aaboud et al. Search for the Higgs boson produced in association with a W boson and decaying to four b -quarks via two spin-zero particles in pp collisions at 13 TeV with the ATLAS detector. *Eur. Phys. J.*, C76(11):605, 2016.
- [51] Morad Aaboud et al. Measurement of forward-backward multiplicity correlations in lead-lead, proton-lead and proton-proton collisions with the ATLAS detector. 2016.
- [52] Georges Aad et al. The performance of the jet trigger for the ATLAS detector during 2011 data taking. *Eur. Phys. J.*, C76(10):526, 2016.

- [53] Morad Aaboud et al. Search for heavy long-lived charged R -hadrons with the ATLAS detector in 3.2 fb^{-1} of proton–proton collision data at $\sqrt{s} = 13$ TeV. *Phys. Lett.*, B760:647–665, 2016.
- [54] Morad Aaboud et al. Searches for heavy diboson resonances in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *JHEP*, 09:173, 2016.
- [55] Morad Aaboud et al. Search for pair production of Higgs bosons in the $b\bar{b}b\bar{b}$ final state using proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Rev.*, D94(5):052002, 2016.
- [56] Morad Aaboud et al. Measurement of the $W^{\pm}Z$ boson pair-production cross section in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector. *Phys. Lett.*, B762:1–22, 2016.
- [57] Morad Aaboud et al. Search for new resonances in events with one lepton and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Lett.*, B762:334–352, 2016.
- [58] Morad Aaboud et al. Search for top squarks in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector. *Phys. Rev.*, D94(5):052009, 2016.
- [59] Morad Aaboud et al. Search for resonances in diphoton events at $\sqrt{s}=13$ TeV with the ATLAS detector. *JHEP*, 09:001, 2016.
- [60] Morad Aaboud et al. Measurement of the $t\bar{t}$ production cross-section using $e\mu$ events with b-tagged jets in pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector. *Phys. Lett.*, B761:136–157, 2016.
- [61] Morad Aaboud et al. Measurement of the Inelastic Proton-Proton Cross Section at $\sqrt{s} = 13$ TeV with the ATLAS Detector at the LHC. *Phys. Rev. Lett.*, 117(18):182002, 2016.
- [62] Georges Aad et al. Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV. *JHEP*, 08:045, 2016.
- [63] Morad Aaboud et al. Search for TeV-scale gravity signatures in high-mass final states with leptons and jets with the ATLAS detector at $\sqrt{s} = 13$ TeV. *Phys. Lett.*, B760:520–537, 2016.
- [64] Morad Aaboud et al. Search for the Standard Model Higgs boson produced by vector-boson fusion in 8 TeV pp collisions and decaying to bottom quarks with the ATLAS detector. 2016.
- [65] Morad Aaboud et al. Measurement of the top quark mass in the $t\bar{t} \rightarrow$ dilepton channel from $\sqrt{s} = 8$ TeV ATLAS data. *Phys. Lett.*, B761:350–371, 2016.

- [66] Morad Aaboud et al. Measurement of the photon identification efficiencies with the ATLAS detector using LHC Run-1 data. 2016.
- [67] Georges Aad et al. Measurement of the double-differential high-mass Drell-Yan cross section in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *JHEP*, 08:009, 2016.
- [68] Morad Aaboud et al. Charged-particle distributions at low transverse momentum in $\sqrt{s} = 13$ TeV pp interactions measured with the ATLAS detector at the LHC. *Eur. Phys. J.*, C76(9):502, 2016.
- [69] Georges Aad et al. Measurement of the angular coefficients in Z -boson events using electron and muon pairs from data taken at $\sqrt{s} = 8$ TeV with the ATLAS detector. *JHEP*, 08:159, 2016.
- [70] Georges Aad et al. Search for pair production of gluinos decaying via stop and sbottom in events with b -jets and large missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Rev.*, D94(3):032003, 2016.
- [71] Morad Aaboud et al. Measurement of the relative width difference of the B^0 - \bar{B}^0 system with the ATLAS detector. *JHEP*, 06:081, 2016.
- [72] Georges Aad et al. Transverse momentum, rapidity, and centrality dependence of inclusive charged-particle production in $\sqrt{s_{NN}} = 5.02$ TeV p+Pb collisions measured by the ATLAS experiment. *Phys. Lett.*, B763:313, 2016.
- [73] Morad Aaboud et al. Search for scalar leptoquarks in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS experiment. *New J. Phys.*, 18(9):093016, 2016.
- [74] Georges Aad et al. Search for gluinos in events with an isolated lepton, jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Eur. Phys. J.*, C76(10):565, 2016.
- [75] Morad Aaboud et al. Search for squarks and gluinos in final states with jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Eur. Phys. J.*, C76(7):392, 2016.
- [76] Georges Aad et al. Measurement of the inclusive isolated prompt photon cross section in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *JHEP*, 08:005, 2016.
- [77] Morad Aaboud et al. Search for new phenomena in final states with an energetic jet and large missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector. *Phys. Rev.*, D94(3):032005, 2016.
- [78] Georges Aad et al. Search for lepton-flavour-violating decays of the Higgs and Z bosons with the ATLAS detector. 2016.

- [79] Georges Aad et al. Measurements of the charge asymmetry in top-quark pair production in the dilepton final state at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Phys. Rev.*, D94(3):032006, 2016.
- [80] Georges Aad et al. Measurements of $Z\gamma$ and $Z\gamma\gamma$ production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Phys. Rev.*, D93(11):112002, 2016.
- [81] Morad Aaboud et al. Search for metastable heavy charged particles with large ionization energy loss in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS experiment. *Phys. Rev.*, D93(11):112015, 2016.
- [82] Morad Aaboud et al. Study of the rare decays of B_s^0 and B^0 into muon pairs from data collected during the LHC Run 1 with the ATLAS detector. *Eur. Phys. J.*, C76(9):513, 2016.
- [83] Georges Aad et al. Search for the Standard Model Higgs boson decaying into $b\bar{b}$ produced in association with top quarks decaying hadronically in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *JHEP*, 05:160, 2016.
- [84] Georges Aad et al. Measurement of fiducial differential cross sections of gluon-fusion production of Higgs bosons decaying to $W\bar{W}$ with the ATLAS detector at $\sqrt{s} = 8$ TeV. *JHEP*, 08:104, 2016.
- [85] Morad Aaboud et al. Search for new phenomena in events with a photon and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *JHEP*, 06:059, 2016.
- [86] Georges Aad et al. Measurement of W^\pm and Z -boson production cross sections in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Lett.*, B759:601–621, 2016.
- [87] Morad Aaboud et al. Search for charged Higgs bosons produced in association with a top quark and decaying via $H^\pm \rightarrow \tau\nu$ using pp collision data recorded at $\sqrt{s} = 13$ TeV by the ATLAS detector. *Phys. Lett.*, B759:555–574, 2016.
- [88] Georges Aad et al. Beam-induced and cosmic-ray backgrounds observed in the ATLAS detector during the LHC 2012 proton-proton running period. *JINST*, 11(05):P05013, 2016.
- [89] Morad Aaboud et al. Search for resonances in the mass distribution of jet pairs with one or two jets identified as b -jets in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Lett.*, B759:229–246, 2016.
- [90] Georges Aad et al. Muon reconstruction performance of the ATLAS detector in protonproton collision data at $\sqrt{s} = 13$ TeV. *Eur. Phys. J.*, C76(5):292, 2016.

- [91] Georges Aad et al. Identification of high transverse momentum top quarks in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *JHEP*, 06:093, 2016.
- [92] Georges Aad et al. Topological cell clustering in the ATLAS calorimeters and its performance in LHC Run 1. 2016.
- [93] Georges Aad et al. Charged-particle distributions in pp interactions at $\sqrt{s} = 8$ TeV measured with the ATLAS detector. *Eur. Phys. J.*, C76(7):403, 2016.
- [94] Georges Aad et al. Measurements of $W^\pm Z$ production cross sections in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector and limits on anomalous gauge boson self-couplings. *Phys. Rev.*, D93(9):092004, 2016.
- [95] Georges Aad et al. Measurement of total and differential W^+W^- production cross sections in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector and limits on anomalous triple-gauge-boson couplings. *JHEP*, 09:029, 2016.
- [96] Georges Aad et al. Search for supersymmetry at $\sqrt{s} = 13$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector. *Eur. Phys. J.*, C76(5):259, 2016.
- [97] Georges Aad et al. Measurement of event-shape observables in $Z \rightarrow \ell^+\ell^-$ events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector at the LHC. *Eur. Phys. J.*, C76(7):375, 2016.
- [98] Georges Aad et al. Search for new phenomena in final states with large jet multiplicities and missing transverse momentum with ATLAS using $\sqrt{s} = 13$ TeV proton-proton collisions. *Phys. Lett.*, B757:334–355, 2016.
- [99] Georges Aad et al. Search for single production of a vector-like quark via a heavy gluon in the $4b$ final state with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV. *Phys. Lett.*, B758:249–268, 2016.
- [100] Georges Aad et al. Search for single production of vector-like quarks decaying into Wb in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Eur. Phys. J.*, C76(8):442, 2016.
- [101] Georges Aad et al. Test of CP Invariance in vector-boson fusion production of the Higgs boson using the Optimal Observable method in the ditau decay channel with the ATLAS detector. 2016.
- [102] Georges Aad et al. Charged-particle distributions in $\sqrt{s}=13$ TeV pp interactions measured with the ATLAS detector at the LHC. *Phys. Lett.*, B758:67–88, 2016.
- [103] Georges Aad et al. Measurement of the charged-particle multiplicity inside jets from $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector. *Eur. Phys. J.*, C76(6):322, 2016.

- [104] Georges Aad et al. A search for top squarks with R-parity-violating decays to all-hadronic final states with the ATLAS detector in $\sqrt{s} = 8$ TeV proton-proton collisions. *JHEP*, 06:067, 2016.
- [105] Georges Aad et al. A search for an excited muon decaying to a muon and two jets in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *New J. Phys.*, 18(7):073021, 2016.
- [106] Georges Aad et al. Probing lepton flavour violation via neutrinoless $\tau \rightarrow 3\mu$ decays with the ATLAS detector. *Eur. Phys. J.*, C76(5):232, 2016.
- [107] Georges Aad et al. Measurement of the charge asymmetry in highly boosted top-quark pair production in $\sqrt{s} = 8$ TeV pp collision data collected by the ATLAS experiment. *Phys. Lett.*, B756:52–71, 2016.
- [108] Georges Aad et al. Reconstruction of hadronic decay products of tau leptons with the ATLAS experiment. *Eur. Phys. J.*, C76(5):295, 2016.
- [109] Georges Aad et al. Search for new phenomena with photon+jet events in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *JHEP*, 03:041, 2016.
- [110] Georges Aad et al. Measurement of the ZZ Production Cross Section in pp Collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector. *Phys. Rev. Lett.*, 116(10):101801, 2016.
- [111] Georges Aad et al. Combination of searches for WW , WZ , and ZZ resonances in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Phys. Lett.*, B755:285–305, 2016.
- [112] Georges Aad et al. Search for charged Higgs bosons in the $H^\pm \rightarrow tb$ decay channel in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector. *JHEP*, 03:127, 2016.
- [113] Georges Aad et al. Search for strong gravity in multijet final states produced in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC. *JHEP*, 03:026, 2016.
- [114] Georges Aad et al. Measurement of the transverse momentum and ϕ_η^* distributions of DrellYan lepton pairs in protonproton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Eur. Phys. J.*, C76(5):291, 2016.
- [115] Georges Aad et al. Search for new phenomena in dijet mass and angular distributions from pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Lett.*, B754:302–322, 2016.
- [116] Georges Aad et al. Search for the Standard Model Higgs boson produced in association with a vector boson and decaying into a tau pair in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Phys. Rev.*, D93(9):092005, 2016.

- [117] Georges Aad et al. Evidence for single top-quark production in the s -channel in proton-proton collisions at $\sqrt{s}=8$ TeV with the ATLAS detector using the Matrix Element Method. *Phys. Lett.*, B756:228–246, 2016.
- [118] Georges Aad et al. A search for prompt lepton-jets in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector. *JHEP*, 02:062, 2016.
- [119] Georges Aad et al. Measurements of top-quark pair differential cross-sections in the lepton+jets channel in pp collisions at $\sqrt{s}=8$ TeV using the ATLAS detector. *Eur. Phys. J.*, C76(10):538, 2016.
- [120] Georges Aad et al. Dijet production in $\sqrt{s}=7$ TeV pp collisions with large rapidity gaps at the ATLAS experiment. *Phys. Lett.*, B754:214–234, 2016.
- [121] Georges Aad et al. Measurement of the correlations between the polar angles of leptons from top quark decays in the helicity basis at $\sqrt{s}=7$ TeV using the ATLAS detector. *Phys. Rev.*, D93(1):012002, 2016.
- [122] Georges Aad et al. Search for dark matter produced in association with a Higgs boson decaying to two bottom quarks in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector. *Phys. Rev.*, D93(7):072007, 2016.
- [123] Georges Aad et al. Identification of boosted, hadronically decaying W bosons and comparisons with ATLAS data taken at $\sqrt{s}=8$ TeV. *Eur. Phys. J.*, C76(3):154, 2016.
- [124] Georges Aad et al. Measurement of the differential cross-section of highly boosted top quarks as a function of their transverse momentum in $\sqrt{s}=8$ TeV proton-proton collisions using the ATLAS detector. *Phys. Rev.*, D93(3):032009, 2016.
- [125] Georges Aad et al. Search for anomalous couplings in the Wtb vertex from the measurement of double differential angular decay rates of single top quarks produced in the t -channel with the ATLAS detector. *JHEP*, 04:023, 2016.
- [126] Georges Aad et al. Measurement of the production cross-section of a single top quark in association with a W boson at 8 TeV with the ATLAS experiment. *JHEP*, 01:064, 2016.
- [127] Georges Aad et al. Search for the production of single vector-like and excited quarks in the Wt final state in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector. *JHEP*, 02:110, 2016.
- [128] Georges Aad et al. Search for magnetic monopoles and stable particles with high electric charges in 8 TeV pp collisions with the ATLAS detector. *Phys. Rev.*, D93(5):052009, 2016.

- [129] Georges Aad et al. Measurements of four-lepton production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Phys. Lett.*, B753:552–572, 2016.
- [130] Georges Aad et al. Search for the electroweak production of supersymmetric particles in $\sqrt{s}=8$ TeV pp collisions with the ATLAS detector. *Phys. Rev.*, D93(5):052002, 2016.
- [131] Georges Aad et al. Search for flavour-changing neutral current top quark decays $t \rightarrow Hq$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *JHEP*, 12:061, 2015.
- [132] Georges Aad et al. Measurement of the $t\bar{t}W$ and $t\bar{t}Z$ production cross sections in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *JHEP*, 11:172, 2015.
- [133] Georges Aad et al. Measurement of jet charge in dijet events from $\sqrt{s}=8$ TeV pp collisions with the ATLAS detector. *Phys. Rev.*, D93(5):052003, 2016.
- [134] Georges Aad et al. Search for new phenomena in events with at least three photons collected in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Eur. Phys. J.*, C76(4):210, 2016.
- [135] Georges Aad et al. Observation of Long-Range Elliptic Azimuthal Anisotropies in $\sqrt{s} = 13$ and 2.76 TeV pp Collisions with the ATLAS Detector. *Phys. Rev. Lett.*, 116(17):172301, 2016.
- [136] Georges Aad et al. Searches for Higgs boson pair production in the $hh \rightarrow b\bar{b}\tau\tau, \gamma\gamma WW^*, \gamma\gamma b\bar{b}, b\bar{b}b\bar{b}$ channels with the ATLAS detector. *Phys. Rev.*, D92:092004, 2015.
- [137] Georges Aad et al. Search for pair production of a new heavy quark that decays into a W boson and a light quark in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Phys. Rev.*, D92(11):112007, 2015.
- [138] Georges Aad et al. Measurement of the charge asymmetry in top-quark pair production in the lepton-plus-jets final state in pp collision data at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Eur. Phys. J.*, C76(2):87, 2016.
- [139] Georges Aad et al. Constraints on new phenomena via Higgs boson couplings and invisible decays with the ATLAS detector. *JHEP*, 11:206, 2015.
- [140] Georges Aad et al. Search for a high-mass Higgs boson decaying to a W boson pair in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *JHEP*, 01:032, 2016.
- [141] Georges Aad et al. Measurements of fiducial cross-sections for $t\bar{t}$ production with one or two additional b-jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector. *Eur. Phys. J.*, C76(1):11, 2016.

- [142] Georges Aad et al. Summary of the ATLAS experiments sensitivity to supersymmetry after LHC Run 1 interpreted in the phenomenological MSSM. *JHEP*, 10:134, 2015.
- [143] Georges Aad et al. Searches for scalar leptoquarks in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Eur. Phys. J.*, C76(1):5, 2016.
- [144] Georges Aad et al. Search for lepton-flavour-violating H decays of the Higgs boson with the ATLAS detector. *JHEP*, 11:211, 2015.
- [145] Georges Aad et al. Constraints on non-Standard Model Higgs boson interactions in an effective Lagrangian using differential cross sections measured in the $H \rightarrow \gamma\gamma$ decay channel at $\sqrt{s} = 8$ TeV with the ATLAS detector. *Phys. Lett.*, B753:69–85, 2016.