

Honors Mathematics III

Review — Midterm 1

CHEN Xiwen

UM-SJTU Joint Institute

June 18, 2018

Table of contents

Linear Maps

Matrices

Theorem of Systems of Linear Equations

Determinant

Overview

Final Remarks

Linear Maps

Matrices

Theorem of Systems of Linear Equations

Determinant

Overview

Final Remarks

Linear maps

To prove a map from vector space (U, \oplus, \odot) to another vector space (V, \boxplus, \boxdot) is **linear**, it needs to be

- ▶ Homogeneous: $L(\lambda \odot u) = \lambda \boxdot L(u)$, and
- ▶ Additive: $L(u \oplus u') = L(u) \boxplus L(u')$.

Examples.

1. For $I \in \mathbb{R}$, the map $D : C^1(I) \rightarrow C(I)$, $f \mapsto f'$ is linear.
2. The complex conjugation map in \mathbb{C} is linear if \mathbb{C} is regarded as a real vector space.
3. **Exercise 2.6.** \mathcal{P}_n is the vector space of real polynomials over \mathbb{R} of degree at most n . The map

$$\alpha : \mathcal{P}_n \rightarrow \mathbb{R}, \quad \alpha(p) = \int_{-1}^1 p(x) dx$$

is linear.

Range and Kernel

Definitions.

- ▶ **Dual space:** $V^* = \mathcal{L}(V, \mathbb{F})$ for a finite-dimensional vector space V with basis $\{b_1, \dots, b_n\}$.
- ▶ **Dual basis:** $\{b_1^*, \dots, b_n^*\}$ with

$$b_k^*(b_j) = \begin{cases} 1 & k = j \\ 0 & k \neq j \end{cases}$$

Exercise 2.5

1. $L = \sum_{i=1}^n L(b_i)b_i^*$ for a linear map $L \in \mathcal{L}(V, \mathbb{F})$.
2. $V \cong V^*$.

Range and Kernel

Definitions.

$$\operatorname{ran} L := \left\{ v \in V : \exists_{u \in U} v = Lu \right\}, \quad \ker L := \{ u \in U : Lu = 0 \}$$

Results.

1. A linear map $L \in \mathcal{L}(U, V)$ is injective iff $\ker L = \{0\}$.
2. **Dimension formula.** $\dim \operatorname{ran} L + \dim \ker L = \dim U$ (for finite-dimensional vector spaces).

Isomorphisms

- ▶ For n -dimensional vector spaces U, V , an isomorphism maps from basis (b_1, \dots, b_n) to basis (Lb_1, \dots, Lb_n) .
- ▶ For finite-dimensional vector spaces U, V , $U \cong V \Leftrightarrow \dim U = \dim V$.
- ▶ If $\dim U = \dim V$, then for a linear map $L \in \mathcal{L}(U, V)$, injective \Leftrightarrow surjective \Leftrightarrow bijective.

Operator Norm

- ▶ Equivalent definitions:

$$\|L\| := \sup_{\substack{u \in U \\ u \neq 0}} \frac{\|Lu\|_V}{\|u\|_U} = \sup_{\substack{u \in U \\ \|u\|_U = 1}} \|Lu\|_V$$

- ▶ Additional property:

$$\|L_2 L_1\| \leq \|L_2\| \cdot \|L_1\|, \quad L_1 \in \mathcal{L}(U, V), \quad L_2 \in \mathcal{L}(V, W)$$

Linear Maps

Matrices

Theorem of Systems of Linear Equations

Determinant

Overview

Final Remarks

Linear Maps as Matrices

Every linear map between finite-dimensional vector spaces can be expressed as a matrix.

$$\begin{array}{ccc} U & \xrightarrow{L} & V \\ \varphi_{\mathcal{A}} \downarrow & & \downarrow \varphi_{\mathcal{B}} \\ \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \end{array} \quad \Phi_{\mathcal{A}}^{\mathcal{B}}(L) = A = \varphi_{\mathcal{B}} \circ L \circ \varphi_{\mathcal{A}}^{-1}$$

- ▶ Matrix multiplication.
- ▶ Transpose A^T and adjoint A^* , $\langle x, Ay \rangle = \langle A^*x, y \rangle$.
- ▶ Inverse A^{-1} of $n \times n$ matrices. (Find inverse using Gauss-Jordan algorithm.)
- ▶ *Change basis.*

Change Basis — Passive Point of View

1. Find basis change matrix T such that $e'_i = Te_i$.
2. Find inverse of T .
3. Find matrix A representing the operation with respect to the new basis.
4. Calculate TAT^{-1} .

Linear Maps

Matrices

Theorem of Systems of Linear Equations

Determinant

Overview

Final Remarks

Solution Set

Structure of solution set.

$$\text{Sol}(A, b) = \{x_0\} + \ker A$$

Fredholm alternatives.

1. $Ax = b$ has a unique solution for any $b \in \mathbb{R}^n$.
 - ▶ A is invertible.
 - ▶ $\det A \neq 0$.
 - ▶ $\ker A = \{0\}$.
2. $Ax = 0$ has a non-trivial solution. ($Ax = b$ either has no solution or infinitely many solutions.)
 - ▶ A is not invertible.
 - ▶ $\det A = 0$.
 - ▶ $\ker A \neq \{0\}$.

In general, a system $Ax = b$ is solvable if b is in the range of A , i.e., in the span of column vectors of matrix A .

Linear Maps

Matrices

Theorem of Systems of Linear Equations

Determinant

Overview

Final Remarks

Determinant

- ▶ Determinants in \mathbb{R}^2 and \mathbb{R}^3
- ▶ Permutations.
- ▶ Group and group actions.
- ▶ Determinant in \mathbb{R}^n .

Properties of Determinant

- ▶ Normed. $\det \text{id} = 1$.
- ▶ Multilinear.
- ▶ Alternating:
 - ▶ \det is alternating.
 - ▶

$$\begin{aligned} & \det(a_1, \dots, a_{(j-1)}, a_j, a_{(j+1)}, \dots, a_{(k-1)}, a_k, a_{(k+1)}, \dots, a_p) \\ &= - \det(a_1, \dots, a_{(j-1)}, a_k, a_{(j+1)}, \dots, a_{(k-1)}, a_j, a_{(k+1)}, \dots, a_p) \end{aligned}$$

- ▶ $\det(a_1, \dots, a_p) = 0$ if a_1, \dots, a_p are linearly dependent.

Properties of Determinant

1. Elementary column operations.

- ▶ $\det(a_2, a_1, \dots, a_n) = -\det(a_1, a_2, \dots, a_n)$.
- ▶ $\det(a_1, \dots, \lambda a_j, \dots, a_n) = \lambda \det(a_1, \dots, a_j, \dots, a_n)$.
- ▶ $\det(a_1, \dots, a_j, \dots, a_k + \lambda a_j, \dots, a_n) = \det(a_1, \dots, a_j, \dots, a_k, \dots, a_n)$.

2. $\det A = \det A^T$.

3. $\det(AB) = (\det A)(\det B)$.

4. $\det A = \sum_{\pi \in S_n} \operatorname{sgn} \pi \, a_{1\pi(1)} \cdots a_{n\pi(n)}$.

5. $\det A = \sum_{i=1}^n (-1)^{i+j} a_{ij} \det A_{ij}$.

Results from Determinant

- Solve linear system of equations.

$$x_i = \frac{1}{\det A} \det(a_1, \dots, a_{i-1}, b, a_{i+1}, \dots, a_n), \quad i = 1, \dots, n$$

- Find inverse.

$$A^{-1} = \frac{1}{\det A} A^\#$$

Calculating Determinants

Few methods to calculate determinant:

1. In \mathbb{R}^3 : expand to the case of \mathbb{R}^2 and calculate directly.
2. In higher dimensions: use properties of determinant. Apply *elementary row or column operations* combined with Laplace expansion.
3. Leibnitz Formula.
4. Induction.

Linear Maps

Matrices

Theorem of Systems of Linear Equations

Determinant

Overview

Final Remarks

Systems of Linear Equations

Gauss-Jordan algorithm.

Finite-Dimensional Vector Spaces

Definition of Linear Independence.

$$v = \sum_{i=1}^n \lambda_i v_i = 0 \quad \Leftrightarrow \quad \lambda_i = 0$$

- ▶ v is a n -dimensional vector.
- ▶ v is a linear map: v acting on any vector (basis) gives 0.

Exercise 2.5. The maps b_1^*, \dots, b_n^* form a basis of V^* .

1. $L = \sum_{i=1}^n \lambda_i b_i^* = 0 \Leftrightarrow Lb_i = 0 \Leftrightarrow \lambda_i = 0$.
2. $\forall L \in \mathcal{L}(V, \mathbb{F}), L = \sum_{i=1}^n L(b_i) b_i^*$.

Finite-Dimensional Vector Spaces

Basis (Any two combined.)

- ▶ $\text{span}\{b_1, \dots, b_n\} = V$.
- ▶ $\{b_1, \dots, b_n\}$ is linearly independent.
- ▶ The length of the set $\{b_1, \dots, b_n\}$ is $\dim V$.

Proof using basis.

- ▶ Definition: unique representation.
- ▶ Characterization of basis:
 1. $\{b_1, \dots, b_n\}$ is independent.
 2. $V = \text{span}\{b_1, \dots, b_n\}$.
- ▶ Finite-dimensional vector spaces (subspaces) — basis extension theorem.

e.g. $\dim(U + W) + \dim(U \cap W) = \dim U + \dim W$.

Inner Product Spaces

Inner Product.

► Definitions of inner product. — proof of inner products.

1. $\langle v, v \rangle \geq 0$ and $\langle v, v \rangle = 0$ iff $v = 0$.
2. $\langle u, v + w \rangle = \langle u, v \rangle + \langle u, w \rangle$.
3. $\langle u, \lambda v \rangle = \lambda \langle u, v \rangle$.
4. $\langle u, v \rangle = \overline{\langle v, u \rangle}$.

► Induced norm. $\|v\|^2 = \langle v, v \rangle$.

e.g. $\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2)$.

► Normalization and projection.

e.g. (2.4) The vector space $C([-1, 1])$ of real continuous functions on $[-1, 1]$ endowed with the scalar product $\langle f, g \rangle := \int_{-1}^1 fg$ and the induced norm $\|f\| = \sqrt{\langle f, f \rangle}$.

1. $m_k(x) = x^k, k = 0, 1, \dots$
2. $\{1, \sin(n\pi x), \cos(n\pi x)\}_{n=1}^{\infty}$.

► **Note:** $V = A \oplus B, V = A \oplus C \not\Rightarrow B = C$.

Linear Maps

- ▶ Vector spaces:
 1. Dimension.
 2. Basis.
- ▶ Linear map:
 1. Range.
 2. Kernel.

Matrices

Matrices as Linear Maps

$$\begin{array}{ccc} U & \xrightarrow{L} & V \\ \varphi_{\mathcal{A}} \downarrow & & \downarrow \varphi_{\mathcal{B}} \\ \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^m \end{array} \quad \Phi_{\mathcal{A}}^{\mathcal{B}}(L) = A = \varphi_{\mathcal{B}} \circ L \circ \varphi_{\mathcal{A}}^{-1}$$

- ▶ $\text{ran } L = \text{span}\{a_{.1}, \dots, a_{.n}\}$.
- ▶ Matrix acting on x .
- ▶ Matrix elements. $\langle e_i, Ae_j \rangle = a_{ij}$.

Basis change. Use a basis that is convenient for the operation.

- ▶ Projection.
- ▶ Rotation.

e.g.(3.5.) Find the matrix describing rotation about the axis through the origin and the point $(1, 2, -1)^T$.

Theory of Systems of Linear Equations

- ▶ Fredholm alternatives.
- ▶ Matrix rank.
- ▶ Presenting solution sets.

Determinants

- ▶ Properties. (Row and column operations.)
- ▶ Matrix implications.
- ▶ Relations to system of linear equations.

Linear Maps

Matrices

Theorem of Systems of Linear Equations

Determinant

Overview

Final Remarks

Final Remarks

1. Vectors and vector spaces. ($\text{ran } V$, $\text{ker } V$, $\text{span } \mathcal{B}$ are all vector spaces!)
2. If the notations do not occur in the problem statement, define them clearly.
3. Do not forget to normalize the vectors using the *specified* inner product in Gram-Schmidt orthonormalization.
4. Consider using properties to reduce work.
5. Be careful with calculations.

Thanks for your attention!
Good Luck!