

Honors Mathematics III

Review — Final

CHEN Xiwen

UM-SJTU Joint Institute

August 7, 2018

Table of contents

Integration in Practice

- Integration Over Ordinate Regions
- Theorems for Integration

Surfaces and Surface Integrals

- Tangent Spaces of Surfaces
- The Normal Vector to Hypersurfaces
- Orientation of Hypersurfaces
- Scalar Surface Integrals
- Flux Through Hypersurfaces

The Theorems of Gauss and Stokes

Summary

- Elements
- Practical Integration

Integration in Practice

Integration Over Ordinate Regions

Theorems for Integration

Surfaces and Surface Integrals

Tangent Spaces of Surfaces

The Normal Vector to Hypersurfaces

Orientation of Hypersurfaces

Scalar Surface Integrals

Flux Through Hypersurfaces

The Theorems of Gauss and Stokes

Summary

Elements

Practical Integration

Integration Over Cuboids

By Fubini's Theorem, we have

$$\int_Q f = \int_{a_n}^{b_n} \cdots \int_{a_2}^{b_2} \int_{a_1}^{b_1} f(x_1, x_2, \dots, x_n) dx_1 dx_2 \dots dx_n.$$

or equivalently,

$$\int_Q f = \int_Q f(x) dx$$

Integration Over Ordinate Regions

Definitions.

- ▶ **Ordinate region (with respect to x_k):** there exists a measurable set $\Omega \subset \mathbb{R}^{n-1}$ and continuous, almost everywhere differentiable functions $\varphi_1, \varphi_2 : \Omega \rightarrow \mathbb{R}$ such that

$$U = \{x \in \mathbb{R}^n : x \in \Omega, \varphi_1(\hat{x}^{(k)}) \leq x_k \leq \varphi_2(\hat{x}^{(k)})\}.$$

- ▶ **Simple region:** U is an ordinate region with respect to each $x_k, k = 1, \dots, n$.

Integration Over Ordinate Regions

For an ordinate region $U \subset \mathbb{R}^n$ with respect to x_k over a measurable set Ω , the indicator function $\mathbb{1}_U$ takes the form

$$\mathbb{1}_U(x) = \mathbb{1}_\Omega \cdot \mathbb{1}_{[\varphi_1(\hat{x}^{(k)}, \varphi_2(\hat{x}^{(k)}))]}(x_k).$$

It then follows that

$$\int_U f(x) dx_1 \dots dx_n = \int_\Omega \left(\int_{\varphi_1(\hat{x}^{(k)})}^{\varphi_2(\hat{x}^{(k)})} f(x) dx_k \right) d\hat{x}^{(k)}$$

if

$$\int_{\varphi_1(\hat{x}^{(k)})}^{\varphi_2(\hat{x}^{(k)})} f(x) dx_k$$

exists for every $\hat{x}^{(k)} \in \Omega$.

Integration in Practice

Integration Over Ordinate Regions

Theorems for Integration

Surfaces and Surface Integrals

Tangent Spaces of Surfaces

The Normal Vector to Hypersurfaces

Orientation of Hypersurfaces

Scalar Surface Integrals

Flux Through Hypersurfaces

The Theorems of Gauss and Stokes

Summary

Elements

Practical Integration

Substitution Rule

3.4.12. Substitution Rule. Let $\Omega \subset \mathbb{R}^n$ be open and $g : \Omega \rightarrow \mathbb{R}^n$ injective and continuously differentiable. Suppose that $\det J_g(y) \neq 0$ for all $y \in \Omega$. Let K be a compact measurable subset of Ω . Then $g(K)$ is compact and measurable and if $f : g(K) \rightarrow \mathbb{R}$ is integrable, then

$$\int_{g(K)} f(x) dx = \int_K f(g(y)) \cdot |\det J_g(y)| dy.$$

Coordinate Systems

- Polar coordinates:

$$x = r \cos \phi, \quad y = r \sin \phi, \quad |\det J(r, \phi)| = r$$

- Cylindrical coordinates:

$$x = r \cos \phi, \quad y = r \sin \phi, \quad z = \zeta, \quad |\det J(r, \phi, \zeta)| = r$$

- Spherical coordinates:

$$x = r \cos \phi \sin \theta, \quad y = r \sin \phi \sin \theta, \quad z = r \cos \theta$$

$$|\det J(r, \phi, \theta)| = r^2 \sin \theta.$$

Coordinate Systems

- Spherical coordinates in \mathbb{R}^n :

$$x_1 = r \cos \theta_1$$

$$x_2 = r \sin \theta_1 \cos \theta_2$$

$$x_3 = r \sin \theta_1 \sin \theta_2 \cos \theta_3$$

$$\vdots$$

$$x_{n-1} = r \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{n-2} \cos \theta_{n-1}$$

$$x_n = r \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{n-2} \sin \theta_{n-1}$$

$$|\det J(r, \theta_1, \dots, \theta_{n-1})| = r^{n-1} \sin^{n-2} \theta_1 \sin^{n-3} \theta_2 \cdots \sin \theta_{n-2}$$

Note. $r > 0, 0 < \theta_k < \pi, k = 1, \dots, n-2, 0 < \theta_{n-1} < 2\pi$.

The Gauss Integral

The Gauss Integral.

$$\lim_{a \rightarrow \infty} I(a) := \int_{-\infty}^{\infty} e^{-x^2/2} dx = \sqrt{2\pi}.$$

Variants. For $k > 0$,

$$\int_{-\infty}^{\infty} e^{-kx^2} dx = \sqrt{\frac{\pi}{k}}.$$

Green's Theorem

Green's Theorem. Let $R \subset \mathbb{R}^2$ be bounded, simple region and $\Omega \supset R$ an open set containing R . Let $F : \Omega \rightarrow \mathbb{R}^2$ be continuously differentiable vector field. Then

$$\int_{\partial R^*} F d\vec{s} = \int_R \left(\frac{\partial F_2}{\partial x_1} - \frac{\partial F_1}{\partial x_2} \right) dx$$

Physical Interpretation of Green's Theorem

► For $F = \begin{pmatrix} F_1(x) \\ F_2(x) \end{pmatrix}$,

$$\begin{aligned} \text{circulation along } \partial R &= \int_{\partial R^*} F d\vec{s} \\ &= \int_R \left(\frac{\partial F_2}{\partial x_1} - \frac{\partial F_1}{\partial x_2} \right) dx \\ &= \int_R \text{rot} F dx \\ &= \text{integral of circulation density over } R. \end{aligned}$$

Physical Interpretation of Green's Theorem

► For $\tilde{F} = \begin{pmatrix} -F_2(x) \\ F_1(x) \end{pmatrix}$,

$$\begin{aligned} \text{flux through } \partial R &= \int_{\partial R^*} \langle F, N \rangle ds = \int_{\partial R^*} \tilde{F} d\vec{s} \\ &= \int_R \left(\frac{\partial \tilde{F}_2}{\partial x_1} - \frac{\partial \tilde{F}_1}{\partial x_2} \right) dx \\ &= \int_R \operatorname{div} F dx \\ &= \text{integral of flux density over } R. \end{aligned}$$

Integration in Practice

Integration Over Ordinate Regions

Theorems for Integration

Surfaces and Surface Integrals

Tangent Spaces of Surfaces

The Normal Vector to Hypersurfaces

Orientation of Hypersurfaces

Scalar Surface Integrals

Flux Through Hypersurfaces

The Theorems of Gauss and Stokes

Summary

Elements

Practical Integration

Tangent Spaces of Surfaces

Definition. Let $\mathcal{S} \subset \mathbb{R}^n$ be a parametrized m -surface with parametrization $\varphi : \Omega \rightarrow \mathcal{S}$. Then

$$t_k(p) = \frac{\partial}{\partial x_k} \begin{pmatrix} \varphi_1(x) \\ \vdots \\ \varphi_2(x) \end{pmatrix} \bigg|_{x=\varphi^{-1}(p)}, \quad k = 1, \dots, m$$

is called the ***k -th tangent vector of \mathcal{S} at $p \in \mathcal{S}$*** and

$$T_p\mathcal{S} := \text{ran} D\varphi|_x = \text{span}\{t_1(p), \dots, t_m(p)\}$$

is called the ***tangent space*** to \mathcal{S} at p . The vector field

$$t_k : \mathcal{S} \rightarrow \mathbb{R}^n, \quad p \mapsto t_k(p)$$

is called the ***k -th tangent vector field*** on \mathcal{S} .

Integration in Practice

Integration Over Ordinate Regions

Theorems for Integration

Surfaces and Surface Integrals

Tangent Spaces of Surfaces

The Normal Vector to Hypersurfaces

Orientation of Hypersurfaces

Scalar Surface Integrals

Flux Through Hypersurfaces

The Theorems of Gauss and Stokes

Summary

Elements

Practical Integration

The Normal Vector to Hypersurfaces

Definition. Let $\mathcal{S} \subset \mathbb{R}^n$ be a hypersurface. Then a unit vector that is orthogonal to all tangent vectors to \mathcal{S} at p is called a **unit normal vector to \mathcal{S} at p** and denoted by $N(p)$. The vector field

$$N : \mathcal{S} \rightarrow \mathbb{R}^n, \quad p \mapsto N(p)$$

is called the **normal vector field** on \mathcal{S} .

Integration in Practice

Integration Over Ordinate Regions

Theorems for Integration

Surfaces and Surface Integrals

Tangent Spaces of Surfaces

The Normal Vector to Hypersurfaces

Orientation of Hypersurfaces

Scalar Surface Integrals

Flux Through Hypersurfaces

The Theorems of Gauss and Stokes

Summary

Elements

Practical Integration

Orientation of Hypersurfaces

Definitions.

- ▶ A hypersurface $\mathcal{S} \subset \mathbb{R}^n$ such that it admits a continuous normal vector field is said to be *orientable*.
- ▶ A choice of direction for the normal vector field is called an *orientation of \mathcal{S}* .
- ▶ A hypersurface that is the boundary of a measurable set $\Omega \subset \mathbb{R}^n$ with non-zero measure is said to be a *closed surface*.
- ▶ A closed hypersurface is said to have *positive orientation* if the normal vector field is chosen so that the normal vectors point *outwards* from Ω .

Integration in Practice

Integration Over Ordinate Regions

Theorems for Integration

Surfaces and Surface Integrals

Tangent Spaces of Surfaces

The Normal Vector to Hypersurfaces

Orientation of Hypersurfaces

Scalar Surface Integrals

Flux Through Hypersurfaces

The Theorems of Gauss and Stokes

Summary

Elements

Practical Integration

The Metric Tensor

Definition. Let $\mathcal{S} \subset \mathbb{R}^n$ be an m -surface with parametrization φ and tangent vector fields t_1, \dots, t_m . Then $G \in \text{Mat}(m \times m; \mathbb{R})$ given by

$$G := \begin{pmatrix} \langle t_1, t_1 \rangle & \cdots & \langle t_1, t_m \rangle \\ \vdots & \ddots & \vdots \\ \langle t_m, t_1 \rangle & \cdots & \langle t_m, t_m \rangle \end{pmatrix}$$

is said to be the **metric tensor** on \mathcal{S} with respect to φ . The coefficients

$$g_{ij} := \langle t_i, t_j \rangle, \quad i, j = 1, \dots, m,$$

are called the **metric coefficients** of G .

Scalar Surface Integrals

Definition. Let $f : \mathcal{S} \rightarrow \mathbb{R}$ be a potential function. \mathcal{S} is a parametrized m -surface with parametrization $\varphi : \Omega \rightarrow \mathcal{S}, \Omega \subset \mathbb{R}^m$. Then the **(s-calar) surface integral of f over \mathcal{S}** is defined as

$$\int_{\mathcal{S}} f dA := \int_{\Omega} f \circ \varphi \sqrt{g(x)} dx$$

Integration in Practice

Integration Over Ordinate Regions

Theorems for Integration

Surfaces and Surface Integrals

Tangent Spaces of Surfaces

The Normal Vector to Hypersurfaces

Orientation of Hypersurfaces

Scalar Surface Integrals

Flux Through Hypersurfaces

The Theorems of Gauss and Stokes

Summary

Elements

Practical Integration

Flux Through Hypersurfaces

Definition. Let $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ be a vector field defined in a neighborhood of a hypersurface \mathcal{S} with parametrization $\varphi : \Omega \rightarrow \mathbb{R}^{n+1}$, $\Omega \subset \mathbb{R}^n$. Then we define the **flux of F through \mathcal{S}** by

$$\begin{aligned}\int_{\mathcal{S}} F d\vec{A} &:= \int_{\mathcal{S}} \langle F, N \rangle dA \\ &= \int_{\Omega} \langle F \circ \varphi(x), N \circ \varphi(x) \rangle \sqrt{g(x)} dx_1 \dots dx_n\end{aligned}$$

Admissible Regions

Definitions.

- ▶ A subset $R \subset \mathbb{R}^n$ is called a **region** if it is open and (pathwise) connected.
- ▶ A region $R \subset \mathbb{R}^n$ is said to be **admissible** if it is bounded and its boundary is the union of a finite number of parametrized hypersurfaces whose normal vectors point outwards from R .
- ▶ A hypersurface $\mathcal{S} \subset \mathbb{R}^3$ with parametrization $\varphi : R \rightarrow \mathcal{S}$ is said to be **admissible** if
 1. the interior $\text{int } R$ is an admissible region in \mathbb{R}^2 with an oriented boundary curve ∂R^* and
 2. R is closed, i.e., $R = \overline{R}$.

Closed Hypersurfaces in \mathbb{R}^3

Definition. Let $\mathcal{S} \subset \mathbb{R}^3$ be an admissible hypersurface with parametrization $\varphi : R \rightarrow \mathcal{S}$. Let $\partial R^* = \mathcal{C}_1^* \cup \mathcal{C}_2^* \cup \cdots \cup \partial_k^*$, where each \mathcal{C}_i^* is an oriented smooth curve in \mathbb{R}^2 and all \mathcal{C}_i^* are pairwise disjoint.

- We say that φ **annihilates** a chain of curves $\mathcal{C}_{i_1} \cup \cdots \cup \mathcal{C}_{i_j}$ if




$$\int_{\varphi(\mathcal{C}_{i_1} \cup \cdots \cup \mathcal{C}_{i_j})} 1 ds = 0.$$

- If φ annihilates ∂R , \mathcal{S} is said to be a **closed surface**.
- Denote by $\mathcal{C}' \subset \partial R$ the largest chain of curves that is annihilated by φ . If $\mathcal{C}' \neq \partial R$ we say that \mathcal{S} is a **surface with boundary** and define

$$\partial \mathcal{C} := \varphi(\partial R \setminus \mathcal{C}').$$

Closed Hypersurfaces in \mathbb{R}^3

Examples.



Stokes's Theorem in \mathbb{R}^3

3.6.7. Stokes's Theorem. Let $\Omega \subset \mathbb{R}^3$ be an open set, $\mathcal{S} \subset \Omega$ a parametrized, admissible surface in \mathbb{R}^3 with boundary $\partial\mathcal{S}$ and let $F : \Omega \rightarrow \mathbb{R}^3$ be a continuously differentiable vector field. Then

$$\int_{\partial\mathcal{S}^*} F d\vec{s} = \int_{\mathcal{S}^*} \operatorname{rot} F d\vec{A}$$

with positive orientation and normal vectors pointing in the direction of the thumb of the right hand if the four fingers point in the direction of the tangent vector to $\partial\mathcal{S}^*$.

Gauss's Theorem

3.6.9. **Gauss's Theorem.** Let $R \subset \mathbb{R}^n$ be an admissible region and $F : \overline{R} \rightarrow \mathbb{R}^n$ a continuously differentiable vector field. Then

$$\int_R \operatorname{div} F \, dx = \int_{\partial R^*} F \, d\vec{A}$$

Green's Identities

3.6.13. Green's Identities. Let $R \subset \mathbb{R}^n$ be an admissible region and $u, v : \bar{R} \rightarrow \mathbb{R}$ be twice continuously differentiable potential functions. Then we have:

► **Green's first identity:**

$$\int_R \langle \nabla u, \nabla v \rangle dx = - \int_R u \cdot \Delta v dx + \int_{\partial R^*} u \frac{\partial v}{\partial n} dA.$$

► **Green's second identity:**

$$\int_R (u \cdot v - v \cdot u) dx = \int_{\partial R^*} \left(u \frac{\partial v}{\partial n} - v \frac{\partial u}{\partial n} \right) dA.$$

Integration in Practice

Integration Over Ordinate Regions

Theorems for Integration

Surfaces and Surface Integrals

Tangent Spaces of Surfaces

The Normal Vector to Hypersurfaces

Orientation of Hypersurfaces

Scalar Surface Integrals

Flux Through Hypersurfaces

The Theorems of Gauss and Stokes

Summary

Elements

Practical Integration

Various Kinds of Elements

- Scalar line element:

$$ds = |\gamma'(t)|dt.$$

Line integral of potential f along \mathcal{C}^* :

$$\int_{\mathcal{C}^*} f ds = \int_I (f \circ \gamma)(t) \cdot |\gamma'(t)| dt.$$

- Vectorial line element:

$$d\vec{s} = \gamma'(t)dt.$$

The line integral of vector field F along \mathcal{C}^* :

$$\int_{\mathcal{C}^*} F d\vec{s} = \int_{\mathcal{C}^*} \langle F, T \rangle ds.$$

Various Kinds of Elements

- **Volume element:** (take spherical coordinates as example.)

$$dx = |\det J_{\Phi}(r, \theta, \varphi)| dr d\theta d\phi.$$

Integration of potentials in a \mathbb{R}^3 region:

$$\int_{\Omega} f = \int_{\Phi^{-1}(\Omega)} f \circ \Phi(r, \theta, \phi) \cdot |\det J_{\Phi}(r, \theta, \varphi)| dr d\theta d\phi.$$

- **Scalar surface element of a hypersurface in \mathbb{R}^n :**

$$dA = |\det(t_1, t_2, \dots, t_{n-1}, N) \circ \varphi| dx_1 dx_2 \dots dx_{n-1}.$$

Volume or area of \mathcal{S} :

$$|\mathcal{S}| = \int_{\Omega} |\det(t_1, \dots, t_{n-1}, N) \circ \varphi(x)| dx_1 dx_2 \dots dx_{n-1}.$$

Various Kinds of Elements

- Infinitesimal surface element of arbitrary surfaces in \mathbb{R}^n :

$$dA = \sqrt{g(x)}dx,$$

where

$$G = \begin{pmatrix} \langle t_1, t_1 \rangle & \cdots & \langle t_1, t_m \rangle \\ \vdots & \ddots & \vdots \\ \langle t_m, t_1 \rangle & \cdots & \langle t_m, t_m \rangle \end{pmatrix}, \quad g(x) = \det G(\varphi(x))$$

The scalar (surface) integral of f over S :

$$\int_S f dA = \int_{\Omega} f \circ \varphi(x) \sqrt{g(x)} dx.$$

Various Kinds of Elements

- Vectorial surface element:

$$d\vec{A} = N(\varphi(x)) \cdot \sqrt{g(x)} dx.$$

The flux of F through \mathcal{S} integral:

$$\int_{\mathcal{S}} F d\vec{A} = \int_{\Omega} \langle F \circ \varphi(x), N \circ \varphi(x) \rangle \sqrt{g(x)} dx_1 \dots dx_n.$$

Integration in Practice

Integration Over Ordinate Regions

Theorems for Integration

Surfaces and Surface Integrals

Tangent Spaces of Surfaces

The Normal Vector to Hypersurfaces

Orientation of Hypersurfaces

Scalar Surface Integrals

Flux Through Hypersurfaces

The Theorems of Gauss and Stokes

Summary

Elements

Practical Integration

Integration in Practice

- Ordinate region:

$$\int_U f(x) dx_1 \dots dx_n = \int_{\Omega} \left(\int_{\varphi_1(\hat{x}^{(k)})}^{\varphi_2(\hat{x}^{(k)})} f(x) dx_k \right) d\hat{x}^{(k)}$$

- Substitution rule:

$$\int_{g(K)} f(x) dx = \int_K f(g(y)) \cdot |\det J_g(y)| dy.$$

- Green's theorem: (\mathbb{R}^2)

$$\int_{\partial R^*} F d\vec{s} = \int_R \left(\frac{\partial F_2}{\partial x_1} - \frac{\partial F_1}{\partial x_2} \right) dx.$$

Integration in Practice

- ▶ Stokes's theorem: (\mathbb{R}^3)

$$\int_{\partial S^*} F d\vec{s} = \int_{S^*} \operatorname{rot} F d\vec{A}.$$

- ▶ Gauss's theorem: (\mathbb{R}^3)

$$\int_R \operatorname{div} F dx = \int_{\partial R^*} F d\vec{A}.$$

- ▶ Green's identities:

$$\begin{aligned} \int_R \langle \nabla u, \nabla v \rangle dx &= - \int_R u \cdot \Delta v dx + \int_{\partial R^*} u \frac{\partial v}{\partial n} dA, \\ \int_R (u \cdot v - v \cdot u) dx &= \int_{\partial R^*} \left(u \frac{\partial v}{\partial n} - v \frac{\partial u}{\partial n} \right) dA. \end{aligned}$$

Thanks for your attention!
Good Luck!