



JOINT INSTITUTE
交大密西根学院

VV285 Honors Mathematics III Solution Manual for RC 4

Chen Xiwen

June 8, 2018

Exercise 1.

Expanding the determinant with respect to the 3rd column, we get

$$\det A = (-1)^{1+3} \det A_{13} = \det A_{13}$$

with

$$A' = A_{13} = \begin{pmatrix} 2 & 1 & 0 & -1 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 1 & 0 & 0 & 2 \end{pmatrix}$$

Expanding the determinant with respect to the second row,

$$\det A = (-1) \cdot (-1)^{2+2} \det A'_{22} = -\det A'_{22}$$

where

$$A'_{22} = \begin{pmatrix} 2 & 0 & -1 \\ 0 & -1 & 1 \\ 1 & 0 & 2 \end{pmatrix}, \quad \det A'_{22} = -5$$

Therefore, $\det A = 5$.

Exercise 2.

The determinant can be found by

$$\det A^{(2n)} = a \cdot (-1)^{1+1} \det A_{11}^{(2n)} + b \cdot (-1)^{2n+1} \det A_{(2n)1}^{(2n)} = a \det A_{11}^{(2n)} - b \det A_{(2n)1}^{(2n)}$$

where

$$A_{11}^{(2n)} = \begin{pmatrix} a & \cdots & b & 0 \\ \vdots & & \vdots & \vdots \\ b & \cdots & a & 0 \\ 0 & \cdots & 0 & a \end{pmatrix} \in \text{Mat}((2n-1) \times (2n-1), \mathbb{R})$$

$$A_{(2n)1}^{(2n)} = \begin{pmatrix} 0 & \cdots & 0 & b \\ a & & b & 0 \\ \vdots & & \vdots & \vdots \\ b & \cdots & a & 0 \end{pmatrix} \in \text{Mat}((2n-1) \times (2n-1), \mathbb{R})$$

Then expanding the $(2n - 1)$ th columns for the two matrices, we obtain

$$\det A = a^2 \det A^{(2n-2)} - b^2 \det A^{(2n-2)}$$

Then we attempt to prove $\det A = (a^2 - b^2)^n$ by induction on $n \geq 1$.

- When $n = 1$, it is clear that

$$\det A^{(2)} = a^2 - b^2$$

- Then based on the hypothesis on $n = k$, from the relation deduced above, we have verified that $\det A^{(2k+2)} = (a^2 - b^2) \det A^{(2k)} = (a^2 - b^2)^n$.

Therefore, the the determinant is given by

$$\det A^{(2n)} = (a^2 - b^2)^n$$

Exercise 3.

The determinant can be transferred to

$$\det D_n = \det \begin{pmatrix} a-b & 0 & 0 & \cdots & 0 \\ c & a & b & \cdots & b \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & b \\ c & \cdots & \cdots & c & a \end{pmatrix} + \det \begin{pmatrix} b & b & b & \cdots & b \\ c & a & b & \cdots & b \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & b \\ c & \cdots & \cdots & c & a \end{pmatrix}$$

$$= (a-b) \det D_{n-1} + b \det \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ c & a & b & \cdots & b \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & b \\ c & \cdots & \cdots & c & a \end{pmatrix}$$

$$= (a-b) \det D_{n-1} + b \det \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ c & a-c & b-c & \cdots & b-c \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & b-c \\ c & \cdots & \cdots & 0 & a-c \end{pmatrix}$$

$$= (a-b) \det D_{n-1} + b(a-c)^{n-1}$$

Similarly, if we decompose the column vector in the first step, then we will get the result

$$\det D_n = (a - c)\det D_{n-1} + c(a - b)^{n-1}$$

Then if $b \neq c$, eliminating D_{n-1} we then obtain

$$\det D_n = \frac{b(a - c)^n - c(a - b)^n}{b - c}$$

When $b = c$, we obtain

$$\begin{aligned}\det D_n &= (a - b)\det D_{n-1} + b(a - b)^{n-1} \\ &= (a - b)^2\det D_{n-2} + 2b(a - b)^{n-1} \\ &= \dots \\ &= (a - b)^{n-1}\det D_1 + (n - 1)b(a - b)^{n-1} \\ &= (a - b)^n + nb(a - b)^{n-1}\end{aligned}$$

Exercise 4.

1.

The system can be described as

$$Ax = b, \quad \text{where } A = \begin{pmatrix} 2 & -1 & 3 & 2 \\ 3 & -3 & 3 & 2 \\ 3 & -1 & -1 & 2 \\ 3 & -1 & 3 & -1 \end{pmatrix}, \quad b = \begin{pmatrix} 6 \\ 5 \\ 3 \\ 4 \end{pmatrix}$$

The determinant of A is (with calculations)

$$\det A = -70$$

Therefore,

$$x_1 = \frac{1}{\det A} \det \begin{pmatrix} 6 & -1 & 3 & 2 \\ 5 & -3 & 3 & 2 \\ 3 & -1 & -1 & 2 \\ 4 & -1 & 3 & -1 \end{pmatrix} = 1$$

$$x_2 = \frac{1}{\det A} \det \begin{pmatrix} 2 & 6 & 3 & 2 \\ 3 & 5 & 3 & 2 \\ 3 & 3 & -1 & 2 \\ 3 & 4 & 3 & -1 \end{pmatrix} = 1$$

$$x_3 = \frac{1}{\det A} \det \begin{pmatrix} 2 & -1 & 6 & 2 \\ 3 & -3 & 5 & 2 \\ 3 & -1 & 3 & 2 \\ 3 & -1 & 4 & -1 \end{pmatrix} = 1$$

$$x_4 = \frac{1}{\det A} \det \begin{pmatrix} 2 & -1 & 3 & 6 \\ 3 & -3 & 3 & 5 \\ 3 & -1 & -1 & 3 \\ 3 & -1 & 3 & 4 \end{pmatrix} = 1$$

2.

The system can be described as

$$Ax = b, \quad \text{where } A = \begin{pmatrix} 1 & 2 & 3 & -2 \\ 2 & -1 & -2 & -3 \\ 3 & 2 & -1 & 2 \\ 2 & -3 & 3 & 1 \end{pmatrix}, \quad b = \begin{pmatrix} 6 \\ 8 \\ 4 \\ -8 \end{pmatrix}$$

The determinant of A is (with calculations)

$$\det A = 360$$

Therefore,

$$x_1 = \frac{1}{\det A} \det \begin{pmatrix} 6 & 2 & 3 & -2 \\ 8 & -1 & -2 & -3 \\ 4 & 2 & -1 & 2 \\ -8 & -3 & 3 & 1 \end{pmatrix} = \frac{11}{10}$$

$$x_2 = \frac{1}{\det A} \det \begin{pmatrix} 1 & 6 & 3 & -2 \\ 2 & 8 & -2 & -3 \\ 3 & 4 & -1 & 2 \\ 2 & -8 & 3 & 1 \end{pmatrix} = \frac{37}{20}$$

$$x_3 = \frac{1}{\det A} \det \begin{pmatrix} 1 & 2 & 6 & -2 \\ 2 & -1 & 8 & -3 \\ 3 & 2 & 4 & 2 \\ 2 & -3 & -8 & 1 \end{pmatrix} = -\frac{9}{10}$$

$$x_4 = \frac{1}{\det A} \det \begin{pmatrix} 1 & 2 & 3 & 6 \\ 2 & -1 & -2 & 8 \\ 3 & 2 & -1 & 4 \\ 2 & -3 & 3 & -8 \end{pmatrix} = -\frac{39}{20}$$