

Honors Mathematics IV

RC 5

CHEN Xiwen

UM-SJTU Joint Institute

November 19, 2018

Table of contents

Complex Analysis

- Complex Differentiability

- Sets in the Complex Plane

- Power Series

Properties of Holomorphic Functions

- Complex Integrals

- Practical Integration

- Analytic Functions

Singularities and Poles

- Classification of Singularities

- Representation Near Poles

Residue Calculus

- The Residue Theorem

Exercises.

Complex Analysis

Complex Differentiability

Sets in the Complex Plane

Power Series

Properties of Holomorphic Functions

Complex Integrals

Practical Integration

Analytic Functions

Singularities and Poles

Classification of Singularities

Representation Near Poles

Residue Calculus

The Residue Theorem

Exercises.

Holomorphic Functions

Definition. A function $f : \mathbb{C} \rightarrow \mathbb{C}$ is **complex differentiable**, or **holomorphic**, as $z \in \mathbb{C}$ if

$$f'(z) := \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h}$$

exists.

- ▶ A holomorphic function is infinitely often differentiable.
- ▶ A holomorphic function is analytic.
- ▶ Any closed integral of a holomorphic function vanished.

Complex Differentiability

Cauchy-Riemann differential equations. The complex function

$$f : \mathbb{C} \rightarrow \mathbb{C}, \quad f(x + yi) = u(x, y) + iv(x, y)$$

is holomorphic only if its components u, v satisfy

$$u_x = v_y, \quad u_y = -v_x.$$

Complex Differentiability

Differential operators.

- Define

$$\frac{\partial}{\partial z} := \frac{1}{2} \left(\frac{\partial}{\partial x} + \frac{1}{i} \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial \bar{z}} := \frac{1}{2} \left(\frac{\partial}{\partial x} - \frac{1}{i} \frac{\partial}{\partial y} \right).$$

- If $f = u + iv : \mathbb{C} \rightarrow \mathbb{C}$ is holomorphic at $z \in \mathbb{C}$, then

$$f'(z) = \frac{\partial f}{\partial z} = 2 \frac{\partial u}{\partial z}, \quad \frac{\partial f}{\partial \bar{z}} = 0.$$

Complex Analysis

Complex Differentiability

Sets in the Complex Plane

Power Series

Properties of Holomorphic Functions

Complex Integrals

Practical Integration

Analytic Functions

Singularities and Poles

Classification of Singularities

Representation Near Poles

Residue Calculus

The Residue Theorem

Exercises.

Sets in the Complex Plane

Let $\Omega \subset \mathbb{C}$, then it is

► **Open:**

$$\forall z \in \Omega, \exists \varepsilon > 0 \text{ s.t. } B_\varepsilon(z) = \{w \in \mathbb{C} : |w - z| < \varepsilon\} \subset \Omega.$$

► **Bounded:** $\Omega \subset B_R(0)$ for some $R > 0$.

► **Compact** (denoted as K):

► **Definition.** Every sequence in K has a converging subsequence with limit in K .

► K is compact $\Leftrightarrow K$ is closed and bounded.

► **Disconnected:** For an open(closed) set Ω , there exists two open(closed) sets $\Omega_1, \Omega_2 \in \mathbb{C}$ such that

$$\Omega = \Omega_1 \cup \Omega_2, \quad \Omega_1 \cap \Omega_2 = \emptyset.$$

► **Region(domain):** An open and connected set in \mathbb{C} .

Complex Analysis

Complex Differentiability

Sets in the Complex Plane

Power Series

Properties of Holomorphic Functions

Complex Integrals

Practical Integration

Analytic Functions

Singularities and Poles

Classification of Singularities

Representation Near Poles

Residue Calculus

The Residue Theorem

Exercises.

Power Series

- **Convergence.** The series of the form

$$f(z) = \sum_{n=0}^{\infty} a_n z^n$$

with radius of convergence ρ : ($1/\rho = \lim_{n \rightarrow \infty} \sup |a_n|^{1/n}$.)

- converges if $|z| < \rho$.
 - diverges if $|z| > \rho$.
- **Differentiation.** Within the disc of convergence,

$$f'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}$$

is the derivative of f and has the same radius of convergence as f .

Trigonometric Functions

- Series definition.

$$\cos z = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!}, \quad \sin z = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!}.$$

- Euler formulas.

$$\cos z = \frac{e^{iz} + e^{-iz}}{2}, \quad \sin z = \frac{e^{iz} - e^{-iz}}{2i}.$$

Analytic Functions

Definition. A function f defined on an open set $\Omega \subset \mathbb{C}$ is said to be **analytic** (or have a power series expansion) at z_0 if there exists a power series centered at z_0 , with positive radius of convergence such that

$$f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$$

for all z in a neighborhood of z_0 . Moreover, with the same radius of convergence,

$$f'(z) = \sum_{n=0}^{\infty} n a_n(z - z_0)^{n-1}.$$

Note. A holomorphic function is automatically analytic.

Complex Analysis

Complex Differentiability

Sets in the Complex Plane

Power Series

Properties of Holomorphic Functions

Complex Integrals

Practical Integration

Analytic Functions

Singularities and Poles

Classification of Singularities

Representation Near Poles

Residue Calculus

The Residue Theorem

Exercises.

Integrals along Complex Curves

Definition. Let $\Omega \subset \mathbb{C}$ be an open set, f is holomorphic on Ω and $\mathcal{C}^* \subset \Omega$ and oriented smooth curve, the integral of f along \mathcal{C}^* is defined as

$$\int_{\mathcal{C}^*} f(z) dz := \int_I f(\gamma(t)) \cdot \gamma'(t) dt.$$

In particular, we have

► **Curve length:**

$$\ell(\mathcal{C}) := \left| \int_{\mathcal{C}} dz \right|.$$

► **Orientation:**

$$\int_{-\mathcal{C}^*} f(z) dz = - \int_{\mathcal{C}^*} f(z) dz.$$

► **Inequality:**

$$\left| \int_{\mathcal{C}^*} f(z) dz \right| \leq \ell(\mathcal{C}) \cdot \sup_{z \in \mathcal{C}} |f(z)|.$$

Primitives

Consider a continuous function f in an open set $\Omega \subset \mathbb{C}$ and an oriented curve \mathcal{C}^* .

- ▶ **2.2.5. Theorem.** If \mathcal{C}^* begins at w_1 and ends at w_2 and has primitive F ,

$$\int_{\mathcal{C}^*} f(z) dz = F(w_2) - F(w_1), \quad \oint_{\mathcal{C}} f(z) dz = 0.$$

- ▶ **2.2.9 and 10. Theorem.** f is holomorphic. $D \subset \Omega$ is a triangle or a rectangle whose interior is contained in Ω

$$\oint_D f(z) dz = 0.$$

- ▶ **2.2.11. Theorem.** f is holomorphic in an open disc $\Rightarrow f$ has a primitive in that disc.

Cauchy's Theorem

- ▶ 2.2.12. **Cauchy's Theorem.** f is holomorphic in a disc, then for any closed curve \mathcal{C} in that disc,

$$\oint_{\mathcal{C}} f(z) dz = 0.$$

- ▶ **Morera's Theorem (converse of Cauchy's theorem).** f is a continuous function in the open disc D , such that for any triangle T contained in D ,

$$\int_T f(z) dz = 0,$$

then f is holomorphic.

Complex Analysis

Complex Differentiability

Sets in the Complex Plane

Power Series

Properties of Holomorphic Functions

Complex Integrals

Practical Integration

Analytic Functions

Singularities and Poles

Classification of Singularities

Representation Near Poles

Residue Calculus

The Residue Theorem

Exercises.

Evaluating Real Integrals using Cauchy's Theorem

1. Transform the integrand into a complex function $f(z)$.
2. Select appropriate contour and use Cauchy's theorem to obtain

$$\int_{-R}^{-\varepsilon} f(z)dz + \underbrace{\int_{-C_\varepsilon} f(z)dz}_{I_\varepsilon} + \int_{\varepsilon}^R f(z)dz + \underbrace{\int_{C_R} f(z)dz}_{I_R} = 0.$$

3. Let $\varepsilon \rightarrow 0, R \rightarrow \infty$. In many cases,

$$I_R \rightarrow 0, \quad I_\varepsilon \rightarrow I_0.$$

4. Then

$$\int_{\mathbb{R}} f(x)dx = -I_0.$$

Note. $-I_0$ is usually not the desired integral. We need to take some transformations.

Evaluating Real Integrals using Cauchy's Theorem

Example 1. Show that if $\xi \in \mathbb{R}$, then

$$e^{-\pi\xi^2} = \int_{-\infty}^{\infty} e^{-\pi x^2} e^{-2\pi i x \xi} dx.$$

This means that $e^{-\pi x^2}$ is its own Fourier transform (later). If $\xi = 0$, the formula is the known integral

$$1 = \int_{-\infty}^{\infty} e^{-\pi x^2} dx.$$

Evaluating Real Integrals using Cauchy's Theorem

Remarks. Evaluate the following integrals.

1. Slide Example 2.2.14. Verify that

$$\int_0^{\infty} \frac{1 - \cos x}{x^2} dx = \frac{\pi}{2},$$

by integrating

$$f(z) = \frac{1 - e^{iz}}{z^2}.$$

2. Assignment 5.4 Show that

$$\int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}.$$

by integrating

$$f(z) = \frac{e^{iz} - 1}{2iz}.$$

Jordan's Lemma

2.2.15. Jordan's Lemma. Assume that for some $R_0 > 0$ the function $g : \mathbb{R} \setminus B_{R_0}(0) \rightarrow \mathbb{C}$ is holomorphic, Let

$$f(z) = e^{iaz} g(z), \quad \text{for some } a > 0.$$

Let

$$C_R = \{z \in \mathbb{C} : z = R \cdot e^{i\theta}, 0 \leq \theta \leq \pi\}$$

be a semi-circle segment in the upper half-plane and assume that

$$\sup_{0 \leq \theta \leq \pi} |g(Re^{i\theta})| \xrightarrow{R \rightarrow \infty} 0.$$

Then

$$\lim_{R \rightarrow \infty} \int_{C_R} f(z) dz = 0.$$

Cauchy's Integral Formula

2.2.16. Cauchy's Integral Formula. Suppose f is a holomorphic function in an open set $\Omega \subset \mathbb{C}$. If D is an open disc whose closure is contained in Ω , then

$$f(z) = \frac{1}{2\pi i} \oint_C \frac{f(\zeta)}{\zeta - z} d\zeta \quad \text{for all } z \in D,$$

where $C = \partial D$ is the (positively oriented) boundary circle of D . Furthermore,

$$f^{(n)}(z) = \frac{n!}{2\pi i} \oint_C \frac{f(\zeta)}{(\zeta - z)^{n+1}} d\zeta \quad \text{for all } z \in D.$$

Complex Analysis

Complex Differentiability

Sets in the Complex Plane

Power Series

Properties of Holomorphic Functions

Complex Integrals

Practical Integration

Analytic Functions

Singularities and Poles

Classification of Singularities

Representation Near Poles

Residue Calculus

The Residue Theorem

Exercises.

Analytic Functions.

2.2.20. Theorem. Suppose f is a holomorphic function in an open set Ω . If D is an open disc centered at z_0 whose closure is contained in Ω , then f has a power series expansion at z_0

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n.$$

for all $z \in D$ and the coefficients are given by

$$a_n = \frac{f^{(n)}(z_0)}{n!}, \quad n \in \mathbb{N}.$$

Analytic Functions

2.2.23. Theorem. Let $\Omega \subset \mathbb{C}$ be a region and $f, g : \Omega \rightarrow \mathbb{C}$ two holomorphic functions. Suppose $S \subset \Omega$ has an accumulation point that is contained in Ω and that

$$f(z) = g(z) \quad \text{for all } z \in S.$$

Then $f(z) = g(z)$ for all $z \in \Omega$.

Analytic Continuation. $g : \Omega \rightarrow \mathbb{C}, \Omega \subset \mathbb{C}$ is an analytic continuation of $f : M \rightarrow \mathbb{C}, M \subset \Omega$ if

- ▶ g is holomorphic.
- ▶ $g(z) = f(z)$ for $z \in M$.

Complex Analysis

Complex Differentiability

Sets in the Complex Plane

Power Series

Properties of Holomorphic Functions

Complex Integrals

Practical Integration

Analytic Functions

Singularities and Poles

Classification of Singularities

Representation Near Poles

Residue Calculus

The Residue Theorem

Exercises.

Classification of Singularities

Definition. $\Omega \subset \mathbb{C}$ is open, $z_0 \in \Omega$ and $f : \Omega \setminus \{z_0\} \rightarrow \mathbb{C}$ is holomorphic. (f has a **point singularity at z_0** .) The singularity is

- ▶ **removable**: there exists an analytic continuation $\tilde{f} : \Omega \rightarrow \mathbb{C}$.
(i.e., $\lim_{z \rightarrow z_0} f(z)$ exists.)
- ▶ a **pole**:
 1. $g = 1/f$ is holomorphic on $\Omega \setminus \{z_0\}$.
 2. g has a removable singularity at z_0 .
 3. $\tilde{g}(z_0) = 0$.
- ▶ **essential**: it is neither removable nor a pole.

2.3.5. Theorem. f is holomorphic in a connected open set Ω with a zero at $z_0 \in \Omega$ and does not vanish identically in Ω . In a neighborhood $U \subset \Omega$ of z_0 ,

$$f(z) = (z - z_0)^n g(z) \quad \text{for all } z \in U,$$

where g is non-vanishing and holomorphic.

- ▶ n, g are both unique.
- ▶ n is the **multiplicity** or **order** of the zero.
- ▶ The zero is **simple** if $n = 1$.

Poles

2.3.8. Theorem. $f : \Omega \rightarrow \mathbb{C}$ has a pole at $z_0 \in \Omega$, then in a neighborhood U of z_0 ,

$$f(z) = (z - z_0)^{-n} h(z) \quad \text{for all } z \in U,$$

where h is non-vanishing and holomorphic.

- ▶ n, h are both unique.
- ▶ n is the **multiplicity** or **order** of the pole.
- ▶ The pole is **simple** if $n = 1$.

Complex Analysis

Complex Differentiability

Sets in the Complex Plane

Power Series

Properties of Holomorphic Functions

Complex Integrals

Practical Integration

Analytic Functions

Singularities and Poles

Classification of Singularities

Representation Near Poles

Residue Calculus

The Residue Theorem

Exercises.

Representation Near Poles

2.3.11. Theorem. If $f : \Omega \rightarrow \mathbb{C}$ has a pole of order n at $z_0 \in \Omega$, then there exists a neighborhood $U \subset \Omega$ of z_0 , numbers $a_{-n}, \dots, a_{-1} \in \mathbb{C}$ and a holomorphic function $G : U \rightarrow \mathbb{C}$ such that

$$f(z) = \frac{a_{-n}}{(z - z_0)^n} + \frac{a_{-n+1}}{(z - z_0)^{n-1}} + \dots + \frac{a_{-1}}{z - z_0} + G(z)$$

for all $z \in U$.

► **Principal part:**

$$P(z) := \frac{a_{-n}}{(z - z_0)^n} + \frac{a_{-n+1}}{(z - z_0)^{n-1}} + \dots + \frac{a_{-1}}{z - z_0}.$$

► **Residue:**

$$\operatorname{res}_{z_0} f := a_{-1} = \frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} ((z - z_0)^n f(z)).$$

Complex Analysis

Complex Differentiability

Sets in the Complex Plane

Power Series

Properties of Holomorphic Functions

Complex Integrals

Practical Integration

Analytic Functions

Singularities and Poles

Classification of Singularities

Representation Near Poles

Residue Calculus

The Residue Theorem

Exercises.

The Residue Theorem

2.4.2. Theorem. Suppose that f is holomorphic in an open set containing a (positively oriented) toy contour \mathcal{C} and its interior, except for poles at the points z_1, \dots, z_N inside \mathcal{C} . Then

$$\int_{\mathcal{C}} f(z) dz = 2\pi i \sum_{k=1}^N \operatorname{res}_{z_k} f.$$

Residue Calculus

2.4.5. Theorem. Let P and Q be polynomials of degree m and n , respectively, where $n \geq m + 2$. If $Q(x) \neq 0$ for $x > 0$, if Q has a zero of order at most 1 at the origin and if

$$f(z) = \frac{z^\alpha P(z)}{Q(z)}, \quad 0 < \alpha < 1,$$

then

$$\int_0^\infty \frac{x^\alpha P(x)}{Q(x)} dx = \frac{2\pi i}{1 - e^{2\pi\alpha i}} \sum_{j=1}^k \operatorname{res}_{z_j} f,$$

where z_1, \dots, z_k are the nonzero poles of P/Q .

Evaluating Real Integrals Using Residue Calculus

1. Extend the real domain to complex domain.
 - ▶ Change $x \in \mathbb{R}$ to $z \in \mathbb{C}$.
 - ▶ Consider e^{iz} for $\sin x, \cos x$.
2. Find a suitable contour and the branch (if needed).
3. Find poles for $f(z)$.
4. Calculate residues for poles. (If the contour cannot be decided yet, find residue for all poles.)
 - ▶ Write out expression near poles.
 - ▶ Use

$$\operatorname{res}_{z_0} f = \frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} ((z - z_0)^n f(z)).$$

5. Write out residue theorem.
6. Save the desired integral and solve other parts.

Exercises.

Exercise 1. Suppose U and V are open sets in the complex plane. Prove that if $f : U \rightarrow V$ and $g : V \rightarrow \mathbb{C}$ are two functions that are differentiable (in the real sense, that is, as functions of the two real variables x and y), and $h = g \circ f$, then

$$\frac{\partial h}{\partial z} = \frac{\partial g}{\partial z} \frac{\partial f}{\partial z} + \frac{\partial g}{\partial \bar{z}} \frac{\partial f}{\partial \bar{z}}$$

and

$$\frac{\partial h}{\partial \bar{z}} = \frac{\partial g}{\partial z} \frac{\partial f}{\partial \bar{z}} + \frac{\partial g}{\partial \bar{z}} \frac{\partial f}{\partial z}.$$

This is the complex version of the chain rule.

Exercises.

Exercise 2. Show that in polar coordinates, the Cauchy-Riemann equations take the form

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta} \quad \text{and} \quad \frac{1}{r} \frac{\partial u}{\partial \theta} = -\frac{\partial v}{\partial r}.$$

Use these equations to show that the logarithm function defined by

$$\ln z = \ln r + i\theta \quad \text{where } z = e^{i\theta} \text{ with } -\pi < \theta < \pi$$

is holomorphic in the region $r > 0$ and $-\pi < \theta < \pi$.

Exercises.

Exercise 3. Evaluate the integrals

$$\int_0^{\infty} e^{-ax} \cos(bx) dx \quad \text{and} \quad \int_0^{\infty} e^{-ax} \sin(bx) dx$$

by integrating e^{-Az} , $A = \sqrt{a^2 + b^2}$, over an appropriate sector with angle ω , with $\cos \omega = a/A$.

Exercises.

Exercise 4. Let Ω be an open subset of \mathbb{C} and let $T \subset \Omega$ be a triangle whose interior is also contained in Ω . Suppose that f is a function holomorphic in Ω except possibly at a point w inside T . Prove that if f is bounded near w , then

$$\int_T f(z) dz = 0.$$

Exercises.

Exercise 5. If f is a holomorphic function on the strip $-1 < y < 1$, $x \in \mathbb{R}$ with

$$|f(z)| \leq A(1 + |z|)^\eta$$

for all z in that strip, where η is a fixed real number. Show that for each integer $n \geq 0$ there exists $A_n \geq 0$ so that

$$|f^{(n)}(x)| \leq A_n(1 + |x|)^\eta$$

for all $x \in \mathbb{R}$.

Thanks for your attention!