

Honors Mathematics IV

RC 8

CHEN Xiwen

UM-SJTU Joint Institute

December 7, 2018

Table of contents

Power Series Solutions

- The Power Series Ansatz
- Euler's Equation
- The Method of Frobenius

Bessel Functions

- The Bessel Equations
- Applications of Bessel Functions

Exercises

Power Series Solutions

The Power Series Ansatz

Euler's Equation

The Method of Frobenius

Bessel Functions

The Bessel Equations

Applications of Bessel Functions

Exercises

The Power Series Ansatz

Equation.

$$x'' + p(t)x' + q(t)x = 0$$

Solution.

1. Make power series ansatz of the solution:

$$x(t) = \sum_{k=0}^{\infty} a_k t^k.$$

2. Plug the ansatz into the equation using

$$x'(t) = \sum_{k=1}^{\infty} k a_k t^{k-1}, \quad x''(t) = \sum_{k=2}^{\infty} k(k-1) a_k t^{k-2}.$$

The Power Series Ansatz

Equation.

$$x'' + p(t)x' + q(t)x = 0$$

Solution.

3. Find the recurrence formula of a_k s.
4. Decide the first few terms (by initial conditions) to obtain two independent solutions $x_1(t), x_2(t)$.
5. Find $x(t)$ by

$$x(t) = x_1(t) + x_2(t).$$

The Power Series Ansatz

Example 1. Determine the terms up to x^5 in each of the two linearly independent power series solutions to

$$y'' + (2 - 4x^2)y' - 8xy = 0$$

centered at $x = 0$. Also find the radius of convergence of these solutions.

Power Series Solutions

The Power Series Ansatz

Euler's Equation

The Method of Frobenius

Bessel Functions

The Bessel Equations

Applications of Bessel Functions

Exercises

Euler's Equation

Euler's equation.

$$t^2 x'' + \alpha t x' + \beta x = 0, \quad \alpha, \beta \in \mathbb{R}.$$

Solution ($t > 0$). We make ansatz $x(t) = t^r$ and obtain

$$(r^2 + (\alpha - 1)r + \beta)t^r = 0, \quad r = -\frac{\alpha - 1}{2} \pm \frac{1}{2}\sqrt{(\alpha - 1)^2 - 4\beta}.$$

Then

► $(\alpha - 1)^2 - 4\beta > 0$. $r_1 \neq r_2 \in \mathbb{R}$.

$$x(t; c_1, c_2) = c_1 t^{r_1} + c_2 t^{r_2}, \quad c_1, c_2 \in \mathbb{R}.$$

Euler's Equation

Euler's equation.

$$t^2 x'' + \alpha t x' + \beta x = 0, \quad \alpha, \beta \in \mathbb{R}.$$

Solution ($t > 0$). We make ansatz $x(t) = t^r$ and obtain

$$(r^2 + (\alpha - 1)r + \beta)t^r = 0, \quad r = -\frac{\alpha - 1}{2} \pm \frac{1}{2}\sqrt{(\alpha - 1)^2 - 4\beta}.$$

Then

$$\blacktriangleright (\alpha - 1)^2 - 4\beta = 0. \quad r_1 = r_2 = \frac{1 - \alpha}{2}.$$

$$x(t; c_1, c_2) = c_1 t^{r_1} + c_2 t^{r_1} \ln t, \quad c_1, c_2 \in \mathbb{R}.$$

Euler's Equation

Euler's equation.

$$t^2 x'' + \alpha t x' + \beta x = 0, \quad \alpha, \beta \in \mathbb{R}.$$

Solution ($t > 0$). We make ansatz $x(t) = t^r$ and obtain

$$(r^2 + (\alpha - 1)r + \beta)t^r = 0, \quad r = -\frac{\alpha - 1}{2} \pm \frac{1}{2}\sqrt{(\alpha - 1)^2 - 4\beta}.$$

Then

► $(\alpha - 1)^2 - 4\beta < 0$. $r_1 = \lambda + i\mu, r_2 = \lambda - i\mu \in \mathbb{C}$, where

$$\lambda = \frac{1 - \alpha}{2}, \quad \mu = \frac{1}{2}\sqrt{4\beta - (\alpha - 1)^2}.$$

Then the two solutions are $x_1(t) = t^{r_1}, x_2(t) = t^{r_2}$ and

$$x(t; c_1, c_2) = c_1 t^\lambda \cos(\mu \ln t) + c_2 t^\lambda \sin(\mu \ln t), \quad c_1, c_2 \in \mathbb{R}.$$

Euler's Equation

Euler's equation.

$$t^2 x'' + \alpha t x' + \beta x = 0, \quad \alpha, \beta \in \mathbb{R}.$$

General solution. For $t > 0$ or $t < 0$,

$$x(t; c_1, c_2) = \xi(-t; c_1, c_2) = \begin{cases} c_1 |t|^{r_1} + c_2 |t|^{r_2}, \\ c_1 |t|^{r_1} + c_2 |t|^{r_1} \ln |t|, \\ c_1 |t|^\lambda \cos(\mu \ln |t|) + c_2 |t|^\lambda \sin(\mu \ln |t|). \end{cases}$$

The solutions may not be defined at $t = 0$.

Power Series Solutions

The Power Series Ansatz

Euler's Equation

The Method of Frobenius

Bessel Functions

The Bessel Equations

Applications of Bessel Functions

Exercises

Regular Singular Points

Definition. The equation

$$x'' + p(t)x' + q(t)x = 0$$

has a **regular singular point** at t_0 if

- ▶ $(t - t_0)p(t)$ and $(t - t_0)^2q(t)$ are analytic in a neighborhood of t_0 or, equivalently,
- ▶ we have convergent power series such that

$$p(t) = \frac{p_{-1}}{t - t_0} + \sum_{j=0}^{\infty} p_j(t - t_0)^j,$$

$$q(t) = \frac{q_{-2}}{(t - t_0)^2} + \frac{q_{-1}}{t - t_0} + \sum_{j=0}^{\infty} q_j(t - t_0)^j$$

in a neighborhood of t_0 .

The Method of Frobenius

Equation.

$$t^2 x'' + t(tp(t))x' + t^2 q(t)x = 0, \quad t > 0$$

Solution.

1. Make **Frobenius ansatz** and identify $tp(t)$, $t^2 q(t)$ and their coefficients,

$$x(t) = t^r \sum_{k=0}^{\infty} a_k t^k, \quad a_0 \neq 0.$$

2. Plug in the ansatz (or memorize) to obtain the **indicial equation**

$$F(r) = 0, \quad F(x) := x(x-1) + p_0 x + q_0,$$
$$a_m F(r+m) = - \sum_{k=0}^{m-1} (q_{m-k} + (r+k)p_{m-k}) a_k, \quad m \geq 1.$$

The Method of Frobenius

3. Find two independent solutions to the ODE. The first solution is (find by recurrent equation)

$$x_t(t) = t^{r_1} \sum_{k=0}^{\infty} a_k(r_1) t^k,$$

and

- ▶ $r_1 - r_2 \notin \mathbb{N}$. Find x_1, x_2 by two different recurrent equations given by r_1 and r_2 .
- ▶ $r_1 - r_2 \in \mathbb{N}$.
 - ▶ Both sides vanish in the indicial equation for $N \in \mathbb{N}$: choose an arbitrary a_N and find the second independent solution.
 - ▶ One side does not vanish for $N \in \mathbb{N}$:

$$x_2(t) = \frac{\partial}{\partial r} \left(t^r \sum_{k=0}^{\infty} a_k(r) t^k \right) \bigg|_{r=r_2} = c \cdot x_1(t) \ln t + t^{r_2} \sum_{k=0}^{\infty} a'_k(r_2) t^k,$$

where $c \in \mathbb{R}$ may vanish. If $r_1 = r_2$, then $c = 1$.

The Method of Frobenius

Example 2. Find a series solution about $x = 0$ of

$$x^2 y'' - xy' + (1 - x)y = 0.$$

The Method of Frobenius

Example 3. Find the first few terms of the second series solution about $x = 0$ of

$$x^2 y'' - xy' + (1 - x)y = 0.$$

Power Series Solutions

The Power Series Ansatz

Euler's Equation

The Method of Frobenius

Bessel Functions

The Bessel Equations

Applications of Bessel Functions

Exercises

The Bessel Functions

Bessel equations. The *Bessel equation* of order ν is given by

$$x^2 y'' + xy' + (x^2 - \nu^2)y = 0, \quad x \in \mathbb{R}.$$

The Bessel Equation of Order 1/2

Equation.

$$t^2 x'' + tx' + \left(t^2 - \frac{1}{4}\right)x = 0.$$

Solution (Frobenius method).

1. Find $tp(t) = 1$, $t^2q(t) = t^2 - 1/4$, which are analytic at $t = 0$ and

$$\begin{aligned} p_0 &= 1, & p_k &= 0 \quad (k \geq 1), \\ q_0 &= -\frac{1}{4}, & q_1 &= 0, & q_2 &= 1, & q_k &= 0 \quad (k \geq 3). \end{aligned}$$

2. Write and solve the indicial equation

$$F(r) = r(r-1) + p_0r + q_0 = \left(r + \frac{1}{2}\right)\left(r - \frac{1}{2}\right) = 0,$$

giving $r_1 = 1/2$ and $r_2 = -1/2$.

The Bessel Equation of Order 1/2

Equation.

$$t^2 x'' + tx' + \left(t^2 - \frac{1}{4}\right)x = 0.$$

Solution (Frobenius method).

3. Write out the recurrent relation:

$$a_m F(r+m) = - \sum_{k=0}^{m-1} (q_{m-k} + (r+k)p_{m-k}) a_k, \quad m \geq 1,$$

which gives

$$-a_1 F(r+1) = 0, \quad -a_m F(r+m) = a_{m-2}$$

for $m \geq 2$. Set $a_0 = 1$.

The Bessel Equation of Order 1/2

Equation.

$$t^2 x'' + tx' + \left(t^2 - \frac{1}{4}\right)x = 0.$$

Solution (Frobenius method).

4. Find the two independent solutions. ($r_1 - r_2 = 1$.)

► $r_1 = \frac{1}{2}$. For $m \geq 2$, the recurrent equation is

$$a_m = -\frac{a_{m-2}}{m(m+1)},$$

giving

$$a_{2k+1} = 0, \quad a_{2k} = \frac{1}{2k(2k+1)} \frac{1}{(2k-2)(2k-1)} a_{2(k-2)}.$$

Then $x_1(t)$ is given by

$$x_1(t) = t^{r_1} \sum_{m=0}^{\infty} a_m t^m = \frac{\sin t}{\sqrt{t}}.$$

The Bessel Equation of Order 1/2

Equation.

$$t^2 x'' + tx' + \left(t^2 - \frac{1}{4}\right)x = 0.$$

Solution (Frobenius method).

4. Find the two independent solutions. ($r_1 - r_2 = 1$.)

- $r_2 = -\frac{1}{2}$. Both sides of the recurrent relation vanish for $N = 1$ in $F(r_2 + N)$. So a_1 is arbitrary (set to 0). For $m \geq 2$, the recurrent relation is

$$a_m = -\frac{a_{m-2}}{m(m-1)},$$

giving

$$a_{2k+1} = 0, \quad a_{2k} = \frac{(-1)^k}{(2k)!}.$$

Then $x_2(t)$ is given by

$$x_2 = t^{r_2} \sum_{m=0}^{\infty} a_m t^m = \frac{\cos t}{\sqrt{t}}.$$

The Bessel Equation of Order 0

Equation.

$$t^2 x'' + tx' + t^2 x = 0.$$

Solution (Frobenius method).

1. Find $tp(t) = 1$, $t^2q(t) = t^2$, which are analytic at $t = 0$ and

$$\begin{aligned} p_0 &= 1, & p_1 &= 0, & p_k &= 0 \quad (k \geq 2), \\ q_0 &= 0, & q_1 &= 0, & q_2 &= 1, & q_k &= 0 \quad (k \geq 3). \end{aligned}$$

2. Write and solve the indicial equation

$$F(r) = r(r-1) + p_0 r + q_0 = r^2 = 0,$$

giving $r_1 = r_2 = 0$.

The Bessel Equation of Order 0

Equation.

$$t^2 x'' + tx' + t^2 x = 0.$$

Solution (Frobenius method).

3. Write out the recurrent relation:

$$a_m F(r+m) = - \sum_{k=0}^{m-1} (q_{m-k} + (r+k)p_{m-k}) a_k, \quad m \geq 1,$$

which gives

$$(r+1)^2 a_1 = 0, \quad a_m(r+m)^2 = -a_{m-2}(r)$$

for $m \geq 2$. Then we know $a_1 = 0$ and set $a_0 = 1$.

The Bessel Equation of Order 0

Equation.

$$t^2 x'' + tx' + t^2 x = 0.$$

Solution (Frobenius method).

4. Find the two independent solutions. $r_1 = r_2 = 0$.

► x_1 . From the recurrent relations we know that

$$a_{2k+1} = 0, \quad a_{2k} = \frac{1}{(r+2k)^2} \frac{1}{(r+2k-2)^2} a_{2k-4}(r).$$

Then $x_1(t)$ is given by

$$x_1(t) = t^{r_1} \sum_{m=0}^{\infty} a_m t^m = \sum_{k=0}^{\infty} \frac{(-1)^k}{2^{2k}(k!)^2} t^{2k}.$$

The Bessel Equation of Order 0

Equation.

$$t^2 x'' + tx' + t^2 x = 0.$$

Solution (Frobenius method).

4. Find the two independent solutions. $r_1 = r_2 = 0$.

► x_2 . The second solution is given by

$$\begin{aligned} x_2(t) &= \left. \frac{d}{dt} \left(t^r \sum_{k=0}^{\infty} a_k(r) t^k \right) \right|_{r=r_2=0} \\ &= x_1(t) \ln t + \sum_{k=0}^{\infty} a'_k(0) t^k, \end{aligned}$$

with

$$a_{2k}(r) = \frac{(-1)^k}{(2+r)^2(4+r)^2 \cdots (2k+r)^2}.$$

The Bessel Equation of Order 0

Equation.

$$t^2 x'' + tx' + t^2 x = 0.$$

Solution (Frobenius method).

4. Find the two independent solutions. $r_1 = r_2 = 0$.

► x_2 . Then we have

$$\frac{a'_{2k}(r)}{a_{2k}(r)} = -2 \sum_{j=1}^k \frac{1}{2j+r}.$$

evaluated at $r_2 = 0$,

$$a'_{2k}(0) = - \left(1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{k} \right) 2_{2n}(0).$$

Therefore,

$$x_2(t) = x_1(t) \ln t + \sum_{k=0}^{\infty} \left(\sum_{j=1}^k \frac{1}{j} \right) \frac{(-1)^{k+1}}{2^{2k}(k!)^2} t^{2k}.$$

The Bessel Functions

Generating function. The function

$$\Psi(x, t) = e^{\frac{x}{2}(t-1/t)}.$$

is a *generating function* for the Bessel functions of the first kind and integer order. More precisely,

$$e^{\frac{x}{2}(t-1/t)} = \sum_{n=-\infty}^{\infty} J_n(x) t^n.$$

The Bessel Functions

Bessel functions of integer order. For $n \in \mathbb{N}$ the function $J_n : \mathbb{R} \rightarrow \mathbb{R}$ is given by

$$J_n(x) := \sum_{r=0}^{\infty} \frac{(-1)^r}{(r+n)!r!} \left(\frac{x}{2}\right)^{2r+n}, \quad J_{-n} = (-1)^n J_n.$$

General Bessel functions. The Bessel functions $J_\nu(x)$ of order $\nu \in \mathbb{R}$ is given by

$$J_\nu(x) = \sum_{r=0}^{\infty} \frac{(-1)^r}{\Gamma(\nu+r+1)r!} \left(\frac{x}{2}\right)^{2r+\nu},$$

where Γ is the Gamma function.

The Bessel Functions


Trigonometric approximation.

$$J_n(x) \sim \frac{1}{\sqrt{x}} \cos\left(x - \frac{2n+1}{4}\pi\right), \quad \text{as } x \rightarrow \infty.$$

Graph of some Bessel functions.

The graph shows

- ▶ J_0 (red),
- ▶ J_1 (yellow),
- ▶ J_2 (green),
- ▶ J_3 (light blue),
- ▶ J_4 (dark blue),
- ▶ J_5 (violet).



The Bessel Functions

Recurrence relations.

$$\begin{aligned}2nJ_n(x) &= xJ_{n+1}(x) + xJ_{n-1}(x), \\ J'_n(x) &= \frac{1}{2}(J_{n-1}(x) - J_{n+1}(x)).\end{aligned}$$

For $\nu \in \mathbb{R}$.

$$\frac{d}{dx}(x^\nu J_\nu(x)) = x^\nu J_{\nu-1}(x), \quad \frac{d}{dx}(x^{-\nu} J_\nu(x)) = -x^{-\nu} J_{\nu+1}(x).$$

Power Series Solutions

The Power Series Ansatz

Euler's Equation

The Method of Frobenius

Bessel Functions

The Bessel Equations

Applications of Bessel Functions

Exercises

Applications of Bessel Functions

1. Use substitution or separation of variables ansatz to obtain a Bessel function

$$x^2 y'' + xy' + (x^2 - \nu^2)y = 0, \quad x \in \mathbb{R}.$$

2. Once the Bessel function is obtained, we can directly write out the solution depending on the practical conditions:
 - ▶ Solution is finite (most common):

$$y(x) = c \cdot J_\nu(x).$$

- ▶ Solution is infinite:

$$y(x) = c \cdot Y_\nu(x).$$

3. Substitute back for the original variables and insert boundary/initial conditions.

The Suspended Chain

Model equation. PDE with boundary condition at $x = l$:

$$\frac{1}{g} u_{tt}(x, t) = x \cdot u_{xx}(x, t) + u_x(x, t), \quad y(l) = 0.$$

Solution.

1. Suppose u is periodic with respect to time:

$$u(x, t) = y(x) \cdot e^{i\omega t}$$

for some frequency ω , then we obtain the ODE

$$\frac{\partial}{\partial x}(x \cdot y') + \frac{\omega^2}{g} y = 0.$$

2. Use substitution

$$x = \frac{gz^2}{4\omega^2}, \quad z = 2\omega \sqrt{\frac{x}{g}}.$$

The Suspended Chain

Model equation. PDE with boundary condition at $x = l$:

$$\frac{1}{g} u_{tt}(x, t) = x \cdot u_{xx}(x, t) + u_x(x, t), \quad y(l) = 0.$$

Solution.

3. Obtain the Bessel equation of order zero:

$$zw'' + w' + zw = 0.$$

4. Write out the finite solution for the Bessel equation

$$w(z) = c \cdot J_0(z).$$

5. Substitute back for the original variables and insert the boundary conditions:

$$y(x) = c \cdot J_0 \left(2\omega \sqrt{\frac{x}{g}} \right), \quad \omega = \frac{1}{2} \sqrt{\frac{g}{l}} \cdot \alpha_{0,n}.$$

Airy's Equation

Airy's equation.

$$y'' + xy = 0.$$

Solution.

1. Use substitution

$$u(t) = x^{-1/2}y(x), \quad t = \frac{2}{3}x^{3/2}$$

to obtain the Bessel equation of order $\nu = 1/3$:

$$\begin{aligned} \frac{du}{dt} &= \frac{du}{dx} \cdot \frac{dx}{dt} = -\frac{1}{2}x^{-2}y + x^{-1}\frac{dy}{dx}, \\ \frac{d^2u}{dt^2} &= \frac{d}{dt} \left(\frac{du}{dt} \right) = x^{-7/2}y - \frac{3}{2}x^{-5/2}\frac{dy}{dx} + x^{-3/2}\frac{d^2y}{dx^2}. \end{aligned}$$

Airy's Equation

Airy's equation.

$$y'' + xy = 0.$$

Solution.

1. Use substitution

$$u(t) = x^{-1/2}y(x), \quad t = \frac{2}{3}x^{3/2}$$

to obtain the Bessel equation of order $\nu = 1/3$: Then

$$\frac{dy}{dx^2} = x^{3/2} \frac{d^2u}{dt^2} - \frac{1}{4}x^{-3/2}u + \frac{3}{2} \frac{du}{dt},$$

substituting $t = \frac{2}{3}x^{3/2}$, we then have

$$t^2 u'' + tu' + (t^2 - \frac{1}{9})u = 0.$$

Airy's Equation

Airy's equation.

$$y'' + xy = 0.$$

Solution.

2. Write out the general solution using Bessel functions:

$$u(t) = c_1 J_{1/3}(t) + c_2 J_{-1/3}(t).$$

3. Substitute back for the original variables.

$$y(x) = c_1 \cdot x^{1/2} J_{1/3} \left(\frac{2}{3} x^{3/2} \right) + c_2 \cdot x^{1/2} J_{-1/3} \left(\frac{2}{3} x^{3/2} \right).$$

Transforming an Equation into a Bessel's Equation

A Bessel function is of the form

$$t^2 u'' + tu' + (t^2 - \nu^2)u = 0.$$

Using the substitution $t = ax^b$, $u = yx^c$, we have

$$y = x^{-c}u(t(x)), \quad \frac{dt}{dx} = abx^{b-1}$$

and thus

$$y' = -cx^{-c-1}u + abx^{b-c-1}u'$$

$$y'' = c(c+1)x^{-c-2}u + ab(b-2c-1)x^{b-c-2}u' + a^2b^2x^{2b-c-2}u''.$$

Transforming an Equation into a Bessel's Equation

Solving for u , u' and u'' , we have

$$u = yx^c, \quad u' = \frac{y' + cx^{-1}y}{abx^{b-c-1}},$$
$$u'' = \frac{y'' - c(c+1)yx^{-2} - ab(b-2c-1)x^{b-c-2} \cdot \frac{y' + cx^{-c-1}u}{abx^{b-c-1}}}{a^2b^2x^{2b-c-2}}$$

Therefore, we can transform the original equation into

$$x^2y'' + x(2c+1)y' + (a^2b^2x^{2b} - b^2\nu^2 + c^2)y = 0.$$

Differential equations of this form can be transformed into a Bessel's equation using substitution

$$t = ax^b, \quad u = yx^c.$$

Transforming an Equation into a Bessel's Equation

Example. For Airy's Equation, we have

$$x^2 y'' + x^3 y = 0,$$

where $c = -\frac{1}{2}$, $b = \frac{3}{2}$ and $a = \frac{2}{3}$, therefore, the transformation is

$$t = ax^b = \frac{2}{3}x^{3/2}, \quad u(t) = yx^c = x^{-1/2}y.$$

and the order is given by

$$c^2 = b^2 \nu^2 \quad \Rightarrow \quad \nu = \frac{1}{3}.$$

Exercises

Exercise 1. The *Legendre differential equation of order* $\lambda \in \mathbb{R}$ is given by

$$(1 - x^2)y'' - 2xy' + \lambda(\lambda + 1)y = 0.$$

1. Find the general solution of the equation in power series form.
2. Verify that if $\lambda \in \mathbb{N}$ there exists a non-zero polynomial solution.

Exercises

Exercise 2. Use the method of Frobenius to solve

$$5x^2y'' + x(1+x)y' - y = 0.$$

Thanks for your attention!