

VE401 Probabilistic Methods in Eng.

RC 7

CHEN Xiwen

UM-SJTU Joint Institute

April 23, 2020

Table of contents

Simple Linear Regression

- Simple Linear Regression Model

- Distributions of Estimators, CIs, and Hypothesis Tests

- Predictions and Model Analysis

- Simple Linear Regression using Software

Multiple Linear Regression

Simple Linear Regression

Simple Linear Regression Model

Distributions of Estimators, CIs, and Hypothesis Tests

Predictions and Model Analysis

Simple Linear Regression using Software

Multiple Linear Regression

Simple Linear Regression Model

Model. With x to be the parameter, the mean of the *response* $Y|x$ is given by

$$\mu_{Y|x} = \beta_0 + \beta_1 x \quad \text{for some } \beta_0, \beta_1 \in \mathbb{R},$$

which is equivalent to

$$Y|x = \beta_0 + \beta_1 x + E,$$

where $E[E] = 0$. We want to find estimators

$$\begin{aligned} B_0 &:= \widehat{\beta_0} = \text{estimator for } \beta_0, & b_0 &= \text{estimate for } \beta_0, \\ B_1 &:= \widehat{\beta_1} = \text{estimator for } \beta_1, & b_1 &= \text{estimate for } \beta_1. \end{aligned}$$

Simple Linear Regression Model

Model. Considering X as a parameter, we have a random sample of size n of $(x, Y|x)$.

$$Y_i := Y|x_i, \quad i = 1, \dots, n.$$

For each measurement y_i , we have a **residual** given by

$$y_i = b_0 + b_1 x_i + e_i.$$

Assumptions.

- ▶ For each value of x , the random variable follows a normal distribution with variance σ^2 and mean $\mu_{Y|x} = \beta_0 + \beta_1 x$.
- ▶ The random variables $Y|x_1$ and $Y|x_2$ are independent if $x_1 \neq x_2$.

Least Squares Estimation

LSE. We have the *error sum of squares*

$$SS_E := \sum_{i=1}^n e_i^2 = \sum_{i=1}^n (y_i - (b_0 + b_1 x_i))^2.$$

To minimize it, we take

$$\begin{aligned}\frac{\partial SS_E}{\partial b_0} &= -2 \sum_{i=1}^n (y_i - b_0 - b_1 x_i) = 0, \\ \frac{\partial SS_E}{\partial b_1} &= -2 \sum_{i=1}^n (y_i - b_0 - b_1 x_i) x_i = 0.\end{aligned}$$

Least Squares Estimation

LSE. We have

$$b_1 = \frac{S_{xy}}{S_{xx}}, \quad b_0 = \bar{y} - b_1 \bar{x},$$

where

$$S_{xx} = \sum_{i=1}^n (x_i - \bar{x})^2 = \sum_{i=1}^n x_i^2 - \frac{1}{n} \left(\sum_{i=1}^n x_i \right)^2 = \sum_{i=1}^n (x_i - \bar{x})x_i,$$

$$S_{yy} = \sum_{i=1}^n (y_i - \bar{y})^2 = \sum_{i=1}^n y_i^2 - \frac{1}{n} \left(\sum_{i=1}^n y_i \right)^2 = \sum_{i=1}^n (y_i - \bar{y})y_i,$$

$$\begin{aligned} S_{xy} &= \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) = \sum_{i=1}^n (x_i - \bar{x})y_i = \sum_{i=1}^n (y_i - \bar{y})x_i \\ &= \sum_{i=1}^n x_i y_i - \frac{1}{n} \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n y_i \right). \end{aligned}$$

Simple Linear Regression

Simple Linear Regression Model

Distributions of Estimators, CIs, and Hypothesis Tests

Predictions and Model Analysis

Simple Linear Regression using Software

Multiple Linear Regression

Distribution of B_1

Theorem. The least squares estimator B_1 for β_1 follows a normal distribution with

$$E[B_1] = \beta_1, \quad \text{Var}[B_1] = \frac{\sigma^2}{\sum (x_i - \bar{x})^2}.$$

Proof. Knowing $Y|x_i = \beta_0 + \beta_1 x_i + E$ and $E[E_i] = 0$, the expectation is given by

$$\begin{aligned} E[B_1] &= E \left[\frac{1}{S_{xx}} \sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y}) \right] = E \left[\frac{1}{S_{xx}} \sum (x_i - \bar{x}) Y_i \right] \\ &= \frac{1}{S_{xx}} \left(\sum (x_i - \bar{x}) E[\beta_0 + \beta_1 x_i + E_i] \right) \\ &= \frac{1}{S_{xx}} \left(\beta_1 \sum (x_i - \bar{x}) x_i \right) \\ &= \beta_1. \end{aligned}$$

Distribution of B_1

Theorem. The least squares estimator B_1 for β_1 follows a normal distribution with

$$E[B_1] = \beta_1, \quad \text{Var}[B_1] = \frac{\sigma^2}{\sum (x_i - \bar{x})^2}.$$

Proof. Similarly, given $\text{Var}[E_i] = \sigma^2$, the variance is given by

$$\begin{aligned} \text{Var}[B_1] &= \frac{1}{S_{xx}^2} \text{Var} \left[\sum (x_i - \bar{x}) Y_i \right] \\ &= \frac{1}{S_{xx}^2} \sum (x_i - \bar{x})^2 \text{Var}[\beta_0 + \beta_1 x_i + E_i] \\ &= \frac{\sigma^2}{\sum (x_i - \bar{x})^2} \\ &= \frac{\sigma^2}{S_{xx}}. \end{aligned}$$

Distribution of B_0

Theorem. The least squares estimator B_0 for β_0 follows a normal distribution with

$$E[B_0] = \beta_0, \quad \text{Var}[B_0] = \frac{\sigma^2 \sum x_i^2}{n \sum (x_i - \bar{x})^2}.$$

Proof. Using $\sum (x_i - \bar{x}) = 0$, the expectation is given by

$$\begin{aligned} E[B_0] &= E \left[\bar{Y} - \frac{\bar{x}}{S_{xx}} \sum (x_i - \bar{x}) Y_i \right] \\ &= \beta_0 + \beta_1 \bar{x} - \frac{\bar{x}}{S_{xx}} \sum (x_i - \bar{x}) (\beta_0 + \beta_1 x_i) \\ &= \beta_0 + \beta_1 \bar{x} - \frac{\bar{x}}{S_{xx}} \sum (x_i - \bar{x}) x_i \beta_1 \\ &= \beta_0. \end{aligned}$$

Distribution of B_0

Theorem. The least squares estimator B_0 for β_0 follows a normal distribution with

$$E[B_0] = \beta_0, \quad \text{Var}[B_0] = \frac{\sigma^2 \sum x_i^2}{n \sum (x_i - \bar{x})^2}.$$

Proof. Similarly, using $\text{Var}[\bar{E}] = \sigma^2/n$, the variance is given by

$$\begin{aligned} \text{Var}[B_0] &= \text{Var}\left[\bar{Y} - \frac{\bar{x}}{S_{xx}} \sum (x_i - \bar{x}) Y_i\right] \\ &= \text{Var}[\beta_0 + \beta_1 \bar{x} + \bar{E}] + \frac{\bar{x}^2}{S_{xx}^2} \sum (x_i - \bar{x})^2 \text{Var}[\beta_0 + \beta_1 x_i + E_i] \\ &= \frac{\sigma^2}{n} + \frac{\bar{x}^2}{S_{xx}} \sigma^2 \\ &= \frac{S_{xx} + \bar{x}^2}{n S_{xx}} \sigma^2 \\ &= \frac{\sum x_i^2}{n S_{xx}} \sigma^2. \end{aligned}$$

Simple Linear Regression

Simple Linear Regression Model

Distributions of Estimators, CIs, and Hypothesis Tests

Predictions and Model Analysis

Simple Linear Regression using Software

Multiple Linear Regression

Simple Linear Regression

Simple Linear Regression Model

Distributions of Estimators, CIs, and Hypothesis Tests

Predictions and Model Analysis

Simple Linear Regression using Software

Multiple Linear Regression

Thanks for your attention!