



Report

Nara Avila Moraes

Physics Institute
IFUSP

2026 / 09 / 04

Summary

① Tasks Status

② Results

③ Selected Article

General Tasks Status

Task	Status	
Correlation Inference Study - Classical Approach		
Correlation Inference Study - Quantum Learning Approach		
2026-11 Paper		
2026-12 LASNPA Presentation		

Correlated Uncertainty Study Tasks Status

Task	Status	Link
Alice ins1222333 Studies		
↪ Uncertainty correlation effect	✓	Plots
RHIC Au-Au/d-Au Studies JETSCAPE		
↪ Uncertainty correlation effect	✓	Plots
Atlas ins1759506 studies		
↪ Uncertainty correlation inference	🕒	Notebook
Alice PbPb2760 Scaled Spectra studies		
↪ Uncertainty correlation inference	🕒	Notebook
Synthetic data studies		
↪ Uncertainty correlation inference	🕒	Notebook

Atlas ins1759506 studies




Figure: GP fit: fixed ell = 1.0

Atlas ins1759506 studies




Figure: GP fit: optimized ell

Atlas ins1759506 studies




Figure: Ratio between white noise and systematic uncorrelated noise

Alice PbPb2760 Scaled Spectra studies

ALICE Pb-Pb 2.76 TeV Grad, 0-5%




Figure: GP fit: fixed ell = 0.2

Alice PbPb2760 Scaled Spectra studies




Figure: GP fit: optimized ell

Selected Article

Responses of multiparticle observables to multidimensional nuclear deformation in relativistic heavy-ion collisions

Ying Shan Zhao¹ and Yifeng Sun^{1,*}

¹*State Key Laboratory of Dark Matter Physics, Shanghai Key Laboratory for Particle Physics and Cosmology,
Key Laboratory for Particle Astrophysics and Cosmology (MOE),
School of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai 200240, China*

Relativistic heavy-ion collisions provide a unique opportunity to probe ground-state nuclear structure through its imprint on the initial collision geometry. We investigate how multiparticle observables respond to combined variations of quadrupole deformation, triaxiality, and hexadecapole deformation, using $^{129}\text{Xe}+^{129}\text{Xe}$ collisions as a representative testing ground. We perform a joint analysis in the three-dimensional $(\beta_2, \gamma, \beta_4)$ parameter space and construct initial-state estimators for several flow and mean transverse momentum correlation observables. At the initial-state level, ρ_2 is primarily sensitive to β_2 and γ , with its sensitivity to γ enhanced at nonzero β_2 . The nonlinear response coefficient $\chi_{4,22}$ is predominantly sensitive to β_4 , while its dependence on β_2 and γ remains comparatively weak. Higher-order correlators exhibit more complex multidimensional response patterns; in particular, ρ_{224} shows a dependence on γ and β_4 that becomes more pronounced at finite β_2 . We further employ the iEBE-VISHNU hybrid model to examine whether these deformation sensitivities survive the subsequent dynamical evolution. The final-state calculations indicate that the sensitivity of ρ_2 to β_2 and γ is largely preserved, whereas the β_4 sensitivity of $\chi_{4,22}$ is substantially reduced. For the other higher-order observables, the initial-state sensitivities are modified by the evolution or cannot be resolved with the present statistics.

Figure: Multidimensional nuclear structure deformation study.