

Topic

Convolution and Correlation

Lecture by: Fatemeh Pishdadian

Convolution

- **convolution** is a mathematical operator which takes two functions x and h and produces a third function that represents the amount of overlap between h and a reversed and translated version of x .
- In signal processing, one of the functions (h) is taken to be a fixed filter *impulse response*, and the other (x) the input signal.

$$(h * x)(t) \equiv \int_{-\infty}^{\infty} h(\tau) x(t - \tau) d\tau$$

↑
Convolution
operator

Discrete Convolution

- **convolution** is a mathematical operator which takes two functions x and h and produces a third function that represents the amount of overlap between h and a reversed and translated version of x .
- In signal processing, one of the functions (h) is taken to be a fixed filter *impulse response*, and the other (x) the input signal.

$$(h * x)[n] \equiv \sum_{k=-\infty}^{\infty} h[k]x[n-k]$$

↑
Convolution
operator

Convolution In Python Code

```
import numpy as np

def convolution(A,B):


    lengthA=np.size(A)
    lengthB=np.size(B)

    C = np.zeros(lengthA + lengthB -1)


    for m in np.arange(lengthA):
        for n in np.arange(lengthB):
            C[m+n] = C[m+n] + A[m]*B[n]

    return C
```


Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution




Let's look at Convolution



Let's look at Convolution



Let's look at Convolution



Cross-correlation

- **Cross-correlation** is a measure of similarity of two functions at time-lag t applied to one of them. It is a LOT like convolution...

$$(h \text{ 🍏 } x)(t) \equiv \int_{-\infty}^{\infty} h^*(\tau) x(t + \tau) d\tau$$

Means "complex conjugate of h"

↑
Cross-correlation operator
Should be a star
Couldn't find "star" in my font

VERY Similar

- Convolution

$$(h * x)(t) \equiv \int_{-\infty}^{\infty} h(\tau) x(t - \tau) d\tau$$

- Cross-correlation

$$(h \text{ 🍏 } x)(t) \equiv \int_{-\infty}^{\infty} h^*(\tau) x(t + \tau) d\tau$$

Cross-correlation in Python Code

We can easily implement cross correlation with convolution as follows:

```
def crosscorrelation(A,B):  
    return convolution(np.conj(A),B[::-1])
```

Better yet, use the built in Python functions...

```
np.convolve(A,B,"full")    # for convolution  
np.correlate(A,B,"full")   # for cross correlation
```

Auto-correlation

- **Auto-correlation** is a measure of similarity of a function to itself at time-lag t . It is a special case of cross-correlation (cross-correlation of a function with itself).


$$(x \text{ 🍏 } x)(t) \equiv \int_{-\infty}^{\infty} x^*(\tau) x(t + \tau) d\tau$$

Means "complex conjugate of f"

↑

Cross correlation
Should be a star
Couldn't find "star" in my font

Relating them all



Convolution and Fourier transform

- An important property of the Fourier transform: converts convolution in the time domain into multiplication in the frequency domain.

Convolution... $y(t) = h(t) * x(t)$

In the time domain: $y(t) = \int h(\tau)x(t - \tau) d\tau$

In frequency domain: $Y(\omega) = H(\omega)X(\omega)$