

Magnetars!

Kristen E. Gardner

UTU - Department of Physics and Astronomy

What is a Magnetar?

Definition:



Figure: Artistic depiction of a magnetar

What is a Magnetar?

Definition:

- neutron star



Figure: Artistic depiction of a magnetar

What is a Magnetar?

Definition:

- neutron star
- massive magnetic field ($\geq 10^{13}$ G)



Figure: Artistic depiction of a magnetar

Discovery of Magnetars

Definition:



Figure: Artistic depiction of a magnetar

Discovery of Magnetars

Definition:

- proposed in 1992 by Duncan Thompson [1]



Figure: Artistic depiction of a magnetar

Discovery of Magnetars

Definition:

- proposed in 1992 by Duncan Thompson [1]
- SGR 0525-66 found in 1979 (only known outside MW)



Figure: Artistic depiction of a magnetar

Discovery of Magnetars

Definition:

- proposed in 1992 by Duncan Thompson [1]
- SGR 0525-66 found in 1979 (only known outside MW)
- short X-Ray bursts



Figure: Artistic depiction of a magnetar

Discovery of Magnetars

Definition:

- proposed in 1992 by Duncan Thompson [1]
- SGR 0525-66 found in 1979 (only known outside MW)
- short X-Ray bursts
- selection bias



Figure: Artistic depiction of a magnetar

Basic Properties

Description:



Cases:

Figure: Artistic depiction of a magnetar

Basic Properties

Description:

- young neutron star



Cases:

Figure: Artistic depiction of a magnetar

Basic Properties

Description:

- young neutron star
- spinning down, timescale ~ 1 kyr



Cases:

Figure: Artistic depiction of a magnetar

Basic Properties

Description:

- young neutron star
- spinning down, timescale ~ 1 kyr
- high-B field ($\geq 10^{13}$ G)



Cases:

Figure: Artistic depiction of a magnetar

Basic Properties

Description:

- young neutron star
- spinning down, timescale ~ 1 kyr
- high-B field ($\geq 10^{13}$ G)
- rotational period range of 2-12s



Cases:

Figure: Artistic depiction of a magnetar

Basic Properties

Description:

- young neutron star
- spinning down, timescale ~ 1 kyr
- high-B field ($\geq 10^{13}$ G)
- rotational period range of 2-12s



Cases:

- bi-modal population of transients and persistents

Figure: Artistic depiction of a magnetar

Basic Properties

Description:

- young neutron star
- spinning down, timescale ~ 1 kyr
- high-B field ($\geq 10^{13}$ G)
- rotational period range of 2-12s



Cases:

- bi-modal population of transients and persistents
- cross-species magnetar/pulsar

Figure: Artistic depiction of a magnetar

Distinctions

Neutron Star

- radius on order of 10 km
- 1.1 - 2.16 solar masses
- $10^9 - 10^{13}$ G (for non-magnetars)

Magnetars

Radio Pulsars

Distinctions

Neutron Star

- radius on order of 10 km
- 1.1 - 2.16 solar masses
- $10^9 - 10^{13}$ G (for non-magnetars)

Magnetars

- 2-12s rotational period

Radio Pulsars

Distinctions

Neutron Star

- radius on order of 10 km
- 1.1 - 2.16 solar masses
- $10^9 - 10^{13}$ G (for non-magnetars)

Magnetars

- 2-12s rotational period

Radio Pulsars

- ~ 100 ms rotational period

Distinctions

Neutron Star

- radius on order of 10 km
- 1.1 - 2.16 solar masses
- $10^9 - 10^{13}$ G (for non-magnetars)

Magnetars

- 2-12s rotational period
- spin-down \rightarrow quiescent X-ray emission and gamma-ray bursts

Radio Pulsars

- ~ 100 ms rotational period

Distinctions

Neutron Star

- radius on order of 10 km
- 1.1 - 2.16 solar masses
- $10^9 - 10^{13}$ G (for non-magnetars)

Magnetars

- 2-12s rotational period
- spin-down \rightarrow quiescent X-ray emission and gamma-ray bursts

Radio Pulsars

- ~ 100 ms rotational period
- spin-down \rightarrow radio emission, and non-thermal X/gamma-ray radiation

Neutron Star Types




Figure: Neutron star types, courtesy of JPL [cite here]

Distribution in Space

Observations:




Figure: Top down view of the Milky Way with known Magnetars (and distance uncertainties) in red.

Distribution in Space

Observations:

- generally within the galactic plane




Figure: Top down view of the Milky Way with known Magnetars (and distance uncertainties) in red.

Distribution in Space

Observations:

- generally within the galactic plane
- mean speed of 200 km s^{-1} ,
 $\sigma = 100 \text{ km s}^{-1}$




Figure: Top down view of the Milky Way with known Magnetars (and distance uncertainties) in red.

Distribution in Space

Observations:

- generally within the galactic plane
- mean speed of 200 km s^{-1} ,
 $\sigma = 100 \text{ km s}^{-1}$
- selection bias




Figure: Top down view of the Milky Way with known Magnetars (and distance uncertainties) in red.

The Magnetars

Name	P (s)	B (10^{14} G)	Age (kyr)	E (10^{33} erg s $^{-1}$)	D (kpc)	L (10^{33} erg s $^{-1}$)	Band
CXOU J010043.1-721134	8.02	3.9	6.8	1.4	62.4	65	...
4U 0142+61	8.69	1.3	68	0.12	3.6	105	OIR/H
SGR 0418+5729	9.08	0.06	36000	0.00021	2	0.00096	...
SGR 0501+4516	5.76	1.9	15	1.2	2	0.81	OIR/H
SGR 0526-66	8.05	5.6	3.4	2.9	53.6	189	...
1E 1048.1-5937	6.46	3.9	4.5	3.3	9.0	49	OIR
(PSR J1119-6127)	0.41	4.1	1.6	2300	8.4	0.2	R/H
1E 1547.0-5408	2.07	3.2	0.69	210	4.5	1.3	O?/R/H
PSR J1622-4950	4.33	2.7	4.0	8.3	9	0.4	R
SGR 1627-41	2.59	2.2	2.2	43	11	3.6	...
CXOU J164710.2-455216	10.6	<0.66	>420	<0.013	3.9	0.45	...
1RXS J170849.0-400910	11.01	4.7	9.0	0.58	3.8	42	O?/H
CXOU J171405.7-381031	3.82	5.0	0.95	45	13	56	...
SGR J1745-2900	3.76	2.3	4.3	10	8.3	<0.11	R/H
SGR 1806-20	7.55	20	0.24	45	8.7	163	OIR/H
XTE J1810-197	5.54	2.1	11	1.8	3.5	0.043	OIR/R
Swift J1822.3-1606	8.44	0.14	6300	0.0014	1.6	>0.0004	...
SGR 1833-0832	7.56	1.6	34	0.32
Swift J1834.9-0846	2.48	1.4	4.9	21	4.2	<0.0084	...
1E 1841-045	11.79	7.0	4.6	0.99	8.5	184	...
(PSR J1846-0258)	0.327	0.49	0.73	8100	6.0	19	...
3XMM J185246.6+003317	11.56	<0.41	>1300	<0.0036	7	< 0.006	...
SGR 1900+14	5.20	7.0	0.9	26	12.5	90	H
SGR 1935+2154	3.24	2.2	3.6	17
1E 2259+586	6.98	0.59	230	0.056	3.2	17	OIR/H
SGR 0755-2933
SGR 1801-23
SGR 1808-20
AX J1818.8-1559
AX J1845.0-0258	6.97	2.9	...
SGR 2013+34

Figure: So far...

Supernova Remnant Associations

Basics:

Consequences:




Figure: Artist's depiction of supernova remnant and magnetar [Cite here]

Supernova Remnant Associations

Basics:

- 8/23 magnetars

Consequences:




Figure: Artist's depiction of supernova remnant and magnetar [Cite here]

Supernova Remnant Associations

Basics:

- 8/23 magnetars
- 2 other possible associations

Consequences:




Figure: Artist's depiction of supernova remnant and magnetar [Cite here]

Supernova Remnant Associations

Basics:

- 8/23 magnetars
- 2 other possible associations
- provides additional evidence to youth requirement

Consequences:




Figure: Artist's depiction of supernova remnant and magnetar [Cite here]

Supernova Remnant Associations

Basics:

- 8/23 magnetars
- 2 other possible associations
- provides additional evidence to youth requirement

Consequences:

- unexpected characteristics




Figure: Artist's depiction of supernova remnant and magnetar [Cite here]

Supernova Remnant Associations

Basics:

- 8/23 magnetars
- 2 other possible associations
- provides additional evidence to youth requirement

Consequences:

- unexpected characteristics
- challenges dynamo model and fossil field theory




Figure: Artist's depiction of supernova remnant and magnetar [Cite here]

Supernova Remnant Associations

Basics:

- 8/23 magnetars
- 2 other possible associations
- provides additional evidence to youth requirement

Consequences:

- unexpected characteristics
- challenges dynamo model and fossil field theory
- no conclusive theory of formation as of yet




Figure: Artist's depiction of supernova remnant and magnetar [Cite here]

Formation

Theories

Common Conditions

Formation

Theories

- Neutron stars with ~ 1 ms period at birth [1]

Common Conditions

Formation

Theories

- Neutron stars with ~ 1 ms period at birth [1]

Common Conditions

- parasitic binary star systems ending in supernova

Formation

Theories

- Neutron stars with ~ 1 ms period at birth [1]
- Leading Theory: divine entity

Common Conditions

- parasitic binary star systems ending in supernova

Formation

Theories

- Neutron stars with ~ 1 ms period at birth [1]
- Leading Theory: divine entity

Common Conditions

- parasitic binary star systems ending in supernova
-

Kinds of Magnetars

Bi-modal

Cross-species

Kinds of Magnetars

Bi-modal

- Persistent Magnetars

Cross-species

Kinds of Magnetars

Bi-modal

- Persistent Magnetars
- Transient Magnetars

Cross-species

Kinds of Magnetars

Bi-modal

- Persistent Magnetars
- Transient Magnetars

Cross-species

- Magnetar Pulsar

Kinds of Magnetars

Bi-modal

- Persistent Magnetars
- Transient Magnetars

Cross-species

- Magnetar Pulsar
- Theoretically: neutron star switch between magnetar, pulsar, and magnetar pulsar

Magnetic Breaking

Materials

Magnetic Breaking

Materials



EM-Bursts

EM-Bursts



Summary and Conclusions

What we know

What we don't know

Summary and Conclusions

What we know



What we don't know

Summary and Conclusions

What we know



What we don't know

- A definitive model for the formation of magnetars

References

- ① Duncan RC, Thompson C. 1992. ApJ 392:L9-L13