

Assignment 1

Hruday Beeravelli

Find Python Codes from below link

[https://github.com/Hruday-Beeravelli/
INTERNSHIP-IITH-1/blob/main/
Assignment1/Assignment1.py](https://github.com/Hruday-Beeravelli/INTERNSHIP-IITH-1/blob/main/Assignment1/Assignment1.py)

and latex-tikz codes from

[https://github.com/Hruday-Beeravelli/
INTERNSHIP-IITH-1/blob/main/
Assignment1/Assignment1.tex](https://github.com/Hruday-Beeravelli/INTERNSHIP-IITH-1/blob/main/Assignment1/Assignment1.tex)

$$= a \sqrt{(m_1^2 - m_2^2)^2 + 4(m_1 - m_2)^2} \quad (1.2.7)$$

Distance between $(am_1^2, 2am_1)$ and $(am_2^2, 2am_2)$ is

$$= a \sqrt{(m_1^2 - m_2^2)^2 + 4(m_1 - m_2)^2} \quad (1.2.8)$$

1 EXAMPLES 1

1.1 Question 7

Find the distance between the following pairs of points

$$\left(\frac{am_1^2}{2am_1}, \frac{am_2^2}{2am_2} \right) \quad (1.1.1)$$

1.2 Solution

The distance between two vectors is given by

$$\|\mathbf{A} - \mathbf{B}\| = \sqrt{(\mathbf{A} - \mathbf{B})^\top \cdot (\mathbf{A} - \mathbf{B})} \quad (1.2.1)$$

$$= \left\| \begin{pmatrix} am_1^2 \\ 2am_1 \end{pmatrix} - \begin{pmatrix} am_2^2 \\ 2am_2 \end{pmatrix} \right\| \quad (1.2.2)$$

$$= \left\| \begin{pmatrix} am_1^2 - am_2^2 \\ 2am_1 - 2am_2 \end{pmatrix} \right\| \quad (1.2.3)$$

$$= \sqrt{\begin{pmatrix} am_1^2 - am_2^2 \\ 2am_1 - 2am_2 \end{pmatrix}^\top \begin{pmatrix} am_1^2 - am_2^2 \\ 2am_1 - 2am_2 \end{pmatrix}} \quad (1.2.4)$$

$$= \sqrt{(am_1^2 - am_2^2) \quad 2am_1 - 2am_2} \begin{pmatrix} am_1^2 - am_2^2 \\ 2am_1 - 2am_2 \end{pmatrix} \quad (1.2.5)$$

$$= \sqrt{(am_1^2 - am_2^2)^2 + (2am_1 - 2am_2)^2} \quad (1.2.6)$$