

Problem Set 4

Tate Mason

Question 1 - Hansen 7.17

Part A

Part B

Part C

Question 2 - Hansen 7.28

```
library(haven)
library(dplyr)
```

Attaching package: 'dplyr'

The following objects are masked from 'package:stats':

`filter`, `lag`

The following objects are masked from 'package:base':

`intersect`, `setdiff`, `setequal`, `union`

```

rm(list = ls())

dat <- read_dta('~/.SchoolWork/Sem2/Metrics/PSets/PS3/cps09mar.dta')
sample <- (dat[,11]==1)&(dat[,2]==0)&(dat[,3]==1)
df <- dat[sample,]

y <- as.matrix(log(df[,5]/(df[,6]*df[,7])))
exp <- df[,1]-df[,4]-6
exp2 <- (exp^2)/100

x_df <- data.frame(
  education = df[,4],
  experience = exp,
  exp_squared = exp2,
  intercept = 1
)
x <- as.matrix(x_df)

xx <- t(x)%*%x
xy <- t(x)%*%y
beta <- solve(xx,xy)

fitted <- x %*% beta
resid <- y - fitted

n <- nrow(x)
k <- ncol(x)
xx_inv <- solve(t(x)%*%x)

df <- n-k

hc0 <- matrix(0, nrow=k, ncol=k)
for (i in 1:n) {
  xi <- matrix(x[i,], nrow=k)
  hc0 <- hc0 + resid[i]^2 * (xi %*% t(xi))
}

```

Part B

```
theta_hat <- beta[1]/(beta[2]+beta[3]/5)
cat("\nPart B: Estimated Theta")
```

Part B: Estimated Theta

```
print(theta_hat)
```

```
[1] 3.468335
```

Part C

```
gradient_g <- c(
  1/(beta[2] + beta[3]/5),
  -beta[1]/(beta[2]+beta[3]/5)^2,
  -beta[1]/(5*(beta[2]+beta[3]/5)^2),
  0
)

theta_se <- sqrt(t(gradient_g) %*% hc0 %*% gradient_g)
print(theta_se)
```

```
      [,1]
[1,] 121935.8
```

Part D

Part E

EQ 1

EQ 2

```

b0 <- 0
b1 <- 1
n <- 100
sim <- function() {
  X1 <- rexp(n)
  e <- mixtools::rnormmix(n,lambda=c(0.5,0.5),mu=c(-1,2),sigma=c(1,1))
  Y <- b0 + b1*X1 + e
  x <- cbind(1,X1)
  xx <- t(x)%*%x
  xy <- t(x)%*%Y
  bhat <- solve(xx,xy)
  return(c(bhat[2]))
}

```

These results show that the average of $\hat{\beta}_1 \rightarrow \beta_1$ as n grows. The variance also approaches 0. This is consistent with what was derived in class, that as $n \rightarrow \infty$, we see the predicted approach the actual, and variance should be 0 as with sufficiently large n , there will be no variance in observations.

```

run_mc <- function(n_sims = 1000) {
  mc_res <- sapply(1:n_sims, function(s) {
    sim()
  })
  cat("Mean b1:", mean(mc_res), "\n")
  cat("Variance of b1:", var(mc_res), "\n")
}

```

```
run_mc()
```

```

Mean b1: 1.00064
Variance of b1: 0.03828634

```

```

n <- 2
run_mc()

```

```

Mean b1: 20.36982
Variance of b1: 280954.1

```

```
n <- 10  
run_mc()
```

Mean b1: 0.9765303
Variance of b1: 0.7344353

```
n <- 50  
run_mc()
```

Mean b1: 1.001743
Variance of b1: 0.07827652

```
n <- 500  
run_mc()
```

Mean b1: 0.9989494
Variance of b1: 0.006521762

As $n \rightarrow \infty$, the mean and variance get closer to the true values. This is a showcase of the WLLN.

EQ 3

Part A

```
b0 <- 0  
b1 <- 1  
num_sims <- 1000  
alpha <- 0.05  
  
sim_test <- function(n, b1_true, b1_null) {  
  X <- rexp(n)  
  e <- mixtools::rnormmix(n, lambda=c(0.5,0.5), mu=c(-2,2), sigma=c(1,1))  
  Y <- b0+b1*X + e  
  x <- cbind(1,X)  
  xx <- t(x)%*%x  
  xy <- t(x)%*%Y
```

```

bhat <- solve(xx,xy)
b0_hat <- bhat[1]
b1_hat <- bhat[2]

yhat <- x %*% bhat
ehat <- Y - yhat

sigma_sq_hat <- sum(ehat^2)/(n-2)
var_cov_matrix <- as.numeric(sigma_sq_hat)*solve(xx)
se_b1_hat <- sqrt(var_cov_matrix[2,2])
t_stat <- (b1_hat - b1_null)/se_b1_hat

df <- n-2
t_crit <- qt(1-alpha/2,df)
reject <- abs(t_stat) > t_crit
p_val <- 2*pt(abs(t_stat), df=df, lower.tail=FALSE)

return(list(
  b1_hat = b1_hat,
  se_b1_hat = se_b1_hat,
  t_stat = t_stat,
  t_crit = t_crit,
  p_val = p_val,
  reject = reject
))
}

```

```

run_hypothesis_test <- function(n, b1_true, b1_null) {
  results <- data.frame(
    b1_hat = numeric(num_sims),
    se_b1_hat = numeric(num_sims),
    t_stat = numeric(num_sims),
    p_val = numeric(num_sims),
    reject = logical(num_sims)
  )

  for (i in 1:num_sims) {
    sim_result <- sim_test(n,b1_true,b1_null)
    results$b1_hat[i] <- sim_result$b1_hat
    results$se_b1_hat[i] <- sim_result$se_b1_hat
    results$t_stat[i] <- sim_result$t_stat
    results$p_val[i] <- sim_result$p_val
  }
}

```

```

    results$reject[i] <- sim_result$reject
  }

  reject_rate <- mean(results$reject)
  mean_b1_hat <- mean(results$b1_hat)
  var_b1_hat <- var(results$b1_hat)
  mean_se_b1_hat <- mean(results$se_b1_hat)
  theoretical_var <- mean(results$se_b1_hat^2)

  return(list(
    results = results,
    reject_rate = reject_rate,
    mean_b1_hat = mean_b1_hat,
    var_b1_hat = var_b1_hat,
    mean_se_b1_hat = mean_se_b1_hat,
    theoretical_var = theoretical_var
  ))
}

```

```

results_100_true <- run_hypothesis_test(n=100,b1_true=1,b1_null=1)
cat("Part a & b: Results for n = 100, H:    = 1 (true value)\n")

```

Part a & b: Results for n = 100, H: = 1 (true value)

```

cat("Theoretical rejection rate at    = 0.05 should be: 0.05\n")

```

Theoretical rejection rate at = 0.05 should be: 0.05

```

cat("Observed rejection rate:", results_100_true$rejection_rate, "\n")

```

Observed rejection rate:

```

cat("Mean ^ :", results_100_true$mean_b1_hat, "\n")

```

Mean ^ : 1.00064

```

cat("Variance of ^ :", results_100_true$var_b1_hat, "\n")

```

Variance of ^ : 0.05304139

```
cat("Mean standard error of  $\hat{\beta}$  :", results_100_true$mean_se_b1_hat, "\n")
```

Mean standard error of $\hat{\beta}$: 0.2296013

```
cat("Theoretical variance (from SE):", results_100_true$theoretical_var, "\n\n")
```

Theoretical variance (from SE): 0.05385484

Part C

```
sample_size <- c(10,50,500,1000)
results_varying_n <- list()

for (n in sample_size) {
  results_varying_n[[paste0("n", n)]] <- run_hypothesis_test(n=n, b1_true=1, b1_null=1)
  cat("Results for n =", n, ", H:    = 1 (true value)\n")
  cat("Rejection rate:", results_varying_n[[paste0("n", n)]]$rejection_rate, "\n")
  cat("Mean  $\hat{\beta}$  :", results_varying_n[[paste0("n", n)]]$mean_b1_hat, "\n")
  cat("Variance of  $\hat{\beta}$  :", results_varying_n[[paste0("n", n)]]$var_b1_hat, "\n")
  cat("Mean standard error of  $\hat{\beta}$  :", results_varying_n[[paste0("n", n)]]$mean_se_b1_hat, "\n")
  cat("Theoretical variance (from SE):", results_varying_n[[paste0("n", n)]]$theoretical_var,
  }
}
```

Results for n = 10 , H: = 1 (true value)

Rejection rate:

Mean $\hat{\beta}$: 1.058527

Variance of $\hat{\beta}$: 1.004367

Mean standard error of $\hat{\beta}$: 0.932711

Theoretical variance (from SE):

Results for n = 50 , H: = 1 (true value)

Rejection rate:

Mean $\hat{\beta}$: 0.9846322

Variance of $\hat{\beta}$: 0.1098474

Mean standard error of $\hat{\beta}$: 0.3363417

Theoretical variance (from SE):

Results for n = 500 , H: = 1 (true value)

Rejection rate:
Mean $\hat{}$: 0.9953861
Variance of $\hat{}$: 0.01004739
Mean standard error of $\hat{}$: 0.1006388
Theoretical variance (from SE):

Results for n = 1000 , H: = 1 (true value)
Rejection rate:
Mean $\hat{}$: 0.9989068
Variance of $\hat{}$: 0.004919958
Mean standard error of $\hat{}$: 0.07101032
Theoretical variance (from SE):

Part D

```
results_100_false <- run_hypothesis_test(n=100,b1_true=1,b1_null=0)
cat("Part d: Results for n = 100, H: = 0 (false null)\n")
```

Part d: Results for n = 100, H: = 0 (false null)

```
cat("Rejection rate (power):", results_100_false$rejection_rate, "\n")
```

Rejection rate (power):

```
cat("Mean  $\hat{\phantom{x}}$  :", results_100_false$mean_b1_hat, "\n")
```

Mean $\hat{}$: 0.994658

```
cat("Variance of  $\hat{\phantom{x}}$  :", results_100_false$var_b1_hat, "\n\n")
```

Variance of $\hat{}$: 0.05434488

```
results_varying_n_false <- list()

for (n in sample_size) {
  set.seed(123)
  results_varying_n_false[[paste0("n", n)]] <- run_hypothesis_test(n = n, b1_true = 1, b1_null = 0)
```

```

cat("Results for n =", n, ", H:   = 0 (false null)\n")
cat("Rejection rate (power):", results_varying_n_false[[paste0("n", n)]]$rejection_rate, "
cat("Mean ^ :", results_varying_n_false[[paste0("n", n)]]$mean_b1_hat, "\n")
cat("Variance of ^ :", results_varying_n_false[[paste0("n", n)]]$var_b1_hat, "\n")
}

```

```

Results for n = 10 , H:   = 0 (false null)
Rejection rate (power):
Mean ^ : 1.017303
Variance of ^ : 1.029224
Results for n = 50 , H:   = 0 (false null)
Rejection rate (power):
Mean ^ : 1.009133
Variance of ^ : 0.1264878
Results for n = 500 , H:   = 0 (false null)
Rejection rate (power):
Mean ^ : 1.004012
Variance of ^ : 0.009162103
Results for n = 1000 , H:   = 0 (false null)
Rejection rate (power):
Mean ^ : 0.997987
Variance of ^ : 0.005327341

```