

Microscopic and Statistical Origin of the Emergent Pseudoscalar Vacuum Mode

Erick Sangalang

ORCID: 0009-0009-4039-2815

2026

Abstract

This paper develops a possible microscopic and statistical interpretation for the emergent pseudoscalar mode introduced in a phenomenological model of vacuum electrodynamics. Rather than treating the field as a fundamental axion-like particle, the present work interprets the mode as a coarse-grained collective variable associated with unresolved mesoscopic vacuum configurations. A statistical ensemble of microscopic vacuum states is introduced, from which an effective order parameter, free-energy functional, temperature-dependent mass scale, damping term, and electromagnetic coupling are motivated. The framework is intentionally exploratory: it does not claim a completed microscopic theory of the vacuum, but instead provides a plausible statistical route by which an emergent pseudoscalar response could arise and generate experimentally testable signatures such as vacuum birefringence, delayed electromagnetic emission, and resonant cavity response.

Keywords: emergent vacuum structure; pseudoscalar order parameter; statistical field theory; modified electrodynamics; vacuum birefringence; resonant cavities; effective field theory

1 Introduction

A previous phenomenological framework considered the possibility that the electromagnetic vacuum may possess an emergent mesoscopic degree of freedom capable of coupling dynamically to electromagnetic fields. In that model, the new mode was represented by an effective pseudoscalar order parameter $\phi(x, t)$ coupled to electromagnetism through a gauge-invariant interaction of the form

$$\mathcal{L}_{\text{int}} = g\phi F_{\mu\nu}\tilde{F}^{\mu\nu}. \quad (1)$$

The resulting phenomenology predicts temperature-dependent vacuum birefringence, delayed electromagnetic emission in resonant systems, and scaling relations for high- Q cavity experiments.

Throughout the paper, natural units with $\hbar = c = 1$ are assumed unless otherwise stated. In these units, the pseudoscalar field has mass dimension one, m_ϕ , ω , and Γ have dimensions of energy, ξ has dimensions of inverse energy, and the effective coupling g_{eff} has dimensions of inverse energy. The quantities m_ϕ , Γ , g_{eff} , and ξ are phenomenological parameters to be constrained experimentally rather than fixed fundamental constants.

The purpose of the present paper is to address a natural follow-up question: what microscopic or statistical vacuum structure could give rise to such an emergent mode?

This work does not propose a final microscopic theory of the vacuum. Instead, it constructs a conservative statistical interpretation in which the pseudoscalar field ϕ is treated as a coarse-grained collective coordinate describing an ensemble of unresolved microscopic or mesoscopic vacuum configurations. The approach is analogous in spirit to the emergence of order parameters in condensed matter systems, where macroscopic response fields may arise from many underlying microscopic degrees of freedom without requiring each collective mode to be fundamental.




Figure 1: Conceptual flow from unresolved microscopic configurations to an effective pseudoscalar vacuum response. The diagram emphasizes that ϕ is not assumed to be fundamental, but is treated as a coarse-grained collective coordinate whose dynamics can generate testable electromagnetic signatures.

2 Conceptual Motivation

2.1 Vacuum as an Effective Medium

In standard classical electrodynamics, the vacuum is described by fixed constants ϵ_0 and μ_0 . In quantum electrodynamics, the vacuum is not empty, but exhibits polarization effects due to quantum fluctuations. At sufficiently low energies, many physical systems can be described by effective field theories, where only long-wavelength collective variables are retained while microscopic details are integrated out.

The guiding hypothesis of the present work is that if the vacuum possesses unresolved microscopic structure, then its macroscopic response to strong electromagnetic fields may be expressible through an effective collective coordinate. This coordinate is not assumed to be a new fundamental particle. Instead, it is treated as a phenomenological order parameter encoding statistical bias among microscopic vacuum configurations.

2.2 Why a Pseudoscalar?

The electromagnetic invariant

$$F_{\mu\nu}\tilde{F}^{\mu\nu} = -4\mathbf{E} \cdot \mathbf{B} \quad (2)$$

is odd under parity and time reversal. Therefore, a field coupled linearly to this invariant must transform as a pseudoscalar if the total interaction is to preserve the desired symmetry structure. In the present model, ϕ is interpreted as a pseudoscalar order parameter measuring handedness, chirality, or orientation bias within the microscopic vacuum ensemble.

This interpretation distinguishes the model from a conventional scalar medium response. The relevant collective variable is not simply a density-like excitation, but a parity-sensitive measure of microscopic organization.

3 Microscopic Ensemble Picture

Let the microscopic vacuum be described by a large set of unresolved configurations $\{\sigma_i\}$, where i labels local microscopic degrees of freedom. These variables are not specified in detail. They may represent topological configurations, fluctuating polarization states, microscopic orientations, or other unknown vacuum substructure.

The partition function is written schematically as

$$Z = \sum_{\{\sigma_i\}} \exp[-\beta H(\{\sigma_i\})], \quad (3)$$

where

$$\beta = \frac{1}{k_B T_{\text{eff}}}. \quad (4)$$

Here T_{eff} is an effective temperature parameter describing statistical disorder in the microscopic ensemble. It need not correspond directly to ordinary thermodynamic temperature in all contexts, but in laboratory systems it may be influenced by environmental temperature, noise, boundary conditions, and cavity preparation.

3.1 Definition of the Order Parameter

A coarse-grained pseudoscalar order parameter may be defined as

$$\phi(x, t) = \langle \chi(\sigma_i) \rangle_{\Delta V}, \quad (5)$$

where $\chi(\sigma_i)$ is a microscopic pseudoscalar quantity associated with local handedness or orientation bias, and $\langle \cdots \rangle_{\Delta V}$ denotes averaging over a mesoscopic volume ΔV .

The order parameter vanishes when the microscopic ensemble is statistically symmetric:

$$\phi = 0. \quad (6)$$

A nonzero value corresponds to a biased ensemble in which one handedness or orientation sector is statistically favored:

$$\phi \neq 0. \quad (7)$$

4 Effective Free Energy

At long wavelengths, the statistical behavior of the order parameter can be described by an effective free-energy functional:

$$\mathcal{F}[\phi] = \int d^3x \left[\frac{1}{2} (\nabla \phi)^2 + \frac{1}{2} m_\phi^2(T_{\text{eff}}) \phi^2 + \frac{\lambda}{4} \phi^4 + \cdots \right]. \quad (8)$$

The quadratic term determines the energetic cost of exciting the collective mode. The quartic term stabilizes the theory for $\lambda > 0$.

4.1 Entropic Contribution

Assume the effective free energy may be decomposed as

$$F(\phi, T_{\text{eff}}) = U(\phi) - T_{\text{eff}} S(\phi). \quad (9)$$

If the entropy is expanded around the unbiased state,

$$S(\phi) = S_0 - \alpha \phi^2 + \mathcal{O}(\phi^4), \quad (10)$$

then

$$F(\phi, T_{\text{eff}}) = U(\phi) + \alpha T_{\text{eff}} \phi^2 + \mathcal{O}(\phi^4). \quad (11)$$

This motivates a temperature-dependent effective mass:

$$m_\phi^2(T_{\text{eff}}) = m_0^2 + c T_{\text{eff}}, \quad (12)$$

where m_0 and c are phenomenological constants. In the simplified limit $m_0^2 \ll c T_{\text{eff}}$, the model reduces to

$$m_\phi^2(T_{\text{eff}}) \propto T_{\text{eff}}. \quad (13)$$

This relation provides a possible statistical explanation for the temperature-dependent response proposed in the companion phenomenological model.

5 Electromagnetic Biasing of the Ensemble

The interaction between the emergent pseudoscalar order parameter and electromagnetism is written as

$$\mathcal{L}_{\text{int}} = g\phi F_{\mu\nu} \tilde{F}^{\mu\nu} = -4g\phi \mathbf{E} \cdot \mathbf{B}. \quad (14)$$

This term implies that regions with nonzero $\mathbf{E} \cdot \mathbf{B}$ bias the microscopic ensemble toward a nonzero pseudoscalar order parameter.

The effective potential in the presence of electromagnetic fields may be written as

$$V_{\text{eff}}(\phi) = \frac{1}{2} m_\phi^2 \phi^2 + \frac{\lambda}{4} \phi^4 - g_{\text{eff}} \phi \mathbf{E} \cdot \mathbf{B}. \quad (15)$$

Figure 2 illustrates the same statement geometrically: a nonzero electromagnetic invariant $\mathbf{E} \cdot \mathbf{B}$ tilts the effective potential and shifts the equilibrium value of the collective pseudoscalar coordinate.

In the linear-response regime, minimizing the effective potential gives

$$\phi \approx \frac{g_{\text{eff}} \mathbf{E} \cdot \mathbf{B}}{m_\phi^2(T_{\text{eff}})}. \quad (16)$$

If $m_\phi^2(T_{\text{eff}}) \propto T_{\text{eff}}$, then

$$\phi \propto \frac{g_{\text{eff}} \mathbf{E} \cdot \mathbf{B}}{T_{\text{eff}}}. \quad (17)$$

This reproduces the leading scaling assumed in the phenomenological treatment.




Figure 2: Schematic effective potential for the pseudoscalar order parameter. With no electromagnetic bias, the equilibrium lies at the statistically symmetric point. A nonzero $\mathbf{E} \cdot \mathbf{B}$ term tilts the potential and shifts the minimum, corresponding to an induced pseudoscalar response of the microscopic ensemble.

6 Dynamical Equation for the Emergent Mode

A statistical order parameter generally does not respond instantaneously. Relaxation, damping, and finite correlation times are expected. The effective equation of motion may therefore be written as

$$\square\phi + \Gamma\partial_t\phi + m_\phi^2\phi + \lambda\phi^3 = g_{\text{eff}}\mathbf{E} \cdot \mathbf{B}, \quad (18)$$

where Γ is a phenomenological damping coefficient.

The damping term represents irreversible relaxation of the mesoscopic ensemble toward equilibrium. Physically, Γ^{-1} defines the characteristic lifetime of the collective excitation.

6.1 Linear Susceptibility

For small oscillations about equilibrium, the nonlinear term may be neglected. In Fourier space, the response is

$$\phi(\omega) = \chi_\phi(\omega)g_{\text{eff}}[\mathbf{E} \cdot \mathbf{B}](\omega), \quad (19)$$

where

$$\chi_\phi(\omega) = \frac{1}{m_\phi^2 - \omega^2 - i\Gamma\omega}. \quad (20)$$

The magnitude of the linear response is therefore

$$|\chi_\phi(\omega)| = \frac{1}{\sqrt{(m_\phi^2 - \omega^2)^2 + \Gamma^2\omega^2}}. \quad (21)$$

This susceptibility produces three important physical features:

1. resonant enhancement when $\omega \approx m_\phi$,

2. phase delay between electromagnetic driving and vacuum response,
3. delayed re-radiation after the external drive is removed.




Figure 3: Normalized susceptibility of the emergent pseudoscalar mode. The response peaks near $\omega \approx m_\phi$, while larger damping Γ broadens and suppresses the resonance. This illustrates how finite relaxation of the microscopic ensemble would appear experimentally as a linewidth and reduced peak response.

7 Statistical Interpretation of Delayed Emission

In the companion phenomenological model, delayed electromagnetic emission was proposed as a possible observable signature. In the present statistical interpretation, delayed emission arises because the driven electromagnetic field temporarily biases the microscopic ensemble. When the drive is removed, the ensemble relaxes back toward equilibrium.

The relaxation process can transfer part of the stored collective excitation back into electromagnetic modes. In the simplest approximation, the decay of the order parameter after the drive is removed is

$$\phi(t) \sim \phi_0 e^{-\Gamma t/2} \cos(\omega_\phi t + \delta), \quad (22)$$

where ω_ϕ is the effective oscillation frequency and δ is a phase offset. The observable delayed signal is therefore expected to exhibit a characteristic decay time

$$\tau_\phi \sim \Gamma^{-1}. \quad (23)$$

This provides a concrete experimental handle: a real delayed-emission signal should not merely appear after the drive is removed, but should also exhibit a reproducible decay constant and scaling behavior with cavity parameters, field strength, and temperature.




Figure 4: Schematic delayed-emission ringdown of the emergent pseudoscalar coordinate after the external drive is removed. A genuine signal should exhibit a reproducible decay time and phase relation rather than an arbitrary transient.

8 Correlation Length and Mesoscopic Structure

A microscopic statistical model also implies a correlation length. Near equilibrium, the two-point correlation function may be written schematically as

$$\langle \phi(x)\phi(0) \rangle \sim \frac{e^{-|x|/\xi}}{|x|}, \quad (24)$$

where ξ is the correlation length of the emergent vacuum mode.

For a massive collective excitation,

$$\xi \sim \frac{1}{m_\phi}. \quad (25)$$

Thus, lower effective mass corresponds to longer-range coherent response. In resonant cavity experiments, observable enhancement would be expected only if the coherence scale is not too small compared with the relevant cavity mode structure.

This suggests an additional experimental consistency requirement:

$$\xi \lesssim L_{\text{cavity}} \quad (26)$$

for localized response, or

$$\xi \sim L_{\text{cavity}} \quad (27)$$

for strong coherent enhancement.

9 Relation to Axion and Axion-Like Models

The interaction $\phi F_{\mu\nu} \tilde{F}^{\mu\nu}$ resembles the coupling used in axion and axion-like particle models. However, the interpretation differs.

In conventional axion models, ϕ is typically treated as a fundamental or particle-like field with a microscopic origin tied to symmetry breaking. In the present framework, ϕ is treated as a collective




Figure 5: Schematic relation between the mesoscopic correlation length ξ and the resonant cavity scale. If ξ is much smaller than the mode structure, the response averages out locally; if ξ approaches the relevant cavity scale, coherent enhancement may become possible.

statistical variable. Its mass, damping rate, and coupling may therefore depend on environmental preparation, boundary conditions, and collective response rather than being fixed fundamental constants.

This distinction is important experimentally. If the mode is emergent rather than fundamental, then cavity geometry, thermal preparation, electromagnetic history, and boundary conditions may influence the observed response.

10 Compatibility Requirements

Any viable microscopic or statistical model must satisfy several constraints:

1. It must preserve gauge invariance.
2. It must preserve charge conservation.
3. It must remain compatible with precision QED tests.
4. It must avoid allowing uncontrolled free-energy extraction.
5. It must explain why the response is weak or absent in ordinary conditions.
6. It must produce falsifiable laboratory signatures.

The present framework satisfies the first two requirements at the effective-field-theory level because the coupling is built from gauge-invariant electromagnetic tensors. The remaining requirements impose constraints on the allowed values of g_{eff} , m_ϕ , Γ , and ξ .

10.1 Lorentz-Symmetry Considerations

The present model does not assume fundamental Lorentz-symmetry breaking. The effective pseudoscalar response is instead treated as a collective variable that becomes dynamically relevant only in the presence of external electromagnetic driving, boundary conditions, or nonequilibrium preparation. Any apparent preferred direction in a laboratory experiment would therefore be associated with the applied fields, cavity geometry, or measurement frame rather than with a permanent background vector or tensor built into empty space.

10.2 Why the Response May Be Weak in Ordinary Conditions

A central consistency requirement is that the proposed response remain weak or absent in ordinary electromagnetic environments. Within the statistical interpretation, suppression may occur through several mechanisms: weak effective coupling g_{eff} , rapid damping Γ , short correlation length ξ , thermal or environmental disorder encoded by T_{eff} , and insufficient coherent electromagnetic biasing through $\mathbf{E} \cdot \mathbf{B}$. Observable effects would therefore be expected only in carefully prepared strong-field, low-noise, resonant configurations rather than in generic laboratory conditions.

11 Possible Microscopic Mechanisms

Although the exact microscopic origin is left open, several broad classes of mechanisms could motivate an emergent pseudoscalar response:

11.1 Topological Configuration Bias

The vacuum ensemble may contain topologically distinct configurations. An applied $\mathbf{E} \cdot \mathbf{B}$ background could statistically favor one orientation sector over another, generating an effective pseudoscalar order parameter.

11.2 Chiral Mesoscopic Fluctuations

If microscopic vacuum fluctuations possess transient handedness, then strong electromagnetic fields may bias the population of left- and right-handed configurations. The order parameter ϕ would then measure the imbalance between those sectors.

11.3 Boundary-Induced Vacuum Organization

In resonant cavities, conducting boundaries restrict allowed electromagnetic modes. If the vacuum response depends on mode geometry, then boundary conditions may enhance or suppress the effective pseudoscalar response.

11.4 Collective Nonequilibrium Response

The emergent mode may be absent in equilibrium but become dynamically accessible under strong coherent driving. In this case, ϕ is best interpreted as a nonequilibrium collective response rather than a permanent vacuum condensate.

12 Experimental Consequences

The microscopic/statistical interpretation sharpens the experimental predictions of the phenomenological model. A genuine emergent collective mode should exhibit:

1. temperature-dependent response,
2. finite relaxation time,
3. resonant enhancement near $\omega \approx m_\phi$,
4. dependence on cavity boundary conditions,

5. reproducible phase lag relative to the electromagnetic drive,
6. scaling with $\mathbf{E} \cdot \mathbf{B}$ rather than with $\mathbf{E}$ or $\mathbf{B}$ alone.

Table 1: Representative experimental signatures implied by the microscopic/statistical interpretation.

Observable	Leading behavior	Experimental handle
Pseudoscalar response	$\phi \propto g_{\text{eff}} \mathbf{E} \cdot \mathbf{B} / T_{\text{eff}}$	field alignment, cryogenic preparation
Resonance enhancement	peak near $\omega \approx m_\phi$	frequency sweep, cavity tuning
Linewidth	set by Γ	ringdown and quality-factor analysis
Delayed emission	$\tau_\phi \sim \Gamma^{-1}$	time-domain detection after drive shutoff
Coherence effects	controlled by ξ / L_{cavity}	cavity geometry and boundary conditions
Pseudoscalar selection rule	scales with $\mathbf{E} \cdot \mathbf{B}$	field reversal and null-orientation tests

A null result across these signatures would constrain or falsify the proposed statistical origin.

13 Limitations

The present paper is intentionally exploratory. Several important limitations remain:

1. The microscopic variables σ_i are not explicitly identified.
2. The effective temperature T_{eff} requires sharper operational definition.
3. The coupling g_{eff} is not derived from first principles.
4. The damping coefficient Γ is phenomenological.
5. Existing experimental bounds must be incorporated quantitatively.
6. A full detector-level cavity simulation has not yet been performed.

These limitations define the next stage of the research program.

14 Conclusion

This work proposes a possible microscopic and statistical interpretation for the emergent pseudoscalar vacuum mode used in a phenomenological model of modified electrodynamics. The central idea is that ϕ need not be a fundamental particle, but may instead represent a coarse-grained collective coordinate associated with unresolved mesoscopic vacuum configurations.

By modeling the vacuum as a statistical ensemble, one can motivate an effective free-energy functional, a temperature-dependent mass scale, a finite damping time, a resonant susceptibility, and coupling to the parity-sensitive electromagnetic invariant $\mathbf{E} \cdot \mathbf{B}$. This provides a conceptual bridge between the phenomenological predictions of emergent vacuum electrodynamics and a possible underlying statistical mechanism.

The framework remains speculative, but it is structured to generate falsifiable predictions. Future work should focus on quantitative bounds, microscopic toy models, detector-level simulations, and comparison with precision electrodynamic experiments.

References

- [1] F. Wilczek, “Two Applications of Axion Electrodynamics,” *Physical Review Letters* **58**, 1799–1802 (1987).
- [2] P. Sikivie, “Experimental Tests of the Invisible Axion,” *Physical Review Letters* **51**, 1415–1417 (1983).
- [3] G. Raffelt and L. Stodolsky, “Mixing of the Photon with Low-Mass Particles,” *Physical Review D* **37**, 1237–1249 (1988).
- [4] S. Weinberg, “Phenomenological Lagrangians,” *Physica A* **96**, 327–340 (1979).
- [5] S. Weinberg, “Effective Field Theory, Past and Future,” arXiv:0908.1964 [hep-th] (2009).
- [6] J. Schwinger, “On Gauge Invariance and Vacuum Polarization,” *Physical Review* **82**, 664–679 (1951).
- [7] W. Heisenberg and H. Euler, “Folgerungen aus der Diracschen Theorie des Positrons,” *Zeitschrift für Physik* **98**, 714–732 (1936).
- [8] M. Born and L. Infeld, “Foundations of the New Field Theory,” *Proceedings of the Royal Society of London A* **144**, 425–451 (1934).
- [9] L. D. Landau and E. M. Lifshitz, *Statistical Physics, Part 1*, 3rd ed., Pergamon Press (1980).
- [10] P. M. Chaikin and T. C. Lubensky, *Principles of Condensed Matter Physics*, Cambridge University Press (1995).
- [11] N. Goldenfeld, *Lectures on Phase Transitions and the Renormalization Group*, CRC Press (1992).
- [12] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, 4th ed., Oxford University Press (2002).
- [13] M. Kardar, *Statistical Physics of Fields*, Cambridge University Press (2007).
- [14] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory*, Westview Press (1995).
- [15] C. P. Burgess, “An Introduction to Effective Field Theory,” *Annual Review of Nuclear and Particle Science* **57**, 329–362 (2007).
- [16] A. Caticha, “Entropic Dynamics, Time and Quantum Theory,” *Journal of Physics A: Mathematical and Theoretical* **44**, 225303 (2011).
- [17] A. Caticha, “Entropic Dynamics,” *Entropy* **17**, 6110–6128 (2015).
- [18] E. Verlinde, “On the Origin of Gravity and the Laws of Newton,” *Journal of High Energy Physics* **04**, 029 (2011).
- [19] X.-L. Qi, T. L. Hughes, and S.-C. Zhang, “Topological Field Theory of Time-Reversal Invariant Insulators,” *Physical Review B* **78**, 195424 (2008).
- [20] M. Z. Hasan and C. L. Kane, “Colloquium: Topological Insulators,” *Reviews of Modern Physics* **82**, 3045–3067 (2010).