

Instructions

Need Help?

Call your primary care provider at 608-271-9000 if you have:

- a fast, irregular, or racing heartbeat that does not settle down, or new palpitations
- chest pain, shortness of breath, fainting, or severe dizziness — call 911
- unusual bruising, blood in your stool, black tarry stools, or bleeding that will not stop while taking Eliquis
- swelling of the ankles, sudden weight gain of more than 3 pounds in a day, or light-headedness when standing after your blood pressure medicine change
- home blood pressure readings over 160/100 on two separate days

Your medications have changed

START taking

chlorthalidone 12.5 MG tablet

Take 1 tablet (12.5 mg total) by mouth daily in the morning.

CHANGE how you take

losartan (COZAAR) 100 MG tablet

Take 1 tablet (100 mg total) by mouth daily. Was: 50-mg-daily

START taking

sodium picosulfate-magnesium oxide-citric acid (CLENPIQ) 10-3.5-12 GM/160 ML oral solution

Bowel preparation for colonoscopy. Dispensed: 4 bottles (2 kits). Follow the split-dose prep timing on the attached bowel prep instruction sheet — do not drink both doses on the same evening.

STOP taking

hydrochlorothiazide 25 MG tablet — replaced by chlorthalidone above. Do not take both.

NO CHANGE

apixaban (ELIQUIS) 5 MG tablet — Take 1 tablet by mouth twice daily. Keep taking as prescribed until Dr. Reyes or the GI team tells you otherwise. See the colonoscopy hold instructions below.

Review your updated medication list below and bring it to every appointment.

Blood Pressure & Atrial Fibrillation

Blood pressure today	152 / 94 mmHg (repeat 148 / 90)
Heart rate / rhythm	78 bpm, irregularly irregular (atrial fibrillation)
Weight / Height	212 lb (96.2 kg) / 5 ft 3 in
BMI	37.6 kg/m ² (obesity, class II)
Home BP goal	Less than 130 / 80 mmHg

- Check your blood pressure at home 2 mornings and 2 evenings each week. Sit quietly for 5 minutes first, feet flat, arm supported at heart level.
- Write down the readings or enter them in MyChart. We will review them at your 4-week recheck.
- Because you have atrial fibrillation, an automatic cuff may read low or give an error. If it does, take three readings and record the average.
- Get up slowly from sitting or lying down for the first 2 weeks on the new medication.

Lifestyle Modification — discussed today

Activity

- Goal: 150 minutes of moderate activity per week — for example, a brisk 30-minute walk 5 days a week.
- Start with 10-minute walks and build up. Add light strength work 2 days a week.
- Stop and rest if you feel chest pressure, severe shortness of breath, or your heart races and will not slow down.

Diet

- DASH-style eating: vegetables, fruits, whole grains, beans, fish, and low-fat dairy.
- Keep sodium under 1,500 mg per day. Most sodium comes from canned soups, deli meat, frozen meals, sauces, and restaurant food — not the salt shaker.
- Limit alcohol to no more than 1 drink per day; alcohol can trigger atrial fibrillation episodes.
- Avoid large swings in vitamin K or grapefruit intake and check with the pharmacist before any new supplement while on Eliquis.

Weight

- A 5–10% weight loss (about 11–21 lb) can meaningfully lower blood pressure and reduce atrial fibrillation episodes.
- Referral placed to the Medical Weight Management clinic — they will call you to schedule.

Colonoscopy Preparation

Procedure: Screening colonoscopy (first screening) • Date: Saturday, 08/01/2026 • Arrival: 7:00 AM
Location: Green Oaks Endoscopy Center, 1725 Parkwood Court, Level 1

- Eliquis hold:** Take your last dose of Eliquis on the morning of Wednesday 07/29/2026. Skip all doses on 07/30 and 07/31. Restart Eliquis the evening after the procedure unless the GI physician tells you otherwise. Do not stop it any earlier than this.
- Five days before (07/27):** stop nuts, seeds, popcorn, corn, and raw vegetables.
- Day before (07/31):** clear liquids only all day — no red or purple liquids. First Clenpiq dose in the early evening as directed.
- Day of (08/01):** second Clenpiq dose about 5 hours before arrival. Nothing by mouth 2 hours before arrival.
- Drink extra clear fluid with each prep dose. Arrange a licensed driver — you cannot drive yourself home after sedation.
- Call the GI office at 608-271-0300 if you cannot finish the prep or if your stool is not clear yellow liquid by bedtime.

Health Maintenance

- Colorectal cancer screening — **ordered today**, colonoscopy 08/01/2026
- Mammogram — due, please schedule at Breast Imaging, 608-271-2400
- Influenza vaccine — due this fall
- Lipid panel, CMP, TSH — drawn today, results in MyChart within 3 business days
- Basic metabolic panel — repeat in 2 weeks to check kidney function and potassium after the chlorthalidone start

Your Next Steps

DESTINATION

Green Oaks Internal Medicine

2100 Parkwood Court, Suite 300

Verona, WI 53593

608-271-9000

PICK UP YOUR MEDICATIONS

Neighborhood Pharmacy — Verona

455 Main Street, Verona, WI 53593

608-845-1120

- chlorthalidone 12.5 mg (new)
- losartan 100 mg (new strength)
- CLENPIQ oral solution — 4 bottles

Pick up the Clenpiq at least 3 days before 08/01 so you have it in hand.

READ THESE ATTACHMENTS

- HIGH BLOOD PRESSURE: HOME MONITORING
- ATRIAL FIBRILLATION AND BLOOD THINNERS
- CLENPIQ SPLIT-DOSE BOWEL PREP INSTRUCTIONS
- COLONOSCOPY: WHAT TO EXPECT
- DASH EATING PLAN AND LOW-SODIUM SHOPPING GUIDE

YOUR FUTURE APPOINTMENTS

08/01/2026 7:00 AM

Screening colonoscopy

Green Oaks Endoscopy Center

608-271-0300

08/15/2026 10:30 AM

Blood pressure recheck (nurse visit)

Thomas Nguyen, RN

Green Oaks Internal Medicine

01/2027

Follow-up with Angela Reyes, MD

You have three future appointments. Please review your full appointment list in MyChart.

OFFICE VISIT — TODAY

Angela Reyes, MD

Internal Medicine

Green Oaks Internal Medicine

608-271-9000

MyChart

Send messages to your doctor, view your test results, review your blood pressure log, schedule appointments, and more.

Go to mychart.greenoakshealth.org

Sign up code: DELG-4482-XKQ7

Activation code expires 10/16/2026.