







Comparación de resultados k-means (2 clases) por condición inicial

Condición inicial	Centroides iniciales	Centroides finales	Iteraciones	Q (WCSS total)	Var. intra (pond.)	Var. entre
Separados bajos (10 y 40)	[10, 40]	[45.6, 71.4]	10	2504834.7	56.9281	138.2470
Separados medios (20 y 80)	[20, 80]	[45.6, 71.4]	7	2504834.7	56.9281	138.2470
Separados altos (60 y 100)	[60, 100]	[45.6, 71.4]	7	2504834.7	56.9281	138.2470
Cercanos bajos (40 y 45)	[40, 45]	[45.6, 71.4]	9	2504834.7	56.9281	138.2470
Cercanos medios (45 y 50)	[45, 50]	[45.6, 71.4]	8	2504834.7	56.9281	138.2470
Invertidos (80 y 20)	[80, 20]	[45.6, 71.4]	7	2504834.7	56.9281	138.2470




Condición inicial: Separados bajos (10 y 40)






Condición inicial: Separados medios (20 y 80)






Condición inicial: Separados altos (60 y 100)



Condición inicial: Cercanos bajos (40 y 45)



Condición inicial: Cercanos medios (45 y 50)



Condición inicial: Invertidos (80 y 20)

