

Name:.....

Class:.....



Mathematics for Engineering

Exercise Book

Trần Trọng Huỳnh

CALCULUS

Chapter 1: Function and Limit

1. Find the domain of each function:

a. $f(x) = \sqrt{x+2}$

b. $f(x) = \frac{1}{x^2 - x}$

c. $f(x) = \ln(x+1) - \frac{x}{\sqrt{x-1}}$

2. Find the range of each function:

a. $f(x) = \sqrt{x-1}$

b. $f(x) = x^2 - 2x$

c. $f(x) = \sin x$

3. Determine whether is even, odd, or neither

a. $f(x) = \frac{x}{x^2 + 1}$

b. $f(x) = \frac{x^2}{x^4 + 1}$

c. $f(x) = \frac{x}{x+1}$

4. Let $f(x) = \sqrt{x}$ and $g(x) = \sqrt{2-x}$. Find each function

a. $f \circ g$

b. $g \circ f$

c. $g \circ g$

d. $f \circ f$

5. Explain how the following graphs are obtained from the graph of $f(x)$

a. $f(x-4)$

b. $f(-x)+3$

c. $f(x-2)-3$

d. $-f(x+5)$

6. Use the table to evaluate each expression

a. $f(g(1))$

b. $g(f(1))$

c. $f(f(1))$


d. $g(g(1))$

e. $g \circ f(3)$

f. $g \circ f(6)$

x	1	2	3	4	5	6
$f(x)$	3	1	4	2	2	5
$g(x)$	6	3	2	1	2	3

$$\begin{array}{llll} \text{a. } \lim_{x \rightarrow 3} \frac{x^2 + x - 12}{x - 3} & \text{b. } \lim_{x \rightarrow 1} \frac{x^6 - 1}{x^{10} - 1} & \text{c. } \lim_{x \rightarrow 0} \frac{\tan 3x}{\tan 5x} & \text{d. } \lim_{h \rightarrow 0} \frac{(3+h)^2 - 9}{h} \\ \text{e. } \lim_{t \rightarrow 0} \frac{\sqrt{t^2 + 9} - 3}{t^2} & \text{f. } \lim_{x \rightarrow +\infty} \frac{x^2 + x - 12}{x^3 - 3} & \text{g. } \lim_{x \rightarrow 0^-} \left(\frac{1}{x} - \frac{1}{|x|} \right) & \text{h. } \lim_{x \rightarrow 1} \frac{x^2 - 1}{|x - 1|} \end{array}$$

(a) 

$$\begin{array}{lll} \text{i. } \lim_{x \rightarrow 2^+} f(x) & \text{ii. } \lim_{x \rightarrow -3^+} f(x) & \text{iii. } \lim_{x \rightarrow -3} f(x) \\ \text{iv. } \lim_{x \rightarrow 4} f(x) & \text{v. } \lim_{x \rightarrow 0} f(x) & \text{vi. } \lim_{x \rightarrow -\infty} f(x) \end{array}$$

c. State the equations of the vertical asymptotes.

d. At what numbers is discontinuous?

10. Find the horizontal and vertical asymptotes of each curve

a. $f(x) = \frac{2x^2 + x - 1}{x^2 + x - 2}$ b. $f(x) = \frac{x - 9}{\sqrt{4x^2 + 3x + 2}}$ c. $f(x) = \frac{x^3 - 1}{x^3 - 3x + 2}$

11. Find the constant m that makes f continuous on \mathbb{R}

a. $f(x) = \begin{cases} x^2 - m^2, & x < 4 \\ mx + 20, & x \geq 4 \end{cases}$ b. $f(x) = \begin{cases} \frac{e^{2x} - 1}{x}, & x \neq 0 \\ m, & x = 0 \end{cases}$

c. $f(x) = \begin{cases} mx^2 + 2x, & x < 2 \\ x^3 - mx, & x \geq 2 \end{cases}$ d. $f(x) = \begin{cases} \frac{x^2 - 1}{\sqrt{x} - 1}, & x \neq 1 \\ mx + 1, & x = 1 \end{cases}$


12. Find the numbers at which the function $f(x) = \begin{cases} x + 2, & x < 0 \\ 2x^2, & 1 \geq x \geq 0 \\ 2 - x, & x > 1 \end{cases}$ is discontinuous.

Chapter 2: Derivatives

1. Use the given graph to estimate the value of each derivative

a. $f'(-3)$ b. $f'(-1)$

c. $f'(0)$ d. $f'(3)$



2. Find an equation of the tangent line to the curve at the given point:

a. $y = \frac{x-1}{x-2}$, $(3, 2)$

b. $y = \frac{2x}{x^2+1}$, $(0, 0)$

c. $y = 3 - 2x + x^2$, $x = 1$

d. $y = \frac{3-2x}{x-1}$, $y = -1$

3. Find y'

a. $y = x^2 - x\sqrt{x} + \frac{1}{x} + 2$

b. $y = \sqrt{x + \sqrt{x}}$

c. $y = \frac{x^2}{x+1}$

d. $y = x\sqrt{x+2}$

e. $y = \ln(x^2 + 1) - \frac{1}{x}$

f. $y = e^x \sin(2x+1)$

4. Find y'''

a. $y = xe^{3x-1}$

b. $y = \sqrt[3]{2x+1}$

c. $y = e^{-x} \cos x$


5. Find the values of a and b that make the following function differentiable for all x-values:

a. $f(x) = \begin{cases} ax+b, & x < 0 \\ x^2 - 3x, & x \geq 0 \end{cases}$ b. $f(x) = \begin{cases} \frac{e^x - 1}{x}, & x > 0 \\ ax+b, & x \leq 0 \end{cases}$


6. For $f(x) = \begin{cases} \frac{\sin x - x}{x^2}, & x \neq 0 \\ m, & x = 0 \end{cases}$, where m is constant. Suppose f is differentiable at zero, find $f'(0)$.

7. The graph of is given. State the numbers at which is not differentiable

a.



b.



8. A table of values for f, f', g and g' is given

x	$f(x)$	$g(x)$	$f'(x)$	$g'(x)$
1	3	2	4	6
2	1	8	5	7
3	7	2	7	9

a. If $h(x) = f(g(x))$, find $h'(1)$

b. If $H(x) = g \circ f(x)$, find $H'(1)$

c. If $F(x) = f \circ f(x)$, find $F'(2)$ d. If $G(x) = g \circ g(x)$, find $G'(3)$

9. If $h(x) = \sqrt{4 + 3f(x)}$, where $f(1) = 7, f'(1) = 4$, find $h'(1)$.

10. For the circle $x^2 + y^2 = 25$.

a. Find dy/dx

b. Find an equation of the tangent to the circle at the point (3, 4).

11. Let $(L): x^3 + y^3 = 6xy$

a. Find dy/dx

b. Find an equation of tangent to the curve (L) at the point (3, 3)

12. Find y'' by implicit differentiation

a. $x^4 + y^4 = 16$ b. $\sqrt{x} + \sqrt{y} = 4$ c. $x^3 + y^3 = 1$

13. A water tank has the shape of an inverted circular cone with base radius 2 m and height 4 m. If water is being pumped into the tank at a rate of $2 \text{ m}^3/\text{min}$, find the rate at which the water level is rising when the water is 3 m deep.

14. The length of a rectangle is increasing at a rate of 8 cm/s and its width is increasing at a rate of 3 cm/s. When the length is 20 cm and the width is 10 cm, how fast is the area of the rectangle increasing?

15. Each side of a square is increasing at a rate of 6 cm/s. At what rate is the area of the square increasing when the area of the square is 16 cm^2 ?

16. If $x^2 + y^2 = 25$ and $dy/dt = 6$, find dx/dt when $y = 4$ and $x > 0$.

17. If $z^2 = x^2 + y^2$ ($z > 0$), $dx/dt = 2, dy/dt = 3$, find dz/dt when $x = 5, y = 12$

18. Find the linearization $L(x)$ of the function at a .

a. $f(x) = \frac{1}{\sqrt{2+x}}$, $a = 0$ b. $f(x) = \sqrt[3]{5-x}$, $a = -3$

19. The radius of a circular disk is given as 24 cm with a maximum error in measurement of 0.2 cm. Use differentials to estimate the maximum error in the calculated area of the disk.

Chapter 3: Applications of Differentiation

1. Find the absolute maximum and absolute minimum values of on the given interval

- a. $f(x) = 3x^2 - 12x + 5$, $[0; 3]$ b. $f(x) = x^3 - 3x + 5$, $[0; 3]$
c. $f(x) = x\sqrt{4 - x^2}$, $[-1; 2]$ d. $f(x) = x - \ln x$, $\left[\frac{1}{2}; 2\right]$

2. Find the critical numbers of the function

- a. $f(x) = 5x^2 + 4x$ b. $f(x) = \frac{x-1}{x^2 - x + 1}$ c. $f(x) = x \ln x$

3. Find all numbers that satisfy the conclusion of the Rolle's Theorem

- a. $f(x) = x\sqrt{x+2}$, $[-2; 0]$ b. $f(x) = (x-2)\sqrt[3]{x^2}$, $[0; 2]$

4. Find all numbers that satisfy the conclusion of the Mean Value Theorem

- a. $f(x) = 3x^2 + 2x + 5$, $[-1; 1]$ b. $f(x) = e^{-2x}$, $[0; 3]$

5. If $f(1) = 10$ and $f'(x) \geq 2, \forall x \in [1; 4]$, how small can $f(4)$ possibly be?

6. Find where the function $f(x) = 3x^4 - 4x^3 - 12x^2 + 1$ is increasing and where it is decreasing.

7. Find the inflection points for the function

- a. $f(x) = x^4 - 4x + 1$ b. $f(x) = x^6$ c. $f(x) = xe^x$

8. Find the point on the parabola $y^2 = 2x$ that is closest to the point $(1; 4)$

9. Find the area of the largest rectangle that can be inscribed in a semicircle of radius r .

10. Find two numbers whose difference is 100 and whose product is a minimum.

11. Find two positive numbers whose product is 100 and whose sum is a minimum.

12. Use Newton's method with the specified initial approximation x_1 to find x_3

a. $x^3 + 2x - 4 = 0, \quad x_1 = 1$


b. $x^5 + 2 = 0, \quad x_1 = -1$

c. $\ln(x^2 + 1) - 2x - 1 = 0, \quad x_1 = 1$


d. $\ln(4 - x^2) = x, \quad x_1 = 1$

13. The figure shows the graphs of f, f' and f'' . Identify each curve, and explain your choices

a.



b.



14. Find the most general anti-derivative of the function.

a. $f(x) = 6x^2 - 2x + 3$

b. $f(x) = \sqrt[6]{x} + \frac{1}{x^2}$

c. $f(x) = \frac{x^2 + x + 2}{x}$

d. $f(x) = 2x(x^2 + 1)$

15. Find the anti-derivative of that satisfies the given condition

a. $f(x) = 5x^4 - 2x^5, F(0) = 4$

b. $f(x) = 4 - \frac{2x}{x^2 + 1}, F(0) = 1$


16. A particle is moving with the given data. Find the position of the particle

a. $v(t) = \sin t - \cos t, s(0) = 0$

b. $a(t) = 10\sin t + 3\cos t, s(0) = 0, s(2\pi) = 12$

c. $a(t) = 10 + 3t - 3t^2, s(0) = 0, s(2) = 10$

17. The figure shows the graph of the derivative f' of a function f



- On what intervals is f increasing or decreasing?
- For what values of x does f have a local maximum or minimum?

Chapter 4 - 6: Integration

1. Estimate the area under the graph of $y = f(x)$ using 6 rectangles and left endpoints

a. $f(x) = \frac{1}{x} + x$, $x \in [1, 4]$

b. $f(x) = x^2 + x - 2$, $x \in [-1, 2]$

c. A table of values for f is given

x	1	2	3	4	5	6	7
$f(x)$	5	6	3	2	7	1	2

3. Repeat part (1) using right endpoints

4. For the function $f(x) = e^{-x^2}$, $x \in [-2, 2]$. Estimate the area under the graph of using four approximating rectangles and taking the sample points to be (i) right endpoints, (ii) left endpoints and (iii) midpoints.

5. Use (a) the Trapezoidal Rule, (b) the Midpoint Rule, and (c) Simpson's Rule to approximate the given integral with the specified value of n .

a. $\int_0^3 \sqrt{x} dx$, $n = 4$

b. $\int_1^3 \frac{\sin x}{x} dx$, $n = 6$

6. Let $I = \int_0^3 \frac{dx}{x^2 + 1}$. Find the approximations L_6 , R_6 , M_6 , T_6 and S_6 for I .

7. Find the derivative of the function $g(x) = \int_0^x \sqrt{t^2 + 1} dt$

8. Find g'

a. $g(x) = \int_1^{x^4} \frac{1}{\cos t} dt$

b. $g(x) = \int_1^{\sqrt{x}} \frac{\sin u}{u} du$

$$c. \ g(x) = \int_{2x}^{x^2+x+2} \frac{e^t}{t} dt$$

$$c. \ g(x) = \int_{\sin x}^{\cos x} (1+v^2)^{10} dv$$

9. Find the average value of the function on the given interval

$$a. \ f(x) = x^2, \quad [-1,1]$$

$$b. \ f(x) = \frac{1}{x}, \quad [1,5]$$

$$c. \ f(x) = x\sqrt{x}, \quad [1,4]$$

$$d. \ f(x) = x \ln x, \quad [1, e^2]$$

10. A particle moves along a line so that its velocity at time t is: $v(t) = t^2 - t - 6$ (m/s)

a. Find the displacement of the particle during the time period $1 \leq t \leq 4$

b. Find the distance traveled during this time period

11. Suppose the acceleration function and initial velocity are $a(t) = t + 3$ (m/s²), $v(0) = 5$ (m/s). Find the velocity at time t and the distance traveled when $0 \leq t \leq 5$.

12. A particle moves along a line with velocity function $v(t) = t^2 - t$, where t is measured in meters per second. Find the displacement and the distance traveled by the particle during the time interval $t \in [0, 2]$.

13. Evaluate the integral

$$a. \int_0^2 x^2 \cdot \sqrt{x^3 + 1} dx$$

$$b. \int x e^{x^2} dx$$

$$c. \int \left(\frac{1}{x} + \sqrt{x} - 3x^2 \right) dx$$

$$d. \int_0^1 y(1+y^2)^5 dy$$

$$e. \int \frac{\ln x}{x} dx$$

$$f. \int \frac{t}{t^2 + 1} dt$$

14. Evaluate the integral

$$a. \int x e^x dx$$

$$b. \int_0^1 x^2 e^{-x} dx$$

$$c. \int x \sin x dx$$

$$d. \int \ln x dx$$

$$e. \int_1^e x \ln x dx$$

$$f. \int e^{\sqrt{x}} dx$$

15. Suppose $f(x)$ is differentiable, $f(1) = 4$ and $\int_0^1 f(x) dx = 5$. Find $\int_0^1 xf'(x) dx$

16. Suppose $f(x)$ is differentiable, $f(1) = 3$, $f(3) = 1$ and $\int_1^3 xf'(x) dx = 13$. What is the average value of f on the interval $[1, 3]$?

17. Let $f(x) = \begin{cases} -x-1, & -3 \leq x \leq 0 \\ -\sqrt{1-x^2}, & 0 < x \leq 1 \end{cases}$. Evaluate $\int_{-3}^1 f(x) dx$

18. Find a function f and a value of the constant m such that $2 \int_m^x f(t) dt = 2 \sin x - 1$

19. Suppose that $f(1) = 2$, $f(4) = 7$, $f'(1) = 5$, $f'(4) = 3$ and f'' is continuous. Find the value of $\int_1^4 xf''(x) dx$.

20. Determine whether each integral is convergent or divergent. Evaluate those that are convergent.

a. $\int_1^{\infty} \frac{dx}{(3x+1)^2}$

b. $\int_{-\infty}^0 \frac{dx}{2x-5}$

c. $\int_{-\infty}^{-1} \frac{dx}{\sqrt{2-x}}$

d. $\int_0^{\infty} \frac{xdx}{(x^2+2)^2}$

e. $\int_4^{\infty} e^{-\frac{y}{2}} dy$

f. $\int_{-\infty}^{-1} e^{-2t} dt$

g. $\int_{2\pi}^{\infty} \sin \phi d\phi$

h. $\int_{-\infty}^{\infty} xe^{-x^2} dx$

i. $\int_0^1 \frac{dx}{4x-1}$

j. $\int_0^1 \frac{dx}{\sqrt{1-x^2}}$

k. $\int_{-1}^1 \frac{e^x}{e^x-1} dx$

l. $\int_0^1 \frac{\ln x}{\sqrt{x}} dx$

21. Use the Comparison Theorem to determine whether the integral is convergent or divergent

a. $\int_1^{\infty} \frac{\cos^2 x dx}{1+x^2}$

b. $\int_1^{\infty} \frac{2+e^{-x}}{x} dx$

c. $\int_1^{\infty} \frac{dx}{x+e^{2x}}$

d. $\int_1^{\infty} \frac{x dx}{\sqrt{1+x^6}}$

e. $\int_0^{\frac{\pi}{2}} \frac{dx}{x \sin x}$

f. $\int_0^1 \frac{e^{-x} dx}{\sqrt{x}}$

Chapter 8: Series

1. Determine whether the sequence converges or diverges. If it converges, find the limit

a. $a_n = \frac{3+2n^2}{n+n^2}$ b. $a_n = \frac{\sqrt{n}}{\sqrt{2n+1}+3}$ c. $a_n = \frac{n}{\sqrt{n}+1}$ d. $a_n = \left(1 + \frac{2}{n}\right)^n$

e. $\left\{\sqrt{2}, \sqrt{2+\sqrt{2}}, \sqrt{2+\sqrt{2+\sqrt{2}}}, \dots\right\}$ f. $\left\{\sqrt{2}, \sqrt{2\sqrt{2}}, \sqrt{2\sqrt{2\sqrt{2}}}, \dots\right\}$

g. $\{0.12, 0.1212, 0.121212, \dots\}$

2. Find the limit of the sequence $\{a_n\}$

a. $a_1 = \sqrt{5}, a_{n+1} = \sqrt{5+a_n}$ b. $a_1 = 2, a_{n+1} = \frac{1}{3-a_n}$ c. $a_1 = 1, a_{n+1} = \frac{1}{1+a_n}$

3. Determine whether the series is convergent or divergent. If it is convergent, find its sum

a. $\sum_{n=1}^{\infty} \frac{1}{n(n+1)}$ b. $\sum_{n=2}^{\infty} \frac{n^2+n-1}{n(n-1)}$ c. $\sum_{n=2}^{\infty} \frac{1}{3 \cdot 2^{n-1}}$
d. $\sum_{n=1}^{\infty} \sin n$ e. $\sum_{n=1}^{\infty} \frac{1+3^n}{2^n}$ f. $\sum_{n=1}^{\infty} (0.8^n + 0.3^{n-1})$

4. Determine whether the series is convergent or divergent

a. $\sum_{n=1}^{\infty} \frac{1}{n^2+1}$ b. $\sum_{n=1}^{\infty} \frac{\sqrt{n}}{n+1}$ c. $\sum_{n=1}^{\infty} \left(\frac{1}{n^6} + \frac{4}{n\sqrt{n}}\right)$ d. $\sum_{n=1}^{\infty} ne^{-n}$
e. $\sum_{n=1}^{\infty} \frac{1}{2n+3}$ f. $\sum_{n=1}^{\infty} \frac{4+3^n}{2^n}$ g. $\sum_{n=1}^{\infty} \frac{n!}{n^2 2^n}$ h. $\sum_{n=1}^{\infty} \frac{\cos n}{n^2+1}$

5. Determine whether the series is convergent or divergent

$$\begin{array}{llll} \text{a. } \sum_{n=1}^{\infty} \frac{n}{2^n} & \text{b. } \sum_{n=1}^{\infty} \frac{(-3)^{n-1}}{\sqrt{n}} & \text{c. } \sum_{n=1}^{\infty} \frac{\sqrt{n+1}}{n^2+n+1} & \text{d. } \sum_{n=1}^{\infty} \frac{(-2)^n}{n!} \\ \text{e. } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2n+3} & \text{f. } \sum_{n=1}^{\infty} \frac{(-1)^{n-1} \ln n}{n} & \text{g. } \sum_{n=1}^{\infty} \frac{\cos \pi n}{\sqrt{n+1}} & \text{h. } \sum_{n=1}^{\infty} \left(\frac{n^2+1}{n^2+2n+3} \right)^n \end{array}$$

6. Determine whether the series is absolutely convergent, conditionally convergent, or divergent.

$$\begin{array}{llll} \text{a. } \sum_{n=1}^{\infty} \frac{(-3)^{n+1}}{n^3} & \text{b. } \sum_{n=1}^{\infty} \frac{n^2}{2^n} & \text{c. } \sum_{n=1}^{\infty} \frac{(-10)^n}{n!} & \text{d. } \sum_{n=1}^{\infty} \frac{(-1)^n n}{n^2+1} \\ \text{e. } \sum_{n=1}^{\infty} \frac{\sin 4n}{n^4} & \text{f. } \sum_{n=1}^{\infty} \frac{1}{n\sqrt{n+1}} & \text{g. } \sum_{n=2}^{\infty} \frac{\cos \pi n}{\ln n} & \text{h. } \sum_{n=1}^{\infty} \frac{(-1)^n n^n}{e^n (n!)^2} \end{array}$$

7. Find the radius of convergence and interval of convergence of the series

$$\begin{array}{llll} \text{a. } \sum_{n=0}^{\infty} \frac{(-3)^n x^n}{\sqrt{n+1}} & \text{b. } \sum_{n=0}^{\infty} \frac{n(x+2)^n}{3^{n+1}} & \text{c. } \sum_{n=0}^{\infty} \frac{x^n}{n!} & \text{d. } \sum_{n=0}^{\infty} \frac{(-1)^n (x-1)^n}{2n+1} \\ \text{e. } \sum_{n=0}^{\infty} n!(2x-1)^n & \text{f. } \sum_{n=0}^{\infty} \frac{x^n}{n^2 3^n} & \text{g. } \sum_{n=0}^{\infty} \sqrt{n+1} x^n & \text{h. } \sum_{n=0}^{\infty} \frac{(-2)^n (x-3)^n}{\sqrt[4]{n}} \end{array}$$

8. Find the first n terms in the Maclaurin series for the given function

$$\begin{array}{ll} \text{a. } f(x) = x \sin x, n = 4 & \text{b. } f(x) = x \cos 2x, n = 3 \\ \text{c. } f(x) = \ln(1+x^2), n = 4 & \text{d. } f(x) = e^x \sin x, n = 3 \end{array}$$

9. Approximate f by a Taylor polynomial with degree at the number a

$$\begin{array}{ll} \text{a. } f(x) = \sqrt{x+1}, n = 1, a = 0 & \text{b. } f(x) = \frac{1}{x}, n = 3, a = 1 \\ \text{c. } f(x) = e^{x^2}, n = 3, a = 0 & \text{d. } f(x) = \cos x, n = 4, a = \frac{\pi}{3} \end{array}$$

LINEAR ALGEBRA

Chapter 1: Systems of Linear Equations

1. Write the augmented matrix for each of the following systems of linear equations and then solve them.

$$\text{a. } \begin{cases} -x + y + 2z = 1 \\ 2x + 3y + z = -2 \\ 5x + 4y + 2z = 4 \end{cases}$$

$$\text{b. } \begin{cases} 2x + 3y + z = 10 \\ 2x - 3y - 3z = 22 \\ 4x - 2y + 3z = -2 \end{cases}$$

$$\text{c. } \begin{cases} x + y + z = 0 \\ 2x - y + 2z = 0 \\ x + z = 0 \end{cases}$$

$$\text{d. } \begin{cases} x_1 + 2x_2 - x_3 + x_4 = 0 \\ 2x_1 + 3x_2 - 2x_3 + 3x_4 = 0 \\ x_1 + x_2 - 3x_3 + x_4 = 0 \end{cases}$$

2. Compute the rank of each of the following matrices.

$$\text{a. } A = \begin{pmatrix} 1 & 1 & 2 \\ 3 & -1 & 1 \\ -1 & 3 & 4 \end{pmatrix}$$

$$\text{b. } B = \begin{pmatrix} -2 & 3 & 3 \\ 3 & -4 & 1 \\ -5 & 7 & 2 \end{pmatrix}$$

$$\text{c. } C = \begin{pmatrix} 1 & 1 & -1 & 4 \\ 2 & 1 & 3 & 0 \\ 0 & 1 & 5 & 8 \end{pmatrix}$$

$$\text{d. } D = \begin{pmatrix} 1 & 1 & -1 & 3 \\ -1 & 4 & 5 & -2 \\ 1 & 6 & 3 & 4 \end{pmatrix}$$

3. Find all values of k for which the system has nontrivial solutions and determine all solutions in each case.

$$\text{a. } \begin{cases} x - y + 2z = 0 \\ -x + y - z = 0 \\ x + ky + z = 0 \end{cases}$$

$$\text{b. } \begin{cases} x - 2y + z = 0 \\ x + ky - 3z = 0 \\ x - 6y + 5z = 0 \end{cases}$$

$$\text{c. } \begin{cases} x + y + z = 0 \\ x + y - z = 0 \\ x + y + kz = 0 \end{cases}$$

$$\text{d. } \begin{cases} x + y - z = 0 \\ ky - z = 0 \\ x + y + kz = 0 \end{cases}$$

4. Determine the values of m such that the system of linear equations has exactly one solution.

$$\text{a. } \begin{cases} x - y + 2z = m \\ -x + y - z = 0 \\ -x + my - z = 1 - m \end{cases}$$

$$\text{b. } \begin{cases} mx + y + z = 1 \\ x + my + z = m \\ x + y + mz = m^2 \end{cases}$$

$$\text{c. } \begin{cases} x + y - z = 1 \\ x + my + 2z = m \\ x + 2y + z = 2 \end{cases}$$

$$\text{d. } \begin{cases} x + my - mz = m \\ 2x + y - z = 2 \\ x + y + z = 0 \end{cases}$$

5. Determine the values of k such that the system of linear equations is inconsistent.

$$\text{a. } \begin{cases} x - y + 2z = m \\ -x + y - z = 0 \\ x - y + 3z = 1 - m \end{cases}$$

$$\text{b. } \begin{cases} x - 2y + 2z = m \\ x + my - z = 0 \\ 2x + y + mz = 2 - m \end{cases}$$

6. Find a , b and c so that the system
$$\begin{cases} x + ay + cz = 0 \\ bx + cy - 3z = 1 \\ ax + 2y + bz = 5 \end{cases}$$
 has the solution $(3, -1, 2)$

7. Consider the matrix $A = \begin{pmatrix} 2 & -1 & 3 \\ -4 & 2 & k \\ 4 & -2 & 6 \end{pmatrix}$

a. If A is the augmented matrix of a system of linear equations, determine the number of equations and the number of variables.

b. If A is the augmented matrix of a system of linear equations, find the value(s) of k such that the system is consistent.

c. If A is the coefficient matrix of a homogeneous system of linear equations, determine the number of equations and the number of variables.

d. If A is the coefficient matrix of a homogeneous system of linear equations, find the value(s) of k such that the system is consistent.

8. Find all values of k so that the system of equations has no solution.

$$\text{a. } \begin{cases} x + y - z = 2 \\ x + 2y + z = 3 \\ x + y + (k^2 - 5)z = k \end{cases}$$

$$\text{b. } \begin{cases} x + y - z = 1 \\ 2x + (k + 5)y - 2z = 4 \\ x + (k + 3)y + (k - 1)z = k + 3 \end{cases}$$

9. Find all values of a and b for which the system of equations
$$\begin{cases} x + y + 3z = 2 \\ x + 2y + 5z = 1 \\ 2x + 2y + az = b \end{cases}$$
 is inconsistent.

Chapter 2: Matrix Algebra

1. Let $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$, $B = \begin{pmatrix} -1 & 1 \\ 3 & 2 \end{pmatrix}$ and $C = \begin{pmatrix} 1 & 3 & -4 \\ -1 & 2 & 1 \end{pmatrix}$. Compute the matrix

- a. $2A - B^T$ b. AB c. BA d. AC
 e. CC^T f. $C^T C$ g. A^3 h. $B^2 A^T$

2. Suppose that A and B are nxn matrices. Simplify the expression

- a. $(A+B)^2 - (A-B)^2$ b. $A(BC - CD) + A(C - B)D - AB(C - D)$

3. Let $A = \begin{pmatrix} 3 & 1 & 2 \\ 4 & 8 & 0 \\ 0 & 1 & 2 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 5 & 2 & 1 \\ 1 & 8 & 0 & -6 \\ 1 & 4 & 3 & 7 \end{pmatrix}$.

- a. Compute AB
 b. Compute $f(A)$ if $f(x) = x^2 - 3x + 2$

4. Find the inverse of each of the following matrices.

- a. $\begin{pmatrix} 1 & 5 \\ 2 & -1 \end{pmatrix}$ b. $\begin{pmatrix} 2 & 1 \\ 2 & -4 \end{pmatrix}$ c. $\begin{pmatrix} 1 & -1 & 2 \\ -5 & 7 & -11 \\ -2 & 3 & -5 \end{pmatrix}$ d. $\begin{pmatrix} 1 & -1 & 3 \\ 2 & 0 & 5 \\ -1 & 1 & 0 \end{pmatrix}$

5. Given $A^{-1} = \begin{pmatrix} 1 & -1 & 3 \\ 2 & 0 & 5 \\ -1 & 1 & 0 \end{pmatrix}$. Find a matrix X such that

- a. $AX = \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix}$ b. $AX = \begin{pmatrix} 1 & -1 & 2 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix}$ c. $XA = \begin{pmatrix} 1 & 2 & -1 \\ 3 & 1 & 1 \end{pmatrix}$

6. Find A when

a. $(3A)^{-1} = \begin{pmatrix} 1 & 2 \\ 0 & -2 \end{pmatrix}$ b. $(I + 2A)^{-1} = \begin{pmatrix} 2 & 1 \\ 3 & 2 \end{pmatrix}$ c. $(A^{-1} - 2I)^T = -2 \begin{pmatrix} 1 & 4 \\ 3 & 11 \end{pmatrix}$

7. Write the system of linear equations in matrix form and then solve them.

a. $\begin{cases} 2x - y = 4 \\ 3x + 2y = -4 \end{cases}$ b. $\begin{cases} 2x + 3y + z = 10 \\ 2x - 3y - 3z = 22 \\ 4x - 2y + 3z = -2 \end{cases}$ c. $\begin{cases} x + y = a \\ 2x + 3y = 1 - 2a \end{cases} (a \in \mathbb{R})$

8. Find A^{-1} if

a. $A^2 - 6A + 5I = 0$ b. $A^2 + 3A - I = 0$ c. $A^4 = I$

9. Solve for X

a. $\begin{pmatrix} 1 & 2 \\ 2 & 3 \end{pmatrix} X = \begin{pmatrix} 1 & -1 \\ 3 & 3 \end{pmatrix}$ b. $ABXC = B^T$ c. $AX^T BC = B$

(where A , B and C are $n \times n$ invertible matrices)

10. Compute $\begin{pmatrix} -1 & 3 \\ 0 & 1 \end{pmatrix}^{101}$

11. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation, and assume that $T(1,2) = (-1,1)$ and $T(0,3) = (-3,3)$

a. Compute $T(11,-5)$ b. Compute $T(1,11)$

c. Find the matrix of T d. Compute $T^{-1}(2,3)$

12. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be reflection in $y = x$ followed by rotation through $\frac{\pi}{2}$. Find T

Chapter 3: Determinants and Diagonalization

1. Evaluate the determinant

$$\begin{array}{llll} \text{a. } \begin{vmatrix} 4u & -1 \\ -1 & 2v \end{vmatrix} & \text{b. } \begin{vmatrix} m-1 & 2 \\ m & 2 \end{vmatrix} & \text{c. } \begin{vmatrix} x-2 & -1 \\ -3 & x \end{vmatrix} & \text{d. } \begin{vmatrix} -2 & 0 & 0 \\ 4 & 6 & 0 \\ -3 & 7 & 2 \end{vmatrix} \\ \text{e. } \begin{vmatrix} -3 & 2 & 1 \\ 4 & 5 & 6 \\ 2 & -3 & 1 \end{vmatrix} & \text{f. } \begin{vmatrix} 2 & -1 & 1 \\ 0 & 2 & 1 \\ 1 & 3 & 4 \end{vmatrix} & \text{g. } \begin{vmatrix} x & y & 1 \\ -1 & -2 & 1 \\ 1 & 5 & 1 \end{vmatrix} & \text{h. } \begin{vmatrix} m & -1 & 0 \\ 1 & 2 & 1 \\ 2 & m & -3 \end{vmatrix} \end{array}$$

2. Find the minors and the cofactors of the matrix

$$\begin{array}{lll} \text{a. } A = \begin{pmatrix} 1 & 3 \\ -2 & 4 \end{pmatrix} & \text{b. } B = \begin{pmatrix} -3 & 4 & 2 \\ 6 & 3 & 1 \\ 4 & -7 & -8 \end{pmatrix} & \text{c. } C = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & 1 \\ 1 & 2 & m \end{pmatrix} \end{array}$$

$$3. \text{ Find the adjugate and the inverse of the matrix } A = \begin{pmatrix} 1 & 0 & 3 \\ 0 & -1 & 2 \\ 2 & 1 & 0 \end{pmatrix}$$

$$4. \text{ Let } A = \begin{pmatrix} 1 & * & * & * \\ 0 & -1 & * & * \\ 0 & 0 & 2 & * \\ 0 & 0 & 0 & 2 \end{pmatrix}. \text{ Find}$$

$$\begin{array}{lll} \text{a. } |2A^{-1}| & \text{b. } |AA^T| & \text{c. } |\text{adj } A| \\ \text{d. } |-A^3| & \text{e. } |(2A)^{-1}| & \text{f. } |A^{-1} - 2\text{adj } A| \end{array}$$

5. Let A and B be square matrices of order 4 such that $|A| = -5$ and $|B| = 3$. Find

$$\begin{array}{llll} \text{a. } |2AB| & \text{b. } |\text{adj}(AB)| & \text{c. } |5A^{-1}B^T| & \text{d. } |A^T B^{-1} A^2| \end{array}$$

6. Find all values of k for which the matrix is invertible

$$\text{a. } A = \begin{pmatrix} 1 & 3 \\ k & 2 \end{pmatrix} \qquad \text{b. } B = \begin{pmatrix} m & 1 & 3 \\ 1 & 3 & 2 \\ -1 & 4 & 5 \end{pmatrix} \qquad \text{c. } C = \begin{pmatrix} m & 2 & 0 \\ 1 & m & 1 \\ 2 & 3 & 1 \end{pmatrix}$$

7. Find the eigenvalues and corresponding eigenvectors of the matrix

$$\text{a. } A = \begin{pmatrix} -3 & 5 \\ 10 & 2 \end{pmatrix} \qquad \text{b. } B = \begin{pmatrix} 5 & 4 \\ 2 & 1 \end{pmatrix}$$
$$\text{c. } C = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 3 & 0 \\ 0 & 0 & 4 \end{pmatrix} \qquad \text{d. } D = \begin{pmatrix} -3 & 2 & -1 \\ 0 & 1 & 0 \\ 4 & 1 & 1 \end{pmatrix}$$

8. Find the characteristic polynomial of the matrix

$$\text{a. } A = \begin{pmatrix} 3 & 5 \\ 1 & 2 \end{pmatrix} \qquad \text{b. } B = \begin{pmatrix} 1 & 3 \\ 1 & 3 \end{pmatrix}$$
$$\text{c. } C = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 2 & 1 \\ 1 & 2 & 1 \end{pmatrix} \qquad \text{d. } D = \begin{pmatrix} 1 & 2 & -1 \\ 0 & 1 & 2 \\ -1 & 1 & 1 \end{pmatrix}$$

9. Find the determinant of the matrix $A = \begin{pmatrix} 5 & 1 & 2 & 4 \\ 1 & 0 & -1 & -3 \\ 1 & 1 & 6 & 1 \\ 1 & 0 & 0 & -4 \end{pmatrix}$

Chapter 5: The Vector Space R^n

1. Let $x = (-1, -2, -2)$, $u = (0, 1, 4)$, $v = (-1, 1, 2)$ and $w = (3, 1, 2)$ in R^3 . Find scalars a, b and c such that $x = au + bv + cw$

2. Write v as a linear combination of u and w, if possible, where $u = (1, 2)$, $w = (1, -1)$

a. $v = (0, 1)$ b. $v = (2, 3)$ c. $v = (1, 4)$ d. $v = (-5, 1)$

3. Determine whether the set S is linearly independent or linearly dependent

a. $S = \{(-2, 2), (3, 5)\}$ b. $S = \{(1, -3), (-3, 9)\}$ c. $S = \{(0, 1)\}$

d. $S = \{(0, 0)\}$ e. $S = \{(-1, 2), (3, 1), (2, 1)\}$ f. $S = \{(-1, 2, 3), (1, 3, 5)\}$

g. $S = \{(1, -2, 2), (2, 3, 5), (3, 1, 7)\}$ h. $S = \{(-1, 2, 1), (2, 4, 0), (3, 1, 1)\}$

i. $S = \{(1, -2, 2, 1), (1, 2, 3, 5), (-1, 3, 1, 7)\}$

j. $\{X - Y + Z, 3X + Z, X + Y - Z\}$, where $\{X, Y, Z\}$ is an independent set of vectors

4. For which values of k is each set linearly independent?

a. $S = \{(-1, 2, 1), (k, 4, 0), (3, 1, 1)\}$ b. $S = \{(-1, k, 1), (1, 1, 0), (2, -1, 1)\}$

c. $S = \{(k, 1, 1), (1, k, 1), (1, 1, k)\}$ d. $S = \{(1, 2, 1, 0), (-2, 1, 1, -1), (-1, 3, 2, k)\}$

5. Find all values of m such that the set S is a basis of R^3

a. $S = \{(1, 2, 1), (m, 1, 0), (-2, 1, 1)\}$ b. $S = \{(-1, m, 1), (1, 1, 0), (m, -1, -1)\}$

6. Find a basis for and the dimension of the subspace U

a. $U = \{(2s - t, s, s + t) \mid s, t \in R\}$ b. $U = \{(s - t, s, t, s + t) \mid s, t \in R\}$

c. $U = \{(0, t, -t) \mid t \in R\}$ d. $U = \{(x, y, z) \mid x + y + z = 0\}$

e. $U = \{(x, y, z) \mid x + y + z = 0, x - y = 0\}$ f. $U = \text{span}\{(1, 2, 3), (2, 3, 4), (3, 5, 7)\}$

g. $U = \text{span}\{(1, 2, 4), (-1, 3, 4), (2, 3, 1)\}$ h. $U = \text{span}\{(1, 2, 1, 1), (2, 1, -1, 0), (3, 3, 0, 1)\}$

7. Find a basis for and the dimension of the solution space of the homogeneous system of linear equations.

a.
$$\begin{cases} -x + y + z = 0 \\ 3x - y = 0 \\ 2x - 4y - 5z = 0 \end{cases}$$

b.
$$\begin{cases} x + 2y - 4z = 0 \\ -3x - 6y + 12z = 0 \end{cases}$$

c.
$$\begin{cases} x + y + z + t = 0 \\ 2x + 3y + z = 0 \\ 3x + 4y + 2z + t = 0 \end{cases}$$

8. Find all values of m for which x lies in the subspace spanned by S

a. $x = (-3, 2, m)$ and $S = \{(-1, -1, 1), (2, -3, -4)\}$

b. $x = (4, 5, m)$ and $S = \{(1, -1, 1), (2, -3, 4)\}$

c. $x = (m+1, 5, m)$ and $S = \{(1, 1, 1), (2, 3, 1), (3, 4, 2)\}$

d. $x = (3, 5, 7, m)$ and $S = \{(1, 1, 1, -1), (1, 2, 3, 1), (2, 3, 4, 0)\}$

9. Find the dimension of the subspace

$U = \text{span}\{(-2, 0, 3), (1, 2, -1), (-2, 8, 5), (-1, 2, 2)\}$

10. Let $A = \begin{pmatrix} 1 & 2 & 2 & -1 \\ 3 & 6 & 5 & 0 \\ 2 & 2 & 1 & 2 \end{pmatrix}$. Find $\dim(\text{col } A)$ and $\dim(\text{row } A)$