

Submitted by
Julian Hofer, B.Sc.


Submitted at
**Institute of Semicon-
ductor and Solid State
Physics**

Supervisor
**Prof. Dr. Armando
Rastelli**

Co-Supervisor
DI Christian Schimpf

month year

Exciting and Resolving Quantum Dot Emission with Adiabatic Rapid Passage and Fabry Perot Interferometer



Master Thesis
to obtain the academic degree of
Diplom-Ingenieur
in the Master's Program
Technische Physik

Eidesstattliche Erklärung

Ich erkläre an Eides statt, dass ich die vorliegende Masterarbeit selbstständig und ohne fremde Hilfe verfasst, andere als die angegebenen Quellen und Hilfsmittel nicht benutzt bzw. die wörtlich oder sinngemäß entnommenen Stellen als solche kenntlich gemacht habe. Die vorliegende Masterarbeit ist mit dem elektronisch übermittelten Textdokument identisch.

Datum

Unterschrift

Acknowledgement

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special contents, but the length of words should match the language.

This document is set in Palatino, compiled with pdfL^AT_EX₂ε and Biber.

The L^AT_EX template from Karl Voit is based on KOMA script and can be found online:

<https://github.com/novoid/LaTeX-KOMA-template>

Abstract

This is a placeholder for the abstract. It summarizes the whole thesis to give a very short overview. Usually, this the abstract is written when the whole thesis text is finished.

Contents

Abstract

1	Introduction	1
2	Quantum Dot	3
2.1	Processing	3
2.2	Properties of our dots	3
2.3	Adiabatic Rapid Passage	3
3	Chirp	5
4	Scanning Fabry-Pérot Interferometer	7
4.1	Motivation	7
4.2	Theory	7
4.2.1	Gaussian Beam	7
4.2.2	Fabry-Pérot Interferometer	7
4.2.3	Resonator losses	7
4.2.4	Resonance frequencies and free spectral range	8
4.2.5	Airy distribution of the Fabry-Pérot interferometer	9
4.2.6	Simulation	9
4.3	Setup	10
4.3.1	Flat mirrors	10
4.3.2	Concave mirrors	10
4.3.3	Confocal setup	10
4.4	Measurements and Results	10
	Bibliography	13

List of Figures

4.1	Fabry-Pérot interferometer with electric field mirror reflectivities r_1 and r_2	9
4.2	9

1 Introduction

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special contents, but the length of words should match the language.

This is the second paragraph. Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special contents, but the length of words should match the language.

And after the second paragraph follows the third paragraph. Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special contents, but the length of words should match the language.

After this fourth paragraph, we start a new paragraph sequence. Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and

some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special contents, but the length of words should match the language.

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special contents, but the length of words should match the language.

2 Quantum Dot

2.1 Processing

2.2 Properties of our dots

Table 2.1: My caption		
Quantum dot emission	Energy	Frequency
Center	(1.38 to 2.07) eV	$(3.33 \text{ to } 5.00) \times 10^{14} \text{ Hz}$
Spectral range	(100 to 500) μeV	$(24.20 \text{ to } 120.90) \times 10^9 \text{ Hz}$

2.3 Adiabatic Rapid Passage

3 Chirp

Hallo [2]

4 Scanning Fabry-Pérot Interferometer

4.1 Motivation

Resolve QD emission line.

4.2 Theory

4.2.1 Gaussian Beam

Dot-Spectra in far field is (TEM₀₀).

4.2.2 Fabry-Pérot Interferometer

The Fabry-Pérot interferometer is an optical resonator developed by Charles Fabry and Alfred Pérot. An incoming light beam will only be transmitted through the resonator consisting of two semi-transparent mirrors if it fulfils the resonance condition.[4]

4.2.3 Resonator losses

For the following discussion of the Fabry-Pérot interferometer, a two-mirror-resonator with the reflecting surfaces facing each other and air as medium in between is assumed. The time the light needs for one roundtrip is then given by [3]

$$t_{RT} = \frac{2l}{c} \quad (4.1)$$

where l is the geometrical length of the resonator and c is the speed of light in air.

The photon-decay time $\tau_c \nu$ of the interferometer is then given by

$$\frac{1}{\tau_c} = -\frac{\ln(R_1 \cdot R_2)}{t_{RT}} \quad (4.2)$$

where R_1 and R_2 are the corresponding intensity reflectivities of the mirrors.

The number of photons at frequency ν inside the resonator is described by the differential rate equation

$$\frac{d}{dt}\varphi(t) = -\frac{1}{\tau_c}\varphi(t). \quad (4.3)$$

With a number φ_s of photons at $t = 0$ the integration gives

$$\varphi(t) = \varphi_s e^{-t/\tau_c} \quad (4.4)$$

4.2.4 Resonance frequencies and free spectral range

The round-trip phase shift at frequency ν is given by

$$2\phi(\nu) = 2\pi\nu t_{RT} = 2\pi\nu \frac{2l}{c} \quad (4.5)$$

where $\phi(\nu)$ is the single-pass phase shift between the mirrors.

Resonances are visible for frequencies ν at which the light interferes constructively after one round trip. Two adjacent resonance frequencies differ in their round trip phase shift by 2π . Hence, the free spectral range $\Delta\nu_{FSR}$, the frequency difference between two adjacent resonance frequencies, can be calculated from equation (4.5)

$$2\Delta\phi = 2\pi \quad (4.6)$$

$$\Rightarrow 2\pi\Delta\nu_{FSR} \frac{2l}{c} = 2\pi \quad (4.7)$$

$$\Rightarrow \Delta\nu_{FSR} = \frac{c}{2l} \quad (4.8)$$

4.2.5 Airy distribution of the Fabry-Pérot interferometer




Figure 4.1: Fabry-Pérot interferometer with electric field mirror reflectivities r_1 and r_2

[1]

4.2.6 Simulation




Figure 4.2:

4.3 Setup

4.3.1 Flat mirrors

4.3.2 Concave mirrors

4.3.3 Confocal setup

4.4 Measurements and Results

Appendix

Bibliography

- [1] *Fabry-Pérot interferometer* - Wikipedia. URL: https://en.wikipedia.org/wiki/Fabry%E2%80%93P%C3%A9rot_interferometer (visited on 05/06/2019) (cit. on p. 9).
- [2] Toshiyuki Hirayama and Mansoor Sheik-Bahae. “Real-time chirp diagnostic for ultrashort laser pulses.” In: *Optics Letters* 27.10 (May 15, 2002), p. 860. ISSN: 0146-9592, 1539-4794. DOI: 10.1364/OL.27.000860. URL: <https://www.osapublishing.org/abstract.cfm?URI=ol-27-10-860> (visited on 12/11/2018) (cit. on p. 5).
- [3] Nur Ismail et al. “Fabry-Pérot resonator: spectral line shapes, generic and related Airy distributions, linewidths, finesses, and performance at low or frequency-dependent reflectivity.” In: *Optics Express* 24.15 (July 25, 2016), p. 16366. ISSN: 1094-4087. DOI: 10.1364/OE.24.016366. URL: <https://www.osapublishing.org/abstract.cfm?URI=oe-24-15-16366> (visited on 02/15/2019) (cit. on p. 7).
- [4] Timo Kaldewey et al. “Coherent and robust high-fidelity generation of a biexciton in a quantum dot by rapid adiabatic passage.” In: *Physical Review B* 95.16 (Apr. 10, 2017). ISSN: 2469-9950, 2469-9969. DOI: 10.1103/PhysRevB.95.161302. arXiv: 1701.01371. URL: <http://arxiv.org/abs/1701.01371> (visited on 12/11/2018) (cit. on p. 7).