



Road to Your First Pull-Up

A 12-week bodyweight program for people who can hang from a bar but can't pull up yet. It takes 2 sessions a week, plus a few minutes of daily posture work.



NAME _____

PHYSIOTHERAPIST _____

START DATE _____

TARGET TEST DATE _____

Phase 1

WEEKS 1-4

Foundation

Shoulder blade control, grip, pulling strength and a strong trunk.

Phase 2

WEEKS 5-8

Build the pull

Band-assisted pull-ups, top-position holds and slow lowering.

Phase 3

WEEKS 9-12

First pull-up

Less help from the band, paused negatives and pull-up attempts.

YOUR WEEK

Every day Wall angels: 2-3 minutes (page 2)

Day A Warm-up (10 min) + phase workout (25-35 min)

Day B Same as Day A, 2-3 days later (e.g. Mon & Thu)

EQUIPMENT

Pull-up bar · resistance loop bands (light, medium, heavy) · wall · mat · sturdy table · step or sturdy box

HOW HARD?

Finish each set with 2-3 good reps left in the tank, about **7-8 out of 10** effort. Rest 90 seconds to 2 minutes between pulling sets.

WHEN TO MOVE ON

Each phase ends with a checklist. Move on when you pass it, not just because the weeks are up. Taking 8 weeks or more than 12 is normal.

SAFETY: STOP AND CONTACT YOUR PHYSIOTHERAPIST IF YOU NOTICE


- sharp or pinching pain in your shoulder, elbow or neck
- pins and needles or numbness down your arm
- your shoulder feels like it slips or gives way
- pain that is worse the next morning

Mild muscle soreness is normal. Any discomfort during exercise should stay at **3/10 or less** and settle by the next day. Check the band and bar before every session.



Wall angels

Do these every day, and as the last part of every warm-up. They teach your shoulder blades to move overhead while your ribs stay down. That's the position you want when you hang from the bar.




Daily • 2 sets × 8–10 slow reps (3 seconds up, 3 seconds down)

SET-UP

1. Stand with your heels about a hand's length from the wall and your knees soft.
2. Rest the back of your head, your upper back and your tailbone on the wall.
3. Start in the "goalpost" position: elbows at shoulder height and bent to 90°, forearms toward the wall.

MOVE

4. **Breathe in** through your nose as you slowly slide your arms up into a "Y". Stop where your ribs start to lift or your arms leave the wall.
5. **Breathe out** through your mouth as you slide your arms back down, drawing your elbows toward your back pockets.



KEEP YOUR RIBS DOWN: STOP THEM FLARING

- ✓ **Breathe in wide, not up.** As your arms rise, let the breath widen your sides and back ribs instead of lifting your chest. Your lower ribs stay down toward your hips.
- ✓ **Breathe out to reset.** As your arms lower, breathe out fully and feel your lower ribs soften down again before the next rep.
- ✓ **Zip up gently.** Tighten your lower tummy to about 20–30% effort. That's just enough to stop your ribs popping forward, not a hard brace.
- ✓ **Keep the low-back gap small.** A flat hand should just fit behind your low back. If the gap grows as your arms go up, you've gone too high.
- ✓ **Keep your chin gently tucked** and the back of your head on the wall. Don't let your chin poke up.
- ✓ **Work within your range.** Your hands don't have to touch the wall at the top. Ribs down matters more than height.

Easier: step your feet further from the wall and bend your knees more, or do the same movement lying on your back with your knees bent ("floor angels").

Harder: slow it down to a 4-second breath in on the way up and a 6-second breath out on the way down.

Warm-up

Do these in order before every workout. Finish with 1 set of 8 wall angels.



Breathe in

Breathe out: ribs down

1 90/90 breathing

5 slow breaths

1. Lie on your back with your feet flat on a wall and your hips and knees bent to 90°. Rest one hand on your lower ribs.
2. Breathe in through your nose for 4 seconds. Feel your ribs widen to the sides and into the floor.
3. Breathe out slowly through your mouth for 6–8 seconds, as if fogging a mirror. Let your lower ribs drop down and in. Your low back settles gently toward the floor.
4. Pause for 2–3 seconds before the next breath.

Tip: This sets your ribs in the 'down' position you will keep during every exercise.



Round up

Let your back sag

2 Cat-camel

8–10 slow reps

1. Start on your hands and knees, with your hands under your shoulders and your knees under your hips.
2. Breathe out as you round your back up, tucking your tailbone and chin.
3. Breathe in as you let your back gently sag and look slightly forward.
4. Move slowly and only in a range that feels comfortable.



Start

Finish


3 Band pull-aparts

2 sets × 12–15

1. Hold a light band at shoulder height. Keep your arms straight and your hands shoulder-width apart.
2. Pull the band apart until it reaches your chest, drawing your shoulder blades back and down.
3. Keep your ribs down and your shoulders away from your ears. Return slowly.

Easier: Use a lighter band or start with your hands wider apart.

Warm-up




4 Scapular push-ups

2 sets × 10

1. Start on your hands and knees, with your hands under your shoulders and your arms straight.
2. Keeping your elbows straight, let your chest sink between your shoulder blades.
3. Push the floor away to spread your shoulder blades apart. Only your shoulder blades move.

Harder: Do these from your knees in a kneeling plank, then from your toes.



5 Dead bug

2 sets × 6 each side

1. Lie on your back with your arms pointing to the ceiling and your hips and knees at 90°.
2. Breathe out to set your ribs down and your low back gently into the floor.
3. Slowly reach one arm overhead and the opposite leg out, only as far as your low back stays down.
4. Return to the start and switch sides. Breathe out on each reach.


Easier: Move only your legs, or keep the knee bent as you tap your heel toward the floor.

6 • Wall angels 1 set × 8 (page 2). Then start your phase workout.

Foundation

Build shoulder blade control, grip endurance and basic pulling strength.

EXERCISE	SETS × REPS / TIME
Scapular pull-ups + active hang	3 × 5-8 (2-second hold), then 2 × 15-30 s hang
Band lat pull-down	3 × 10-12
Prone Y raise	2 × 10 (2-second hold)
Inverted row	3 × 6-10
Hollow body hold (tucked)	3 × 15-30 s




A Scapular pull-ups (active hang)

3 sets × 5-8, 2-second hold

1. **Relaxed hang (left photo):** hang from the bar with straight arms and let your shoulders ride up toward your ears.
2. **Active hang (right photo):** keep your arms straight and pull your shoulder blades down into your back pockets. Your shoulders move away from your ears and your whole body rises a few centimetres.
3. Hold for 2 seconds, then slowly let your shoulders ride back up. That's 1 rep.
4. Finish with 2 holds of 15-30 seconds in the active position.

Tip: Your elbows never bend. Only your shoulder blades move. This is the first thing that happens in every pull-up. Keep your toes on a box if your grip needs a break.



B Band lat pull-down

3 sets × 10-12

1. Loop a band over the pull-up bar. Kneel tall facing the bar, with your ribs down.
2. Start with your arms reaching up. Pull your elbows down and back toward your ribs until your hands are near your shoulders.
3. Squeeze for 1 second, then return over 2 seconds without shrugging.

Harder: Use a heavier band, or pause for 3 seconds at the bottom.

Ready for the next phase when you can do:

- 3 × 8 scapular pull-ups with 2-second holds
- a 30-second active hang
- 3 × 10 inverted rows
- a 30-second tucked hollow hold

Foundation, continued



Top: arms resting • Bottom: arms lifted, thumbs up

C Prone Y raise

2 sets × 10, 2-second hold

1. Lie face down with your forehead on a folded towel and your arms overhead in a 'Y', thumbs up.
2. Draw your shoulder blades down toward your back pockets, then lift your arms a few centimetres.
3. Hold for 2 seconds and lower slowly. Keep your neck long and don't arch your back.

Tip: These target the lower trapezius, which helps start the pull-up.



Start

Finish


D Inverted row

3 sets × 6–10

1. Set a sturdy bar at about hip height (a Smith machine or rack bar at the gym). At home, the edge of a heavy table works: check that it cannot tip.
2. Lie under the bar and grip it with your hands shoulder-width apart. Keep your body straight from ears to ankles and squeeze your glutes.
3. Pull your chest to the bar, elbows angled toward your ribs. Pause for 1 second, then lower over 3 seconds.

Easier: Bend your knees with your feet flat.

Harder: Straighten your legs, or add a 3-second pause at the top.



Tucked

Full

E Hollow body hold

3 sets × 15–30 seconds

1. Lie on your back. Breathe out to set your ribs down and press your low back into the floor.
2. Lift your head and shoulders. Start tucked: knees over hips, arms reaching toward your feet.
3. Hold with small breaths. Stop the set if your low back lifts or your ribs flare.


Harder: Straighten one leg, then both. Then take your arms overhead ('Full').

Build the pull

Build strength at the top and on the way down first, while you're fresh. Then practise the whole pull-up path with a band.

EXERCISE	SETS × REPS / TIME
Active hang primer	2 × 20 s
Negative pull-up	3 × 3 (3–5 s lower)
Flexed-arm hang	3 × 10–20 s
Band-assisted pull-up	3 × 4–6
Inverted row, legs straight	3 × 8–10
Hollow body hold, legs straighter	3 × 20–30 s

Start with an active hang primer: 2 × 20 seconds in the active position from Phase 1 (page 5), then begin exercise A.




A Negative (lowering) pull-up

3 sets × 3, lowering over 3–5 seconds

1. Start at the top (right photo): step up on a box so your chin is over the bar.
2. Lift your feet and lower yourself slowly, counting, until your arms are straight (left photo). Keep your shoulders 'active' at the bottom.
3. Step back up for the next rep. Don't jump up from the bottom.

Tip: Slow lowering (eccentric) work builds strength quickly. Expect some muscle soreness in the first week.

Easier: If you drop in under 2 seconds, do the lowering with your knee in a band until you can control it.




B Flexed-arm hang

3 sets × 10–20 seconds

1. Step up on a sturdy box or chair so your chin is above the bar.
2. Grip the bar, lift your feet and hold. Keep your shoulders down, your elbows tucked in and your ribs down.
3. Step back down to finish. Don't drop.

Tip: Keep breathing. Don't hold your breath.

Build the pull, continued



C Band-assisted pull-up

3 sets × 4–6

1. Loop a strong band over the bar and step up on a box. Put one foot in the loop (most help) or kneel into it with one knee (less help).
2. Start from an active hang: shoulders down, ribs down.
3. Pull your elbows down to your ribs until your chin clears the bar. Lower over 3 seconds.
4. To finish, step back onto the box before you take your foot or knee out of the band.

Tip: Choose the set-up where the last rep is hard but still clean. Progress in this order: foot in the loop → knee in the loop → next lighter band. Move one step once you can do 3 × 6 with good form.

D • Inverted row 3 × 8–10 with legs straight (page 6). **E • Hollow body hold** 3 × 20–30 s, straighten your legs as you are able (page 6).

Ready for the next phase when you can do:


- a 20-second flexed-arm hang
- 3 × 3 negatives, lowering over 5 seconds
- 3 × 6 band-assisted pull-ups with a medium band

Your first pull-up

Less help, more control. Keep every rep clean. Grinding out ugly reps won't get you there faster.

EXERCISE	SETS × REPS / TIME
Active hang primer	2 × 20 s
Pull-up attempts	3–5 singles, 2–3 min rest
Slow negatives with pauses	3 × 2 (8–10 s each)
Band-assisted pull-up, less help	3 × 5–6
Hollow body hold (full)	3 × 30 s
Inverted row, 2-second pause	3 × 10
Flexed-arm hang	2 × 20–30 s

Start with an active hang primer: 2 × 20 seconds in the active position from Phase 1 (page 5), then begin exercise A.




A Pull-up attempts

3–5 single attempts, 2–3 minutes' rest

1. Do these straight after your warm-up, while you are fresh.
2. From an active hang, pull your elbows down to your ribs and drive your chest toward the bar.
3. If you don't reach the bar, pull as high as you can, hold for 2 seconds, then lower slowly. That still counts as training.

Tip: Test day: in week 12, or once you pass the test-day checklist on page 10. Rest well the day before, warm up, and give it 1–2 attempts.




B Slow negatives with pauses

3 sets × 2, 8–10 seconds per rep

1. Start at the top as for negatives.
2. Lower slowly, pausing for 2 seconds at the top, at halfway (elbows at 90°) and just before your arms are straight. Each rep takes 8–10 seconds from top to bottom.
3. Step back up between reps.

Your first pull-up, continued




C Band-assisted pull-up

3 sets × 5–6, starting one step less help than you ended Phase 2. Move down a step once 3 × 6 feels clean (usually every 1–2 weeks).

1. Loop a strong band over the bar and step up on a box. Put one foot in the loop (most help) or kneel into it with one knee (less help).
2. Start from an active hang: shoulders down, ribs down.
3. Pull your elbows down to your ribs until your chin clears the bar. Lower over 3 seconds.
4. To finish, step back onto the box before you take your foot or knee out of the band.

Tip: Choose the set-up where the last rep is hard but still clean. Progress in this order: foot in the loop → knee in the loop → next lighter band. Move one step once you can do 3 × 6 with good form.



D Hollow body hold

3 sets × 30 seconds, working toward the full position

1. Lie on your back. Breathe out to set your ribs down and press your low back into the floor.
2. Lift your head and shoulders. Start tucked: knees over hips, arms reaching toward your feet.
3. Hold with small breaths. Stop the set if your low back lifts or your ribs flare.

Harder: Straighten one leg, then both. Then take your arms overhead ('Full').

E • Inverted row 3 × 10, legs straight, with a 2-second pause at the top (page 6). **F • Flexed-arm hang** 2 × 20–30 seconds (page 7).

Ready for test day when you can do:

- a 30-second flexed-arm hang
- a slow negative taking 8–10 seconds, with control
- 3 × 6 band-assisted pull-ups with your lightest band, knee in the loop
- a 30-second full hollow hold

After your first pull-up: do 3 sets of 1–2 reps, then work toward 3 reps in a row.

Progress log

Write down your best set for each session. Small changes add up.

Wk	Date	Main pull (scap pull-ups / band pull-ups / attempts)	Hang / hold (seconds)	Negatives (seconds lowering)	Rows · hollow	Pain 0–10 · notes
1						
2						
3						
4						
5						
6						
7						
8						
9						
10						
11						
12						

WHY THESE EXERCISES: THE RESEARCH

- Pull-ups mainly work the lats (about 117–130% of maximum voluntary contraction), biceps and infraspinatus. The **lower trapezius and pecs switch on first** to start the pull, which is why scapular pull-ups and prone Y raises come first.
Youdas JW et al. J Strength Cond Res 2010;24(12):3404–14. doi:10.1519/JSC.0b013e3181f1598c
- Different grips (overhand, underhand, neutral) activate the muscles similarly over a full rep, so use whichever grip is comfortable.
Dickie JA et al. J Electromyogr Kinesiol 2017;32:30–36. doi:10.1016/j.jelekin.2016.11.004
- Face-down (prone) exercises target the middle and lower trapezius with little upper-trapezius (shrugging) activity.
Cools AM et al. Am J Sports Med 2007;35(10):1744–51. doi:10.1177/0363546507303560
- Wall slides activate the serratus anterior, and push-up "plus" variations favour serratus over upper trapezius.
Castelein B et al. J Orthop Sports Phys Ther 2016;46(3):184–93. doi:10.2519/jospt.2016.5927 · Maenhout A et al. Br J Sports Med 2010;44(14):1010–5. doi:10.1136/bjsm.2009.062810
- Gently tightening the abdominals increased serratus and trapezius activity during wall slides. This supports the "ribs down, zip up" cue.
Vega Toro AS et al. Man Ther 2016;25:11–8. doi:10.1016/j.math.2016.05.331
- Eccentric (lowering) training gives strong strength gains, which is why negatives are used in Phases 2–3.
Douglas J et al. Sports Med 2017;47(5):917–41. doi:10.1007/s40279-016-0628-4

References sourced via PubMed. This program is general education prepared by Holistic Physiotherapy & Wellness. It does not replace an individual assessment. Talk to your physiotherapist before starting if you have had shoulder surgery, a dislocation or ongoing neck or shoulder pain.