

Exercise 4

Holo

Sunday 30th July, 2023

Exercise 1. *Dense projections and Quotient posets*

Part 1.1.

Let $\pi, \mathbb{P}, \mathbb{Q}$ and $\Gamma_{\mathbb{Q}}^{\mathbb{P}}$ be as in the question, let G be any M -generic subset of \mathbb{P} and let $\Gamma = (\Gamma_{\mathbb{Q}}^{\mathbb{P}})_G, \Gamma' = (\tilde{\pi}''\Gamma_{\mathbb{P}})_G$.

Let $p, q \in \Gamma$ and $p', q' \in \Gamma'$ be strengthening of p, q respectively, let $t, r \in G$ be such that $\pi(t) = p', \pi(r) = q'$, because G is generic we have $t, r < w \in G$, and because π is a weak homomorphism, $\pi(w) > q', p'$ hence $\pi(w) > q, p$, and almost by definition $\pi(w) \in \Gamma$.

Because Γ is closed downwards by definition, we only left to show that it intersect with every maximal anti-chain of \mathbb{Q} that is in M , let A be such maximal anti-chain. For each $x \in \mathbb{Q}$ let x^\dagger be a maximal anti-chain of $\{y \in \pi''\mathbb{P} \mid y \geq x\}$, and let $A^\dagger = \bigcup \{y^\dagger \mid y \in A\}$, because $\pi''\mathbb{P}$ is dense, we have that A^\dagger is a maximal anti-chain as well.

Let A^\ddagger be the inverse image of A^\dagger by π , because π preserve incompatibility we must have that A^\ddagger contains a maximal anti-chain, let $t \in A^\ddagger \cap G$, $\pi(t) \in A^\dagger \cap \Gamma$. By construction of A^\dagger there exists some $p \in A$ such that $\pi(t) > p$, and because Γ is closed downwards we have that $p \in A \cap \Gamma$, hence $M[G] \models \text{"}\Gamma \text{ is generic over } M \text{ for } \mathbb{Q}\text{"}$.

Finally because G was arbitrary we can conclude that $0_{\mathbb{P}}$ forces that.

Part 1.2.

Let $G \subseteq \mathbb{P}$ be generic.

Notice that H is exactly Γ from (1.1), so H is generic over M for \mathbb{Q} .

Because being downwards closed is downwards absolute (from \mathbb{P} to $\mathbb{P}/_{\pi}H$) and every 2 elements having a common extension in the subset is absolute (again from \mathbb{P} to $\mathbb{P}/_{\pi}H$) we only need to care about it intersecting every maximal anti-chain of $\mathbb{P}/_{\pi}H$ in $M[H]$.

Let $p, q \in \mathbb{P}/_{\pi}H$, by definition $\pi(p), \pi(q) \in H \implies \pi(p) \parallel \pi(q) \implies p \parallel q$, the last implication is because π is homomorphism and the one before that is because H is a generic filter.

Let $A \subseteq \mathbb{P}/_{\pi}H$ be any maximal anti-chain in $M[H]$, that is $A = \{p\}$ for some $p \in \mathbb{P}/_{\pi}H$ (as any 2 elements of the quotient poset are compatible), so $A \in M$. Extend A to A' a maximal anti-chain in \mathbb{P} , because G is M generic for \mathbb{P} we know that $G \cap A' \neq \emptyset$, but $G \subseteq \mathbb{P}/_{\pi}H$ so $G \cap A' = A \implies G \cap A = A \neq \emptyset$.

For the other direction, let G be any subset of \mathbb{P} such that H is M -generic for \mathbb{Q} and G is $M[H]$ -generic for $\mathbb{P}/_\pi H$.

Similarly to before, every 2 elements in $\mathbb{P}/_\pi H$ are compatible, in particular any 2 elements in G are compatible, as $G \subseteq \mathbb{P}/_\pi H$.

Let $p \in G$ and $q < p$, because π is a weak homomorphism we have that $\pi(q) < \pi(p) \in H$, so $\pi(q) \in H \implies q \in \mathbb{P}/_\pi H$, because $q < p \in \mathbb{P}/_\pi H$ and G is closed downwards in $\mathbb{P}/_\pi H$ we must have that $q \in G$.

Lastly, let $p, q \in \mathbb{P}/_\pi H$, let $D_{p,q} = \{r \in \mathbb{P} \mid r \geq p, q \vee r \perp p \vee r \perp q\}$ and $D' = \pi'' D_{p,q}$, because $D_{p,q}$ is dense and π is dense homomorphism we know that D' is dense hence intersecting H , let $r \in D_{p,q}$ be such that $\pi(r) \in D' \cap H$. Because $\pi(r) \in H$ we have that $r \in \mathbb{P}/_\pi H$ but because every 2 elements of $\mathbb{P}/_\pi H$ are compatible in \mathbb{P} , we have that $r \parallel p$ and $r \parallel q$, by construction of $D_{p,q}$ we get that $r \geq p, q$.

That means that every 2 elements in $\mathbb{P}/_\pi H$ are compatible in $\mathbb{P}/_\pi H$, so the only generic filter is $\mathbb{P}/_\pi H$, so $G = \mathbb{P}/_\pi H$.

Now take A to be any maximal anti-chain, $\pi'' A$ is a maximal anti-chain as well, hence $\{\pi(p)\} = H \cap \pi'' A$ hence $G \cap A = \mathbb{P}/_\pi H \cap A = \{p\} \neq \emptyset$

Exercise 2. Collapse Criterion

First note that \mathbb{Q}/q absorbs \mathbb{Q} , indeed for each M -generic for \mathbb{Q}/q we can take the downwards closure and get an M -generic for \mathbb{Q} .

Therefore $0_{\mathbb{Q}/q} \Vdash |\check{\omega}| = |\check{\delta}|$, so \mathbb{Q}/q is not δ .c.c. and has a maximal anti-chain of size δ .

Let \dot{g} be a \mathbb{Q} name that $0_{\mathbb{Q}} \Vdash \dot{g} : \omega \rightarrow \Gamma_{\mathbb{Q}}$ is surjective"

We will construct recursively an embedding from $\delta^{<\omega}$ to \mathbb{Q} as a tree T :

- $T_\Lambda = 0_{\mathbb{Q}}$
- Let $t \in \delta^{n+1}$ and assume T_t is defined, let $C_t = \{c_t^{(\alpha)} \mid \alpha \in \delta\}$ be a maximal anti-chain of size δ in \mathbb{Q}/T_t such that for each $c \in C_t$ we have that c decides the value of $\dot{g}(\check{n})$ and let $T_{t \smallfrown \{i\}} = c_t^{(i)}$

Note that the anti-chain above is always well defined, let $p \in \mathbb{Q}$ be any element and let $A \subseteq \mathbb{Q}/p$ be any maximal anti-chain that decides $\dot{g}(\check{n})$, let $q \in A$ be any element and let $B \subseteq \mathbb{Q}/q$ of cardinality δ , then $A' = A \setminus q \cup B$ is a maximal anti-chain of size δ that decides $\dot{g}(\check{n})$.

This map is clearly an injective homomorphism, to see that it is dense let $p \in \mathbb{Q}$ be any element, remember that $p \Vdash \dot{g} : \omega \rightarrow \Gamma_{\mathbb{Q}}$ is surjective and $\check{p} \in \Gamma_{\mathbb{Q}} = \text{range}(\dot{g})$ which implies $p \Vdash \exists n \in \check{\omega} \dot{g}(n) = \check{p}$ in particular there exists a name \check{n} such that $p \Vdash \dot{g}(\check{n}) = \check{p}$, let m be a real natural such that $p \Vdash \check{m} = \check{n}$, thus $p \Vdash \dot{g}(\check{m}) = \check{p}$.

The set $\text{Lev}_T(m+1)$ is clearly a maximal anti-chain, so either $p \in \text{Lev}_T(m+1)$, in which case we are done, or there exists $q \in \text{Lev}_T(m+1)$ that is compatible with p , because p, q are compatible and both decide $\dot{g}(\check{m})$ they must agree with one another, that is $q \Vdash \dot{g}(\check{m}) = \check{p}$, in particular $q \Vdash \check{p} \in \text{range}(\dot{g}) = \Gamma_{\mathbb{Q}}$ therefore p is inside every generic G that q is in, in other words $p < q$.

Finally, let G be M -generic for \mathbb{Q} , and let H' be the intersection of G with the image of the tree, it is simple to see that H' is M -generic over T let H'' be the pre-image of H' into $\delta^{<\omega}$, and let H be the downwards closure of H'' into an M -generic for $\text{Coll}(\omega, \delta)$.

To see that $M[G] = M[H]$ first note that because H is definable in $M[G]$ (with parameters) it means that $M[H] \subseteq M[G]$, to see the other direction, let H^\dagger be the restriction of H to $\delta^{<\omega}$, let H^\ddagger be the image of H^\dagger under the embedding to T , we can verify that G is exactly the downward closure of H^\ddagger .

First because $H^\ddagger \subseteq G$ the downwards closure of H^\ddagger is clearly a subset of G , and the downwards closure is M -generic for \mathbb{Q} (as T is dense).

Let $p \in G$ be any element, extend $\{p\}$ to a maximal anti-chain A , A intersects with the downwards closure of H^\ddagger , but p is compatible with every element of the downwards closure of H^\ddagger , so the intersection must be $\{p\}$, so p is in the downwards closure of H^\ddagger .

So we got that G is definable in $M[H]$ (with parameters) and hence $M[G] \subseteq M[H]$ and we are done.