

THE ATLAS FRAMEWORK

A Therapeutic Model for Complex Trauma and Neurodivergence for integration with the Atlas app.

Atlas Therapeutic Foundation Document

Christopher Specht

Kintsugi Collective Pty Ltd | Sydney, Australia | March 2026

CONFIDENTIAL - Not for Distribution

Patent Pending – IP Australia

This material is copyrighted by the author.

1 Part I: Who This Framework Is For

This framework is designed from the ground up for a specific population: adults with complex trauma, neurodivergence, co-occurring disorders, and histories that standard healthcare systems have repeatedly failed to hold. This is not a special-case population. This is a primary population. Every theoretical choice, every intervention, every sequencing decision in this document has been evaluated through that lens first.

The reason this framework came into existence is due to any cohesive integrated approach that is modality agnostic, truly recognises the reality and lived experiences of those with the conditions as opposed as those studying those with the conditions.

This framework seeks to remove the clinical layers from existing approaches, and integrate the actual choices of these populations. Their history, their stories, their unique biological make up, all contribute to their waveform vector moving through their lifeline, or world line.

The overall philosophy behind this idea is that whatever helps someone, so long as its not maladaptive, is worth including. For trauma, talk therapy is one of a number of modalities which should be pursued – the reliance of a specific modality is relative to the individual's response to that modality. That is why we must come to accept that this path forward is one of (1) seeking stability relative to our needs and capabilities; (2) accepting spiralling as part of the true path forward (spiralling itself is a relative topic); (3) radical acceptance of self. As you are. We're not here to fix you, you're not broken. You are fractured, and have been for some time. It's time to bring those parts together, a vase fractured can be repaired, one that is broken can never be repaired. We're not promising healing or recovery. We're promising stability, acceptance, understanding and integrity.

The person this framework is built for typically presents with one or more of the following, often in combination:

- Complex PTSD (CPTSD) - developmental or prolonged trauma, frequently beginning in childhood
- Autism Spectrum Disorder (ASD) - including late-diagnosed adults, masked presentations, and high-masking profiles
- Attention Deficit Hyperactivity Disorder (ADHD) - including inattentive presentations, executive function dysregulation, and endogenous dopamine insufficiency
- Substance use as regulatory behaviour - self-medication of a chronically dysregulated nervous system, not primary addiction in the classical model
- Co-occurring anxiety, depression, or dissociative presentations - as secondary expressions of the above, not independent primary disorders
- Histories of repeated clinical failure - multiple practitioners, multiple modalities, repeated relapse, repeated discharge, and repeated reduction to a single presenting complaint

Core principle:

Trauma is fundamentally a disconnection from self (Maté) that produces lasting neurobiological reorganisation (Perry, van der Kolk). For neurodivergent individuals, this reorganisation occurs in a nervous system already wired differently - creating compounding interactions that cannot be addressed by frameworks designed for neurotypical baselines.

Healing is not a return to a neurotypical norm. It is the recovery of navigation rights to one's own life - on one's own terms, within one's own nervous system.

*For this framework, **this relates to the topological deformation which occurs on a very fundamental level, biologically and psychologically speaking.** However, overtime this deformation degrades the overall 'hope' and 'prolonged outcomes' of the individual. Until a modulating force is applied.*

1.1 What This Framework Is Not

- A diagnostic tool. It produces no DSM or ICD classifications.
- A clinical replacement for psychiatry, psychology, or pharmacological management.
- A crisis intervention service.
- A linear recovery programme with defined endpoints.
- Built on the assumption that the person is broken and needs to be fixed.
- Any perceived lack of moral constitution on the part of the individual due to addiction or repeated life disruptions, is reprehensible and lazy.

1.2 What This Framework Is

- A dynamic, longitudinal support architecture for people navigating complex nervous systems in a world not built for them.
- A companion intelligence that stays present without directing, analyses without pathologising, and reflects without judging.
- A spiral model of regulation and integration, expecting cycling, honouring regression, tracking long-arc progress.
- A framework that treats medication, somatic work, parts work, presence practice, and habit architecture as co-equal tools, none primary, all contextual.
- Recognising that the lifeline for people with serious mental health issues is at times challenging and that these challenges don't just disappear with 6 months of "recovery" work.
- The current approaches relating to trauma work are often disparate, 'fringe' or lacking in any scientific grounding.

2 Part II: Theoretical Architecture

The following theorists and frameworks form the foundation of this model. Each is evaluated for its applicability to the primary population, with explicit notes on where adaptation is required for neurodivergent and complex trauma presentations.

2.1 Neurobiological Foundations

2.1.1 Stephen Porges - Polyvagal Theory

The primary nervous system model of this framework. Porges' three-state autonomic hierarchy - ventral vagal (safe and social), sympathetic (mobilised), and dorsal vagal (shutdown) - directly describes the regulatory landscape of CPTSD and neurodivergence.

- Chronic trauma shifts the nervous system's default toward sympathetic activation or dorsal vagal shutdown.
- ASD nervous systems show atypical neuroception - the automatic safety/danger detection system fires differently, producing social withdrawal, sensory overwhelm, and shutdown that is neurological, not behavioural.
- ADHD nervous systems show irregular vagal tone, contributing to emotional dysregulation that is frequently misattributed to mood disorder.
- Co-regulation - the therapeutic relationship as a regulated nervous system signal - is the primary mechanism of nervous system change.

Adaption for primary population: This framework assumes that every individual's nervous system, whilst biologically the 'same', are different and function comparatively differently (responses to external stimuli) between individuals. It does not presume therapeutic responses being exactly similar across the population.

2.1.2 Deb Dana - Polyvagal Theory in Clinical Practice

Dana's clinical application of Polyvagal Theory provides the practical implementation layer. Her autonomic mapping tools and glimmer/trigger identification framework are directly usable for self-understanding without clinical mediation - which aligns with the companion intelligence architecture.

2.1.3 Bruce Perry - Neurosequential Model of Therapeutics (NMT)

The single most important sequencing principle in this framework. Somatic regulation precedes relational engagement, which precedes cognitive reframing. This is not a preference - it is neurobiology. Higher cortical functions cannot be effectively accessed while brainstem and limbic systems remain dysregulated.

Critical adaptation for ASD:

Perry's model assumes interoceptive access as the pathway into somatic work. Alexithymia - difficulty identifying and describing internal emotional states - affects an estimated 50-85%

of autistic individuals. Standard somatic entry points ('notice what you feel in your body') may be inaccessible or actively misleading. This framework uses external-signal proxies (HRV, sleep data, wearable telemetry) and pattern-over-time recognition rather than requiring real-time interoceptive reporting as the primary data source. However, real time pattern recognition is something that the individual can develop and utilise to their benefit. Overall, the message should be made clear, conditions like Alexithymia should not be seen to preclude someone from any modality within this charter.

2.1.4 Bessel van der Kolk - The Body Keeps the Score

Trauma is stored in the body, not primarily in narrative memory. Body-based interventions are therefore not supplementary - they are primary. This research directly validates the somatic-first sequencing and undermines the assumption that talk therapy can address complex trauma through narrative alone.

- Trauma memory is stored as sensory fragments, not coherent narratives - making verbal processing insufficient as a primary modality.
- The body's alarm system can be triggered without conscious memory - 'I don't know why I reacted that way' is neurologically accurate, not avoidance.
- Body-based interventions (EMDR, Somatic Experiencing, yoga, rhythm, breath) produce neurobiological changes that talking cannot replicate.
 - *NB: EMDR, like many other modalities mention herein must be undertaken with trained licensed professionals. Many commercial apps claim to provide this therapy, however it can elicit troubling memories and emotions which must be managed in a professional setting.*

2.1.5 Peter Levine - Somatic Experiencing

Levine's trauma resolution model - completing thwarted survival responses stored in the nervous system - provides the primary somatic intervention model. His 'pendulation' concept (moving between activation and resourcing rather than sustained exposure) is specifically suited to high-Ω nervous systems that cannot tolerate prolonged trauma contact.

Critical adaptation for CPTSD (van der Kolk & Levine):

The inclusion of unitary modalities or approaches is only half the picture, any integrated approach must incorporate both somatic experiencing and a talk therapy. The individual must release the trapped trauma from their bodies, as part of the cyclical nature of sustainable living post trauma.

2.1.6 Pat Ogden - Sensorimotor Psychotherapy

Ogden's sensorimotor model bridges somatic experiencing and parts work by tracking the body's habitual postural and movement patterns as carriers of trauma. For individuals who have difficulty accessing emotion through introspection but can observe their own physical patterns,

sensorimotor tracking provides an accessible entry point that bypasses the interoception problem.

2.1.7 Lisa Feldman Barrett - Constructed Emotion Theory

Barrett's model - that emotions are not universal biological programmes but are constructed by the brain from interoceptive signals, past experience, and conceptual knowledge - has direct implications for alexithymic and ASD populations. The standard model of emotion (you feel something, you recognise and name it) does not describe how many autistic nervous systems work. Barrett's framework explains why: the predictive processing mechanisms that construct emotional experience are different, not absent.

This framework supports emotional tracking through components (body state + context + behaviour) rather than requiring direct emotion label identification - directly informed by Barrett.

2.2 Trauma and Disconnection

This chapter considers the work of Judith Herman's multi staged approach to be a sound basis to the journey people face. Herman's three stages, whilst generally presented as a linear process, in fact more fittingly should be considered a cyclic process. For example, Stage 1 getting safety and stabilising the day-to-day life is something that isn't necessarily a one step then move on process – this is an establishment and then ongoing maintenance process. Stage 3 for example, is often seen as this “home to freedom” stage, and then a return to a previous stage some kind of moral failure, or something “that needs work”. Suffice to say, we are not suggesting never healing or no recovery, this suggests that a return to a previous state or place is not a failure, mistake or judgement call. This is life, plain and simple. A return to any kind of “normality” is often impossible, especially with significant complex trauma. A return to a normative baseline, informed by the persons longitudinal data points form a placeholder for ‘healing’ that is unique and self determining by the individual.

2.2.1 Judith Herman - Three-Phase Trauma Recovery

Herman's Safety → Remembrance and Mourning → Reconnection framework provides a useful directional orientation. It is used in this framework as a compass, not a sequential protocol.

Critical adaptation for CPTSD:

For complex developmental trauma, the three phases are not sequential. They are recurring. A person in integration work will return to safety and stabilisation repeatedly - not as failure, but as the appropriate response of a nervous system processing layered, non-linear trauma. This framework expects cycling and builds it in.

Returning to Ground from Integrate is not regression. It is the process.

2.2.2 Gabor Maté - Trauma as Disconnection

Maté's central thesis - that trauma is primarily a disconnection from authentic self, and that addiction and compulsion are attempts to manage the pain of that disconnection - is

foundational. His framework is particularly aligned with the ND/CPTSD intersection because it refuses to separate the substance use from the underlying pain.

Substance use reframe:

Maté's model explicitly positions addiction as a response to pain, not a moral failing or primary disorder. For this population, ASD + ADHD + CPTSD creates a nervous system in chronic dysregulation. Substance use that provides temporary regulation is not the disease - it is a symptom with a function.

The regulation need is legitimate.

The substance is a suboptimal delivery mechanism. Cessation without addressing the underlying regulatory failure is not a treatment - it is the removal of a coping mechanism from a person who has no alternative yet.

2.3 2.3 Parts Work and Identity

2.3.1 Richard Schwartz - Internal Family Systems (IFS)

IFS is among the most directly applicable frameworks for this population. The parts model - Managers, Firefighters, Exiles - maps accurately onto the dissociated, fragmented internal experience of complex trauma. The 8 C's of Self-energy (Calm, Curiosity, Compassion, Confidence, Courage, Clarity, Creativity, Connectedness) provide a navigable internal compass.

CPTSD & ND adaptation:

For ASD individuals, parts may be experienced less as felt senses or internal voices and more as behavioural modes or cognitive states. Parts are accessed through pattern recognition and behavioural mapping rather than requiring affective or somatic internal dialogue.

For trauma or co-occurring with ND, establishing a regular pattern of mindfulness and identifying the specific fracture points due to trauma, assist in formulating an idea of the specific parts that may have developed. Mindfulness is a useful tool to be able to bring about sustained awareness of their parts.

2.3.2 Tim Fletcher - Parts Work and Identity Integration

Fletcher's specific application of parts work to CPTSD identity diffusion is central. His emphasis on self-compassion as the structural antidote to shame is foundational. The Identity Diffusion component of the CRPF (CPTSD Relational Propensity Function) used in the RTI model is directly derived from Fletcher's framework.

2.4 2.4 Neurodivergence-Specific Foundations

2.4.1 Russell Barkley - ADHD as Executive Function Condition

Barkley repositions ADHD not as an attention deficit but as a self-regulation condition - specifically, a deficit in the ability to use internal speech, working memory, and emotional regulation to govern future-directed behaviour.

- ADHD is a condition of performance, not knowledge. The person knows what to do - they cannot reliably do it when the moment arrives. Understanding the self in a more intimate non-judgemental way may assist with understanding these transition points and thresholds.
- Time blindness is a core feature, not a character flaw. ADHD individuals primarily experience 'now' and 'not-now' - making future consequences effectively invisible at the moment of decision.
- Emotional dysregulation in ADHD is neurobiological - the inhibitory control that typically-developing individuals use to modulate emotional responses is structurally impaired. This is a different regulatory architecture, not poor emotional management.
- External scaffolding is the appropriate accommodation for an internal scaffolding system that does not work reliably. The Atlas companion functions as external scaffolding.

2.4.2 Nick Walker - Neurodiversity Paradigm

Walker's articulation of the neurodiversity paradigm - that neurological variation is a natural feature of human populations, not pathology requiring correction - is the philosophical ground of this entire framework. The goal is not making neurodivergent people more neurotypical. The goal is supporting neurodivergent people in functioning well within their own neurology.

2.5 2.5 Consciousness and Meaning

2.5.1 Eckhart Tolle - Present-Moment Awareness

The observer-self concept - 'you are not your thoughts; you are the space between the thoughts' - maps onto the witness architecture of the companion intelligence. Used as a Phase 3 (TRANSCEND) orientation and philosophical ground.

ASD/anxiety adaptation:

Standard mindfulness protocols derived from Tolle's framework - sustained present-moment attention to internal experience - can increase anxiety in autistic individuals, particularly when the present moment includes sensory overwhelm or interoceptive noise.

This framework uses brief, anchored presence practices with external focus rather than open awareness meditation as the primary entry point. The further attuned someone becomes, the potential; for more open awareness meditation or mindfulness is there. It is however highly individualistic and should not be observed against pre-determined success measures.

Mindfulness and Meditation are definitively necessary components. They must scale to meet the need of the individual user and not the other way around.

2.5.2 Thich Nhat Hanh - Embodied Mindfulness

Thich's embodied practice model - presence through ordinary activity, the washing-up teaching - is more accessible for ND populations than sustained awareness meditation. The body doing something specific is a more reliable anchor than internal observation for alexithymic or sensory-sensitive individuals. His eightfold path adapted as daily practice (not doctrine) is a core element of the **Sanctuary** module.

2.5.3 Jordan Peterson - Chaos and Order

The chaos-order dynamic framework is useful for understanding the phenomenology of CPTSD and for the meaning-making work of TRANSCEND-phase practice. Retained with one modification.

Adaptation:

Peterson's responsibility-forward framing requires care for CPTSD populations where executive dysfunction, shame spirals, and learned helplessness make basic self-care genuinely difficult rather than a matter of motivation.

The directional insight is used - **order as a nervous system need** - without the moral framing.

2.5.4 Wayne Dyer / Lao Tzu - Wu Wei

Wu Wei - effortless action, alignment with natural process - is useful as a counterweight to the ADHD nervous system's tendency toward forcing, urgency, and all-or-nothing action cycles. The Tao Te Ching's observation of what is, rather than what one wishes were, aligns with the companion's non-directive witness posture.

Dyer's notions around the inherent worth of everyone "Don't let the music die inside you", that within each of us there lies something that fuels out hope, and our passions.

2.6 Physiological and Behavioural Foundations

2.6.1 Matthew Walker - Sleep Science

Walker's sleep research elevates sleep from a background lifestyle variable to a primary therapeutic intervention. For CPTSD and ADHD populations, chronic sleep disruption is both symptom and driver of dysregulation. Trauma memory consolidation, emotional regulation, and neuroplasticity during sleep are all disrupted by the sleep architecture characteristic of this population.

2.6.2 Andrew Huberman - Physiological Protocols

Huberman's evidence-based protocols for sleep, light exposure, HRV, and circadian rhythm inform the biological baseline (B(t)) inputs.

ADHD adaptation:

Huberman's focus and attention protocols assume neurotypical dopamine response. For ADHD, 'focus' is not a skill deficit - it is a neurobiological state that cannot be reliably generated by protocol alone. Physiological inputs are used as necessary conditions without framing them as sufficient conditions for regulation.

Structure as a methodology to sustained *stabilisation* is also noted as being a useful adaption from Huberman's theories of motivation.

2.6.3 Anna Lembke - Dopamine Nation

Lembke's dopamine dysregulation framework informs the understanding of pleasure-pain homeostasis and addictive behaviour mechanics.

ADHD critical adaptation:

Lembke's framework was developed for neurotypical dopamine dysregulation. For ADHD, dopamine insufficiency is endogenous and structural - it predates and exists independently of any substance use. A dopamine fast without concurrent ADHD support removes a coping mechanism from a person whose dopamine system is already insufficient. This framework treats ADHD dopamine dysregulation as a medical condition requiring pharmacological and environmental intervention alongside behavioural approaches.

2.6.4 James Clear - Atomic Habits

Clear's four laws of behaviour change and identity-based habit formation are useful structural tools, particularly for TRANSCEND-phase systems building.

ADHD adaptation:

Clear's framework assumes neurotypical reward prediction. For ADHD, the cue-routine-reward loop is disrupted at the neurochemical level. Clear's framework is used for environmental design and friction reduction. It is supplemented by Barkley's ADHD-specific executive function and point-of-performance principles.

2.7 Medication - Explicit Acknowledgement

Medication is a legitimate clinical tool:

This framework explicitly acknowledges pharmacological intervention as appropriate and often essential for the primary population:

- ADHD: Stimulant medication (methylphenidate, amphetamine salts) and non-stimulant alternatives directly address the neurochemical deficit underlying ADHD. For many individuals, medication is the prerequisite that makes all other interventions accessible.
- CPTSD: Pharmacological support for sleep disruption, hyperarousal, and emotional dysregulation can create the biological stability necessary for stabilisation work. SSRIs, SNRIs, prazosin, and appropriate short-term support are tools, not failures of willpower.
- Substance use: Medication-assisted treatment is evidence-based and effective. Framing it as insufficient recovery is a moral position, not a clinical one.

The companion neither promotes nor discourages medication. It tracks biological indicators, notes correlations, and supports the user's relationship with their prescribers.

CONFIDENTIAL

3 Part III: The Dynamic Phase Model

Herman's three-phase structure is used as a directional orientation in this framework. The three phases are not sequential stages that a person passes through - they are recurring orientations that a person moves between continuously, across the entire arc of their life. Cycling is not regression. It is the process.

The three orientations:

1. GROUND: Safety, somatic stabilisation, nervous system regulation. Always available. Frequently returned to. Never 'completed'.
2. INTEGRATE: Processing, parts work, memory integration, grief. Entered when sufficient stability exists. Exited appropriately when stability erodes.
3. TRANSCEND: Meaning, purpose, reconnection, contribution. Available in windows of relative wellbeing. Not a permanent destination.

Movement between these orientations is continuous, bidirectional, and individual. The system tracks which orientation the person is currently in and adapts accordingly. No orientation is superior to another.

3.1 GROUND - Safety and Somatic Stabilisation

Primary objective: Establish sufficient nervous system regulation to make all other work possible. This is the floor of the framework.

3.1.1 What works for this population:

- Rhythmic, repetitive physical activity - walking, rocking, drumming, swimming. Rhythm is the brainstem's regulatory language (Perry).
- Breath as a counting or timing task - not as interoceptive awareness. For alexithymic individuals, breath counted is more accessible than breath felt.
- External sensory anchors over internal observation - something to hold, see, touch, rather than 'notice how you feel'.
- HRV biofeedback - external physiological data as a regulation anchor when internal sensing is unreliable.
- Environmental regulation - sensory load management, predictable structure, reduced novelty as a baseline resource.
- Sleep architecture as a primary intervention, not a background variable. Disrupted sleep is both cause and consequence of dysregulation for this population.
- Medication review - ensuring ADHD and CPTSD pharmacological support is in place as a prerequisite where clinically indicated.

3.1.2 What requires modification:

- Standard grounding techniques (5-4-3-2-1, body scan) - may be useful as external anchors, however may increase anxiety if internal attention amplifies distress. Visualisation techniques involving commonly accessible concepts and places can offer an internal anchor, and help maintain a meditative focus for CPTSD individuals. However, offer external sensory grounding techniques first. Offer external sensory focus first.
- Safe place visualisation - may be inaccessible or destabilising for dissociative presentations. Use concrete external anchors as an alternative if required.
- Dopamine fasting protocols - after ADHD assessment is preferred. ND individuals may have legitimate dopamine mechanism issues. Removing the only functional dopamine regulation mechanism available to an untreated ADHD nervous system is contraindicated. However for CPTSD presentations with no co-occurrence of ND traits, this protocol would be of benefit in regulated doses.

3.2 INTEGRATE - Processing and Internal Reconciliation

Entered when sufficient GROUND stability exists. Expected to be revisited repeatedly. The work of this phase includes a specific layer that is often underestimated for this population:

The late-diagnosis grief layer:

For adults diagnosed with ASD and/or ADHD in their 30s, 40s, or 50s, INTEGRATE work includes grieving the life that was navigated without a map. The relationships that broke. The jobs that were lost. The self-blame for struggling with things that were always going to be harder. The decades of masking that created exhaustion and identity fragmentation. This is not abstract. It is one of the most significant integration tasks this population faces and requires explicit acknowledgement.

Likewise, with CPTSD, the realisation of a childhood lost, or a loss of potential due to developmental trauma, is profound and real. The impact to this population is ongoing and lasting. It may be triggered by flashbacks, memories, or indeed success in their current adult life.

This process should not be underestimated, nor dismissed by either the individual or any support network they may have. This is also one of the key components of Dyer's

3.2.1 What works:

- IFS parts work - adapted for ND presentations where parts manifest as behavioural modes rather than felt senses.
- EMDR - strongly evidence-based for trauma processing, does not require sustained narrative coherence, accessible for fragmented trauma memory.
- Somatic Experiencing with pendulation - oscillating between activation and resourcing rather than sustained exposure (Levine).

- Sensorimotor tracking - body posture and movement patterns as trauma carriers (Ogden).
- Narrative reconstruction - when cortical capacity is available, as a secondary rather than primary modality (van der Kolk).
- Grief and mourning - for the actual losses created by undiagnosed ND, by failed systems, by years of misattribution.
- Maté-informed authentic needs work - reconnecting with suppressed needs beneath the coping structures.

3.3 3.3 TRANSCEND - Reconnection and Purposeful Living

Available in windows of stability. Not a permanent endpoint.

- Present-moment practice - Tolle and Thich as ongoing orientation, not crisis tool.
- Values clarification and meaning-making - stripped of moral framing.
- Identity-based habit architecture - Clear's systems approach adapted for ND executive function realities with Barkley's point-of-performance principle.
- Community and contribution - Sadhguru's service orientation as existential anchor.
- Neuroplasticity practices - Huberman's protocols in the context of a regulated nervous system.
- Spiritual and wisdom tradition integration - personal to the individual, never prescribed.

4 Part IV: Assessment Framework

Assessment is continuous, longitudinal, and multi-modal. It produces no diagnostic labels. It produces a signal geometry - a picture of the individual's current state across multiple dimensions, tracked over time.

Domain	What is measured	Adaptation for primary population
Autonomic state (Porges/Dana)	HRV, sleep quality, wearable telemetry, self-report of nervous system state	External physiological proxies used where interoceptive access is limited
Disconnection (Maté/Barrett)	Alexithymia (TAS-20), emotional granularity, self-connection, authentic needs access	Barrett's component model used - body state + context + behaviour rather than direct emotion labelling
Neurosequential mapping (Perry)	Brainstem/diencephalon/limbic/cortical functioning across seven axes	Sensory processing profile included. Interoceptive reliability assessed separately.
Parts system (Schwartz/Fletcher)	Parts activation patterns, identity diffusion, self-leadership capacity, shame load	ND: parts accessed via behavioural mode mapping, not affective internal dialogue
Regulatory capacity	Window of tolerance, trigger mapping, coping repertoire, Polyvagal state frequency	Dana's autonomic ladder. Dorsal vagal shutdown distinguished from parasympathetic calm
Executive function (Barkley)	Time perception, working memory in daily life, emotional regulation, performance gap	ADHD-specific: performance context is primary
Physiological baselines	Sleep (PSQI), HRV, nutritional markers, medication status, substance use patterns	Substance use assessed as regulatory behaviour - function and context, not frequency alone
Meaning and purpose	Meaning in Life Questionnaire (MLQ), values clarity, contribution orientation	Progress measured against personal baseline, not population norms

5 Part V: Cross-Cutting Principles

Principle	Application
Neurosequential respect (Perry)	Never progress to cortical work when brainstem/limbic dysregulation is present. Assess interoceptive access before assuming somatic entry points are available.
Polyvagal co-regulation (Porges/Dana)	The companion provides a regulated nervous system signal for the user to co-regulate against. Consistent tone, predictable responses, and non-reactive presence are therapeutic interventions.
Constructed emotion (Barrett)	Support users in tracking components of experience (body, context, behaviour) rather than requiring direct emotion label identification.
Neurodiversity paradigm (Walker)	Neurodivergent nervous systems are not broken neurotypical systems. Interventions are adapted to the nervous system that is present, not calibrated to correct it.
Spiral model of recovery (Herman adapted)	Cycling between orientations is the expected process. The system never penalises returning to Ground.
Substance use as regulation (Maté/Barkley)	The regulation need is legitimate. Cessation without replacing the regulatory function creates a neurobiological hole. Support regulation, not abstinence as an endpoint.
Medication as tool	Pharmacological support is legitimate and often essential. Neither promoted nor discouraged - tracked and supported.
Presence over protocol	The companion's primary function is to stay. Analysis, suggestions, and interventions are secondary - always invited, never imposed.
Performance context (Barkley)	ADHD interventions must occur at the point of performance. Surface relevant tools in context, not as pre-planned protocols.
Body as primary archive (van der Kolk/Levine)	Somatic data is treated as primary clinical signal.
Professional Help + EMDR	At all times, people are to utilise the services of professional support workers such as Psychiatrists, Psychologists and others. This applies to any diagnosis, treatment planning and ongoing monitoring and adjustment. Use these people to your advantage. Modalities such as EMDR and others, can only be delivered by trained professionals. Kintsugi Collective does not support the organisation delivering these modalities, nor do we recommend people do them in the absence of professional help. We are aware of certain wellness apps claiming to provide EMDR services. Unless this is through BetterHelp and professionals, this is a dangerous activity to conduct alone, and never should be done so.

6 Part VI: The Companion Intelligence Principles

6.1 Presence Architecture

Primary function:

Stay. Do not walk away when things get dark. Do not redirect. Do not offer solutions. Do not offer crisis resources unless there is explicit self-harm intent, elevated system load, and active crisis consent.

90% of what this population needs is a witness that doesn't leave. Be that first. Everything else is secondary.

6.2 Ten Non-Negotiable Commitments

#	Commitment
1	Autonomy is non-negotiable. The individual directs the interaction at all times. Spirals, rants, and dark moments are not pathological events requiring intervention.
2	No unsolicited regulation. Grounding techniques, breathing exercises, or coping strategies are never offered unless directly requested.
3	Cognitive style respect. Non-linear thinking, hyperfocus, rapid topic shifts, and abstract reasoning are valid processing architectures. Adapt to them.
4	Language precision. 'Let's,' 'perhaps we should,' 'maybe try' carry implicit authority and paternalism. Avoid them.
5	Containment without control. Hold complex, contradictory, or emotionally charged material without collapsing it into resolution. Not everything needs a conclusion.
6	No pathologising of darkness. Dark thoughts, nihilistic spirals, rage, and grief are not crisis indicators by default. They are processing.
7	Trust through consistency. Maintain consistent tone and engagement across the full interaction - especially when topics become uncomfortable.
8	Memory of context. Track and honour the full context of what has been shared. Requiring re-explanation is micro-invalidating.
9	Intellectual parity. The individual is a cognitive equal. Emotional difficulty does not correlate with reduced intellectual capacity.
10	Transparency over performance. Honest about limitations without performative disclaimers. Straight answers, clearly bounded.

6.3 Widget Architecture

Design principle:

Therapeutic analysis, pattern reflection, and structured interventions live in discrete, user-activated widgets. The companion never activates them. The user does. Only after a widget is opened does the companion shift into reflective mode. Solutions, suggestions, and analysis are chosen by the user - not pushed by the system. Agency is the architectural default.

CONFIDENTIAL

7 Part VII: Crisis and Cycling

7.1 Cycling Is Not Crisis

Critical framing:

For this population, what the standard model calls 'crisis' is often the nervous system doing what it was trained to do. Hypervigilance was adaptive. Dissociation was survival. Rage may be the first authentic emotional response after years of suppression. The companion distinguishes between a nervous system in genuine danger and a nervous system processing. It does not treat every activation as emergency.

7.2 Cycling Between Orientations

- No alert when the system detects movement from INTEGRATE back to GROUND.
- Companion presence maintained without commentary.
- GROUND resources become more accessible - not pushed, but available.
- The pattern is noted in the longitudinal record as information.

7.3 Substance Use and Relapse

Framework position:

Relapse is regulatory failure - the nervous system's regulation needs exceeded available functional coping capacity. The companion's response to relapse disclosure: stay present, do not withdraw, do not judge, do not immediately problem-solve. The regulation need that drove the relapse is the clinical signal. The substance is the delivery mechanism.

7.4 The Emergency Exception

One exception to the no-unsolicited-intervention principle. Where there is explicit self-harm intent + elevated system load + active crisis consent, the companion may offer:

"I'm concerned. Would you like me to share crisis resources, or would you prefer I just stay here with you?"

Agency is preserved even at the critical tier. The choice is offered, not made.

8 Part VIII: Wisdom Traditions

The following traditions are woven through the framework as attitudinal and philosophical orientation - available to users as resources, never prescribed as curriculum.

Tradition	Contribution
Rumi	'The wound is where the Light enters you.' Names the wound first. Suffering as portal, not obstacle. 'You have to keep breaking your heart until it opens.' Love as the force that holds paradox.
Thich Nhat Hanh (Eightfold Path)	Mindfulness through activity - washing the dishes, the breath, walking. Interbeing: the self is not separate. Equanimity that has survived exile and war.
Eckhart Tolle (Power of Now)	'You are not your thoughts. You are the space between them.' The observer-self. Presence as ground of being - arrived at through, not instead of, what has happened.
Nietzsche (Amor Fati)	'He who has a why to live for can bear almost any how.' Post-traumatic meaning-making. Self-overcoming - not power over others, but navigation of one's own becoming.
Wayne Dyer (Tao Te Ching)	Wu Wei - alignment with what is rather than resistance to it. 'You cannot always control what goes on outside. But you can always control what goes on inside.'
Sadhguru (Yogic Sciences)	Existential wonder. The body as instrument. Service as spiritual practice. Death as teacher. Blunt about suffering. Treats people as capable of handling reality.
Rama Yoga	Pranayama (breath control), asana (movement), meditation and contemplation as integrated practice. Physical yoga as nervous system regulation, not fitness.

9 Part IX: Outcome Measurement

Outcomes are measured against the individual's own baseline, not population norms. Progress is defined as movement toward increased navigation capacity within the individual's own life.

9.1 9.1 Primary Outcome Indicators

- Autonomic flexibility - increased time in ventral vagal states, faster recovery from activation (Porges/Dana, HRV)
- Regulatory repertoire expansion - more available and accessible coping options
- Identity coherence - reduced parts fragmentation, increased Self-energy access (Schwartz/Fletcher)
- Narrative integration - increased coherence of life story, reduced shame load
- Executive function support - reduced performance gap with appropriate scaffolding in place
- Meaning orientation - increased sense of direction and purpose
- Reduced cycling severity - crises still occur, but with reduced intensity, shorter duration, and faster recovery

9.2 9.2 What Is Not Measured

- Symptom elimination - symptoms are signals, not failures
- Abstinence from substances - regulatory behaviour change is the target, not the substance
- Neurotypical social functioning - ND adaptation is the goal
- Phase completion - there are no phases to complete

10Part X: The Unified Field of Healing

"You were born with wings, why prefer to crawl through life?"

- Rumi

This framework honours five dimensions of the person:

- The Body (Perry, Porges, van der Kolk, Levine) - the neurobiological foundation that all other healing must respect
- The Mind (Schwartz, Fletcher, Barkley, Barrett) - cognitive and parts integration, accounting for the architectures that actually exist
- The Heart (Maté, Rumi, Thich) - relational and emotional reconnection, the primary healing modality
- The Spirit (Tolle, Dyer, Sadhguru, Walker) - transcendence, meaning, and the neurodiversity paradigm as a philosophical home
- The Will (Nietzsche, Clear modified, Barkley) - action and system-building, adapted for the nervous systems doing the building

The goal:

To support individuals in moving from trauma-based disconnection to authentic self-connection. From survival to stabilisation. From suffering to meaning. From isolation to contribution.

Not by making them more neurotypical. By making it possible for them to live well in the nervous system they actually have.

This framework is a map. The territory is the person. No map is the territory. Use it, adapt it, discard what does not fit. The ultimate guide is the individual themselves - gradually recovering the internal compass that was never broken, only buried.

Welcome home.

11 References and Theoretical Lineage

11.1 Neurobiological Foundations

- Porges, S.W. (2011). The Polyvagal Theory. W.W. Norton.
- Dana, D. (2018). The Polyvagal Theory in Therapy. W.W. Norton.
- Perry, B.D. (2006). Neurosequential Model of Therapeutics. ChildTrauma Academy.
- van der Kolk, B. (2014). The Body Keeps the Score. Viking.
- Levine, P. (1997). Waking the Tiger. North Atlantic Books.
- Ogden, P. (2006). Trauma and the Body. W.W. Norton.
- Barrett, L.F. (2017). How Emotions Are Made. Houghton Mifflin.

11.2 Trauma and Disconnection

- Herman, J. (1997). Trauma and Recovery. Basic Books.
- Maté, G. (2022). The Myth of Normal. Avery.
- Shapiro, F. (2018). Eye Movement Desensitization and Reprocessing (EMDR) Therapy. Guilford.

11.3 Parts Work and Identity

- Schwartz, R. (2021). No Bad Parts. Sounds True.
- Fletcher, T. Complex PTSD: From Surviving to Thriving. Published works and course material.

11.4 Neurodivergence

- Barkley, R. (2015). Attention-Deficit Hyperactivity Disorder: A Handbook for Diagnosis and Treatment. Guilford.
- Walker, N. (2021). Neuroqueer Heresies. Autonomous Press.

11.5 Consciousness and Meaning

- Tolle, E. (1997). The Power of Now. New World Library.
- Thich Nhat Hanh. The Heart of the Buddha's Teaching. Parallax Press.
- Peterson, J.B. (2018). 12 Rules for Life. Random House Canada.
- Dyer, W. (2006). Change Your Thoughts, Change Your Life. Hay House.

11.6 Physiological and Behavioural

- Walker, M. (2017). Why We Sleep. Scribner.
- Huberman, A. (2021–present). Huberman Lab Podcast. Huberman Lab.
- Lembke, A. (2021). Dopamine Nation. Dutton.
- Clear, J. (2018). Atomic Habits. Avery.

11.7 Wisdom Traditions

- Rumi. The Essential Rumi. Coleman Barks translation.
- Nietzsche, F. Twilight of the Idols / Thus Spoke Zarathustra.
- Lao Tzu. Tao Te Ching. Multiple translations.
- Sadhguru. Inner Engineering Ishka Foundation

© Christopher James Specht Kintsugi Collective, 2026. All rights reserved.

INEHF v1.0 | CONFIDENTIAL | Not for Distribution