

SUPPLEMENTARY INFORMATION FOR

On the Origin and Spread of Feral Pigeons

George Pacheco^{1✉}, Filipe G. Vieira¹, Michael D. Martin¹, Morten Tange Olsen¹, Pavel Hulva¹, Tânia de Freitas Raso¹, Peter Njoroge¹, Concepción Salaberria¹, Isabel López-Rull¹, Carles Lalueza-Fox¹, Oscar Ramírez¹, María C. Ávila-Arcos¹, Patricia Rosas Escobar¹, Rui Faria¹, Miguel Carneiro¹, Graciela Sotelo¹, Jóhannis Danielsen¹, Nizar Haddad¹, Fares Khoury¹, Roi Dor¹, Ali Halajian¹, María Belén Arias¹, Oliver Krone¹, Susanne Auls¹, Sampath S. Seneviratne¹, Kajanka Mathiaparanam¹, Michael Bunce¹, Megan L. Coghlan¹, Jon Fjeldsø¹ & M. Thomas P. Gilbert^{1✉}

Supplementary Notes

Sampling effort. To investigate the genomic patterns of current pigeon populations of different evolutionary histories, we intentionally targeted our sampling to cover four distinct categories (Figure 1). Furthermore, to help root the evolutionary relationships between these groups, we also included a small number of individuals representing the *Columba livia* intermedia (Strickland, 1844) subspecies. Specifically in this regard, we sampled five populations from Sri Lanka, where two of them were from urban localities (Colombo and Trincomalee), one was from a Conservation National Park (Pigeon Island) and two others were captive populations maintained by local breeders (Wattala and Wellawatte) (Supplementary Fig.


1). To check for data reproducibility, we sequenced two of the samples twice (Tehran_16-GBS and Perth_02-GBS) to serve as replicates. Finally, to serve as external outgroups, we also generated data from five samples of (*Columba palumbus* Linnaeus, 1758) captured in Copenhagen (Denmark), one captive sample of (*Streptopelia risoria* Linnaeus, 1758), and additionally incorporated previously published whole genome resequence data from a *Columba rupestris*⁸. (we also included the WGS library to serve as another replicate) (Supplementary Spreadsheet).

Methods


¹Section for Evolutionary Genomics, The GLOBE Institute, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark. ¹Natural History Museum of Denmark, University of Copenhagen, Øster Voldgade 5–7, 1350 Copenhagen, Denmark. ¹NTNU University Museum, Norwegian University of Science and Technology, Trondheim, Norway ¹Department of Zoology, Charles University, Prague, Czech Republic. ¹Departamento de Patologia, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, São Paulo, Brazil. ¹Ornithology Section, Department of Zoology, National Museums of Kenya, Nairobi, Kenya. ¹Centro de Investigación en Ecosistemas, Universidad Nacional Autónoma de México, Michoacán, México. ¹Departamento de Ecología Evolutiva, Museo Nacional de Ciencias Naturales, Madrid, Spain. ¹Avian Evolution Node, Department of Zoology and Environment Sciences, University of Colombo, Colombo, Sri Lanka. ¹Institute of Evolutionary Biology, Universitat Pompeu Fabra, Barcelona, Spain. ¹Department of Animal and Plant Sciences, University of Sheffield, Sheffield, UK. ¹Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, Vairão, Portugal. ¹Institute of Evolutionary Biology, Department of Experimental and Health Sciences, University, Pompeu Fabra, Spain. ¹Departamento de Biología, Faculdade de Ciências, Universidade do Porto, Porto, Portugal. ¹University of the Faroe Islands, Tórshavn, Faroe Islands. ¹National Center for Agricultural Research and Extension, Al-Baqah, Jordan. ¹Department of Biology and Biotechnology, American University of Madaba, Madaba, Jordan. ¹Department of Zoology, Tel Aviv University, Tel Aviv, Israel. ¹Natural History Museum, Imperial College of London, London, United Kingdom. ¹Department of Biodiversity, Turfloop Campus, University of Limpopo, Polokwane, South Africa. ¹Department of Wildlife Diseases, Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany. ¹Vetgenomics SL, Edifici Eureka, Campus UAB, Barcelona, Spain. ¹Trace and Environmental DNA (TyEnD) Laboratory, Department of Environment and Agriculture, Curtin University, Perth, Australia.

✉Correspondence should be addressed to ganpa@aquas.dtu.dk (G.P.) & tgilbert@sund.ku.dk (M.T.P.G.)

References



Supplementary Fig. 1. Coverage heatmap. Heatmap based on the presence/absence matrix. Individual samples are represented by each column, whereas clusters of loci are represented by the rows.



Supplementary Fig. 1. Global depth distribution. Heatmap based on the presence/absence matrix. Individual samples are represented by each column, whereas clusters of loci are represented by the rows.




Fig. 3 Estimation of Admixture proportions. Individuals are represented by columns, while rows depict the Admixture proportions based on the assumption of different numbers of ancestral populations ($K = 2 - 15$). Individuals are sorted by sampling locality (light grey upper labels) and grouped per biological status




Fig. 2 Multidimensional scaling analysis. a) Dimensions 1 and 3 are plotted. b) Dimensions 2 and 3 are plotted. Each point on the plot represents a single individual. The ellipses represent the rough distribution of the two most distinct groups.