



Erratum: Development of Pyridine-Boron Trifluoride Electrolyte Additives for Lithium-Ion Batteries [*J. Electrochem. Soc.*, 162, A1186 (2015)]

Mengyun Nie, Jian Xia, and J. R. Dahn

Department of Physics and Atmospheric Science, Dalhousie University, Halifax, Nova Scotia B3H4R2, Canada

© 2015 The Electrochemical Society. [DOI: [10.1149/2.0241508jes](https://doi.org/10.1149/2.0241508jes)] All rights reserved. Published May 6, 2015.

On page A1188, left column, the second paragraph under the Results and Discussion heading should be

Figure 3 plots the capacity of the plateau at 2.3 V from Figure 2a versus the concentration of PBF in the electrolyte. Figure 3 shows a linear relationship. If all the PBF added to the cell is reduced during the first charge, then the slope of the line in Figure 3 yields a 1.52 electron reaction. More likely is that the reaction is a two-electron reaction and that the PBF in portions of the electrolyte outside the jelly roll is not reduced. The linearity in Figure 3 suggests that PBF does not passivate the graphite electrode and, instead, passivation occurs in the following EC reduction. Further research is required to determine the mechanism of the apparent two-electron reduction of PBF.

On page A1188, right column, Figure 3 should be




Figure 3. Capacity of the plateau at 2.3 V vs. the concentration of PBF added to the electrolyte.